

THE INDIANA PREVENTION RESOURCE CENTER
2011 PREVALENCE STATISTICS
MAIN FINDINGS

ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS



THE INDIANA PREVENTION RESOURCE CENTER
2011 PREVALENCE STATISTICS
MAIN FINDINGS

ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS

Survey Conducted February through April 2011
Report Dated August 2011

Conducted by:
Indiana Prevention Resource Center

Published by:
Institute for Drug Abuse Prevention

Department of Applied Health Science
School of Health, Physical Education, and Recreation
Indiana University

Contact:
Indiana Prevention Resource Center
501 North Morton Street Suite 110
Bloomington, IN 47404

<http://www.drugs.indiana.edu>
drugprc@indiana.edu
Toll Free: 1-800-346-3077
Fax: 812-855-4940

Funded, in part by a contract with the
Indiana Family and Social Services Administration
Division of Mental Health and Addiction, and the
School of Health, Physical Education, and Recreation

Source Citation:

Gassman R., Jun, M. K., Samuel, S., Agley, J. D., Lee, J., Morrison, A. M., Agley, B. D., Carey, J. M., Karim, A. S., Morales, N., Oi, S. E., Pardue, N. L., Pardue, S. E., Quiroz, J. E., Shadley, B. G., Smith, J. N., Smith, M. D., Subramanian, R., & Zvolner, L. E. (2011). Alcohol, Tobacco, and Other Drug Use by Indiana Children and Adolescents: The Indiana Prevention Resource Center Survey – 2011 (IDAP Monograph No. 11-01). Bloomington, IN: Indiana Prevention Resource Center.

Table of Contents

| | |
|------------------------------------|----|
| I. Report | 1 |
| Introduction | 2 |
| Results | 8 |
| Methodology | 38 |
| How to Interpret the Results | 52 |
| References | 56 |

| | |
|---------------------------------------|----|
| II. Statewide Prevalence of Use | 61 |
|---------------------------------------|----|

| | Grade | 6 th | 7 th | 8 th | 9 th | 10 th | 11 th | 12 th |
|---|-------|-----------------|-----------------|-----------------|-----------------|------------------|------------------|------------------|
| Percentage Reporting Lifetime Use of ATOD | | 62 | 64 | 66 | 68 | 70 | 72 | 74 |
| Percentage Reporting Monthly Use of ATOD | | 63 | 65 | 67 | 69 | 71 | 73 | 75 |
| Mean Age of First Time Use of Alcohol, Tobacco, and Other Drugs | | | | | | | | 76 |

| | |
|--|----|
| III. Prevalence of Use by Race/Ethnicity | 77 |
|--|----|

| | Grade | 6 th | 7 th | 8 th | 9 th | 10 th | 11 th | 12 th |
|---|-------|-----------------|-----------------|-----------------|-----------------|------------------|------------------|------------------|
| Percentage Reporting Lifetime Use of ATOD | | 78 | 80 | 82 | 84 | 86 | 88 | 90 |
| Percentage Reporting Monthly Use of ATOD | | 79 | 81 | 83 | 85 | 87 | 89 | 91 |

| | |
|---------------------------------------|----|
| IV. Prevalence of Use by Gender | 93 |
|---------------------------------------|----|

| | Grade | 6 th | 7 th | 8 th | 9 th | 10 th | 11 th | 12 th |
|---|-------|-----------------|-----------------|-----------------|-----------------|------------------|------------------|------------------|
| Percentage Reporting Lifetime Use of ATOD | | 94 | 95 | 96 | 97 | 98 | 99 | 100 |
| Percentage Reporting Monthly Use of ATOD | | 94 | 95 | 96 | 97 | 98 | 99 | 100 |

V. Prevalence of Use by Region 101

| | Grade | 6 th | 7 th | 8 th | 9 th | 10 th | 11 th | 12 th |
|---|-------|-----------------|-----------------|-----------------|-----------------|------------------|------------------|------------------|
| Percentage Reporting Lifetime Use of ATOD | 102 | 104 | 106 | 108 | 110 | 112 | 114 | |
| Percentage Reporting Monthly Use of ATOD | 103 | 105 | 107 | 109 | 111 | 113 | 115 | |

VI. Prevalence of Use by DSA 117

| | Grade | 6 th | 7 th | 8 th | 9 th | 10 th | 11 th | 12 th |
|---|-------|-----------------|-----------------|-----------------|-----------------|------------------|------------------|------------------|
| Percentage Reporting Lifetime Use of ATOD | 118 | 120 | 122 | 124 | 126 | 128 | 130 | |
| Percentage Reporting Monthly Use of ATOD | 119 | 121 | 123 | 125 | 127 | 129 | 131 | |

VII. Trends and Prevalence Graphs 133

Trends Graphs 134

Prevalence Graphs 148

VIII. Frequency Tables 159

General 160

Male 194

Female 228

Appendix: Instrument 263

Data tables and graphs can be downloaded from the IPRC website:
<http://www.drugs.indiana.edu/IndianaSurvey>



I. REPORT

| | |
|--------------------------------------|----|
| • Introduction | 2 |
| • Results | 8 |
| • Methodology | 38 |
| • How to Interpret the Results | 52 |
| • References | 56 |

INTRODUCTION

Overview

In spring, 2011, the Indiana Prevention Resource Center (IPRC) conducted the 21st Annual Survey of Alcohol, Tobacco, and Other Drug Use by Indiana Children and Adolescents, referred to hereafter as the Indiana Survey. The IPRC has conducted the Indiana Survey, and has managed and reported the resulting data, since 1991. This project is administered through a contract with the Division of Mental Health and Addiction (DMHA) of the Indiana Family and Social Services Administration (FSSA). The purpose of this project is to provide data for state and local planning with respect to the use of alcohol, tobacco, and other drugs (ATOD), gambling behaviors, and risk and protective factors.

In February, March, and April, 2011, local school officials administered surveys to students in Grades 6 through 12 in 478 schools throughout Indiana. A total of 168,801 youth from both public and nonpublic schools completed a written survey that asked about their use of various drugs, their age of first use of various drugs, and risk and protective factors. This process resulted in 152,678 usable surveys, the primary source of the data reported here. The Methodology section describes this process in more detail. The instrument is located in the appendix.

This report summarizes the findings of the 2011 Indiana Survey. These findings include statistics and other information about the prevalence of ATOD use, gambling behaviors, and risk and protective factors. Statewide prevalence-of-use data are shown for previous years to help distinguish trends and facilitate comparisons among years. National prevalence estimates, reported by Monitoring the Future (MTF) (Johnston, O'Malley, Bachman, & Schulenberg, 2011) and the US Centers for Disease Control and Prevention (CDC, 2009), are provided to facilitate comparison with the United States as a whole. For a summary of findings, see the Results section.

Other surveys, such as the Youth Risk Behavior survey (YRBS) (CDC, 2009) and MTF, collect representative state-level data for Indiana. However, the results of these data are not reported for sub-state regions, nor are local results available to local participants. The strength of the Indiana Survey is its ability to describe reported ATOD use at the local level. This reporting facilitates local needs assessment, planning, and evaluation of drug abuse prevention activities. The IPRC provides a report of local results to each participating school corporation free of charge. Data from an individual school corporation is the property of that corporation. Local data may neither be released nor obtained without the written permission of the corporation where the survey was conducted.

National Outcome Measures (NOMs)

The Government Performance and Results Act of 1993 requires government agencies to report the results of their activities. In response, the United States Department of Health and Human Services' Substance Abuse and Mental Health Services Administration (SAMHSA) designated a set of measures, called the National Outcome Measures (NOMs), for substance abuse prevention (Substance Abuse and Mental Health Services Administration [SAMHSA], 2008). SAMHSA collects the NOMs through its National Survey on Drug Use and Health (NSDUH) for all states, and it reports these data for states and sub-state regions (SAMHSA, 2008). The Indiana Survey collects data on the majority of the substance abuse prevention NOMs for youth, and the IPRC also reports that data for Indiana as a whole and for the sub-state regions.

Increasingly, federal and state agencies require the NOMs for needs assessments and evaluations of local prevention programs that they fund. To compete for grants and to measure outcomes without developing new surveillance capacity, communities will need to be able to capture, track, and analyze the NOMs at the local level.

A unique benefit of participating in the Indiana Survey is that participants receive NOMs data on their own schools and communities. The Indiana Survey collects NOMs data on alcohol, tobacco, marijuana and a variety of other illicit drugs for the following measures:

- thirty day prevalence of use,
- perceived risk of harm,
- age of first use, and
- perception of peer disapproval.

Communities That Care (CTC)

Since 2005, Indiana has received funding through the Strategic Prevention Framework State Incentive Grant (SPF-SIG) to expand its prevention infrastructure and decrease substance use and related consequences. A total of 20 communities were sub-recipients of these funds. With the end of this grant funding in sight, the State has adopted an evidence-based planning model, called Communities That Care (CTC), to facilitate further expansion of the Strategic Prevention Framework (SPF) across Indiana's fourteen Defined Service Areas (DSAs). The CTC complements the SPF through its focus on risk and protective factors among youth and adolescents, and its promotion of data-driven decision-making and evaluation. This year, for the second time, the item content in the Indiana Survey has been modified to align

with the essential risk & protective factors from the CTC Survey. The 2011 Indiana Survey results will provide DSAs with additional data needed to assess prevention needs and to identify appropriate evidence-based programs, policies, and practices.

Changes to the Survey Instrument

Where applicable, the category of influence that an item belongs to according to CTC is identified in brackets [] in the right margin.

Changes to Existing Items

The following items were modified in the 2011 survey. Words added this year appear in *italics*.

■ How easy would it be for you to get...

Cigarettes?

Beer, wine or liquor (*for example, vodka, whiskey or gin*)?

Marijuana?

A drug like cocaine, LSD or amphetamines?

- Very difficult
- Fairly difficult
- Fairly easy
- Easy

■ During the past year, did you attend or participate in an “*Afternoons R.O.C.K. in Indiana*” *after-school prevention program, L.E.A.D. Initiative, SADD or other prevention group at your school?*

- Yes
- No

New Items

We added the following items to the 2011 survey. Response choices follow the item in a bulleted list.

■ The next few questions ask about your family and friends. [Family]

My parents ask me what I think before most family decisions affecting me are made.

If I had a personal problem, I could ask my mom or dad for help.

My parents give me lots of chances to do fun things with them.

- YES!
- yes
- no
- NO!

- Think of your four best friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have... [Peer-Individual]

Been members of a gang?

Participated in clubs, organization, or activities at school?

Made a commitment to stay drug-free?

Liked school?

Regularly attended religious services?

Tried to do well in school?

- None of my friends
- 1 of my friends
- 2 of my friends
- 3 of my friends
- 4 of my friends

- Please answer the following questions: [School]

In my school, students have lots of chances to help decide things like class activities and rules.

Teachers ask me to work on special classroom projects.

My teacher(s) notices when I am doing a good job and lets me know about it.

There are lots of chances for students in my school to get involved in sports, clubs, and other school activities outside of class.

There are lots of chances for students in my school to talk with a teacher one-on-one.

I feel safe at my school.

The school lets my parents know when I have done something well.

My teachers praise me when I work hard in school.

I have lots of chances to be part of class discussions or activities.

- YES!
- yes
- no
- NO!

- These questions ask about the neighborhood and community where you live. [Community]

My neighbors notice when I am doing a good job and let me know.

There are people in my neighborhood who are proud of me when I do something well.

There are people in my neighborhood who encourage me to do my best.

- YES!
- yes
- no
- NO!

Regional Reporting

This year, for the fifth time, this report presents prevalence data broken out at the level of FSSA's sub-state planning regions. These eight regions are aggregates of the State's 92 counties, as indicated in Table 1. Prevalence data for sub-state planning regions provides more targeted information than statewide data. Regional results are detailed in the Results section.

Table 1. *Indiana Family and Social Services Administration Planning Regions*

| Sub-state Regions | Counties |
|-------------------|---|
| Central | Boone, Hamilton, Hancock, Hendricks, Johnson, Marion, Morgan, Shelby |
| East | Blackford, Delaware, Fayette, Grant, Henry, Jay, Madison, Randolph, Rush, Union, Wayne |
| North Central | Cass, Elkhart, Fulton, Howard, Kosciusko, La Porte, Marshall, Miami, St. Joseph, Tipton, Wabash |
| Northeast | Adams, Allen, De Kalb, Huntington, LaGrange, Noble, Steuben, Wells, Whitley |
| Northwest | Jasper, Lake, Newton, Porter, Pulaski, Starke |
| West | Benton, Carroll, Clay, Clinton, Fountain, Monroe, Montgomery, Owen, Parke, Putnam, Sullivan, Tippecanoe, Vermillion, Vigo, Warren, White |
| Southeast | Brown, Bartholomew, Decatur, Franklin, Lawrence, Jackson, Jennings, Ripley, Dearborn, Orange, Washington, Scott, Jefferson, Ohio, Switzerland, Crawford, Clark, Harrison, Floyd |
| Southwest | Greene, Knox, Daviess, Martin, Gibson, Pike, Dubois, Posey, Vanderburgh, Warrick, Spencer, Perry |

Defined Service Area Reporting

This year, for the second time, this report presents prevalence data broken out at the level of defined service areas (DSAs). The DSAs are predefined geographic areas with a minimum of approximately 10,000 targeted youth, consisting of one or more counties in Indiana, and served by a single “Local Prevention Services Coalition.” These fourteen areas are aggregates of the State’s 92 counties, as indicated in Table 2. DSAs will continue to be the units for direct service implementation of Afternoons R.O.C.K. in Indiana afterschool programs. In 2010, DMHA initiated the expansion of the SPF and CTC at the DSA level. Prevalence, risk and protective factor data for DSAs provide more targeted information than statewide or regional data. DSA results are detailed in the Results section.

Table 2. *Defined Service Areas*

| DSA | Counties |
|-----|---|
| 1 | Lake |
| 2 | Benton, Carroll, Cass, Jasper, La Porte, Newton, Porter, Pulaski, Starke, White |
| 3 | Fulton, Kosciusko, Marshall, St. Joseph |
| 4 | De Kalb, Elkhart, LaGrange, Noble, Steuben |
| 5 | Allen, Whitley |
| 6 | Boone, Clay, Fountain, Montgomery, Parke, Tippecanoe, Vermillion, Vigo, Warren |
| 7 | Adams, Blackford, Delaware, Grant, Huntington, Jay, Miami, Randolph, Wabash, Wells |
| 8 | Clinton, Hamilton, Howard, Tipton |
| 9 | Marion |
| 10 | Hendricks, Johnson, Morgan, Owen, Putnam |
| 11 | Fayette, Franklin, Hancock, Henry, Madison, Rush, Shelby, Union, Wayne |
| 12 | Bartholomew, Brown, Dearborn, Decatur, Jackson, Jennings, Lawrence, Martin, Monroe, Ohio, Ripley |
| 13 | Daviess, Gibson, Greene, Knox, Pike, Posey, Sullivan, Vanderburgh, Warrick |
| 14 | Clark, Crawford, Dubois, Floyd, Harrison, Jefferson, Orange, Perry, Scott, Spencer, Switzerland, Washington |

RESULTS

This section of the monograph summarizes

- statistically significant changes in statewide prevalence measures from 2010 to 2011
- regional findings
- Defined Service Area findings
- important patterns relating risk and protective factors to substance use, and
- consequences of use

Alcohol, Tobacco, and Other Drugs

Statewide Prevalence Rates

For many measures and grades, the Indiana Survey results yielded no statistically significant differences between this year and last year. Because this section focuses on changes, prevalence rates for certain grades and drugs may not be mentioned in this section. Complete results from the Indiana Survey appear in subsequent sections of this report.

Gateway Drugs

Research has shown that the majority of individuals' illicit drug use occurs only after they use cigarettes, alcohol, or marijuana. These three substances are known as the "gateway drugs" (Bailey, 1992; Donovan & Jessor, 1983; Fleming, Leventhal, Glynn, & Ershler, 1989; Golub & Johnson, 2001; Kandel & Yamaguchi, 1993).

Gateway Drug Use in Grades 6 to 8

Tobacco use in Grades 6 to 8

The lifetime use of cigarettes decreased among youth in Grades 6 to 8. Monthly use of cigarettes decreased among youth in Grades 6 and 8, and held steady among youth in Grade 7. While this year's prevalence rates of cigarette use may be compared with last year's rates, they should not be compared with those previous to 2010. Last year, the response options to the cigarette item were modified to give respondents the option of having smoked at least 1 time (i.e. "1-5 times"), whereas previously the lowest level of smoking possible to select was "a few times."

The lifetime use of cigars decreased for youth in Grades 6 and 8, and held steady for youth in Grade 7, while monthly use of cigars decreased among youth in Grades 6 to

8. Lifetime use of smokeless tobacco decreased among youth in Grades 6 and 8, and monthly use of smokeless tobacco decreased among students in Grade 6. However, lifetime use of tobacco in pipes increased among students in Grade 7.

Alcohol use in Grades 6 to 8

The reported lifetime and monthly use of alcohol among youth in Grades 6 to 8 decreased compared to last year's data. These changes continue a downward trend in lifetime use of alcohol since the early 1990s.

The reported prevalence of binge drinking decreased among students in Grade 8 and held steady among students in Grades 6 and 7.

Marijuana use in Grades 6 to 8

Reported lifetime use of marijuana increased for youth in Grades 6 and 7, and decreased for youth in Grade 8. In addition, the past-month prevalence rates for marijuana use held steady for youth in Grades 6 and 7, and decreased for youth in Grade 8. Among youth in Grade 8, this change marks the end of an upward trend in monthly use of marijuana that occurred between 2008 and 2010 (see Figure 1). However, a non-significant increase among youth in Grade 7 continues a significant upward trend in monthly marijuana use (see Figure 1). The prevalence rate for past 30-day marijuana use among Indiana's eighth-grade youth is higher than the 2010 national prevalence rate (Johnston, O'Malley, Bachman, & Schulenberg, 2011).




Figure 1. Monthly use of marijuana by Indiana 6th through 8th graders

Note. * Statistically significant (R-square > .80).

Gateway Drug Use in Grades 9 to 12

Tobacco use in Grades 9 to 12

Reported lifetime use of cigarettes among youth in Grades 9 to 12 decreased in a manner consistent with the trend observed in previous years. Reported monthly use of cigarettes among youth in Grades 9 to 11 decreased, but the monthly prevalence rate held steady among youth in Grade 12.

Lifetime use of cigars decreased for youth in Grades 10 and 11, and monthly use decreased for youth in Grade 10. Lifetime and monthly use of tobacco in pipes held steady for youth in Grades 9 to 12, except for a decrease in monthly pipe use among youth in Grade 9.

In addition, lifetime smokeless tobacco use increased for youth in Grade 12, and held steady for youth in Grades 9 to 11. Monthly use of smokeless tobacco decreased among youth in Grade 9, and held steady for youth in Grades 10 to 12. These changes, which include a non-significant increase in monthly use among youth in Grade 12, mark an upward trend in lifetime and monthly use of smokeless tobacco among 12th graders observed since 2007 (see Figure 2).




Figure 2. Monthly and lifetime use of smokeless tobacco by Indiana 12th graders

Note. * Statistically significant (R-square > .80).

Alcohol use in Grades 9 to 12

The reported lifetime prevalence of alcohol use among students in Grades 9 to 12 decreased, as did monthly alcohol use among youth in Grades 10 and 12. These changes continue a downward trend in lifetime use of alcohol since the early 1990s.

The reported prevalence of binge drinking held steady for students in Grades 9 to 12.

Marijuana use in Grades 9 to 12

The reported prevalence of lifetime and past-month use of marijuana held steady among youth in Grades 9 to 12. While these rates are not significantly different from those reported last year, they represent a continuation of the increasing trend of monthly marijuana use that was observed beginning in 2008 (see Figure 3). Prevalence rates of past 30-day marijuana use among Indiana's tenth and twelfth-grade youth are lower than the 2010 national prevalence rates (Johnston, O'Malley, Bachman, & Schulenberg, 2011).




Figure 3. Monthly use of marijuana by Indiana 9th through 12th graders

Note. * Statistically significant (R-square > .80).

Methamphetamine

This is the seventh year that the Indiana Survey included questions about methamphetamine.

Compared to last year, the reported prevalence rates of lifetime use of methamphetamine held steady for youth in Grades 6 to 12, and monthly use increased only for youth in Grade 12 (see Table 3). Prevalence rates of monthly methamphetamine use among Indiana's eighth, tenth, and twelfth-grade youth are higher than 2010 national prevalence rates reported in Monitoring the Future (Johnston, O'Malley, Bachman, & Schulenberg, 2011).

Table 3. *Methamphetamine use by youth in grades 6 to 12*

| Grade | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Nation 2010 [#] |
|----------|------|------|------|------|------|------|--------------------------|
| 6 | | | | | | | |
| Lifetime | 0.5 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | |
| Monthly | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | |
| 7 | | | | | | | |
| Lifetime | 1.2 | 1.0 | 0.9 | 0.9 | 0.9 | 1.0 | |
| Monthly | 0.6 | 0.5 | 0.4 | 0.5 | 0.5 | 0.5 | |
| 8 | | | | | | | |
| Lifetime | 2.0 | 1.6 | 1.5 | 1.5 | 1.5 | 1.5 | 1.8 |
| Monthly | 0.9 | 0.7 | 0.7 | 0.6 | 0.8 | 0.9 | 0.7 |
| 9 | | | | | | | |
| Lifetime | 3.0 | 2.2 | 2.0 | 2.1 | 2.0 | 1.9 | |
| Monthly | 1.2 | 0.8 | 0.9 | 0.8 | 0.9 | 0.9 | |
| 10 | | | | | | | |
| Lifetime | 3.5 | 3.0 | 2.5 | 2.3 | 2.2 | 2.2 | 2.5 |
| Monthly | 1.2 | 1.0 | 1.0 | 0.9 | 1.0 | 0.9 | 0.7 |
| 11 | | | | | | | |
| Lifetime | 4.2 | 3.3 | 2.8 | 2.6 | 2.6 | 2.7 | |
| Monthly | 1.5 | 1.1 | 0.9 | 0.9 | 1.2 | 1.3 | |
| 12 | | | | | | | |
| Lifetime | 5.0 | 3.4 | 2.7 | 2.7 | 2.5 | 2.8 | 2.3 |
| Monthly | 1.5 | 1.0 | 0.9 | 0.9 | 1.0 | 1.3* | 0.5 |

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

[#] Johnston, O'Malley, Bachman, & Schulenberg, 2011.

Other Drugs

In the 2008 Survey, the word "Adderall" was added to the item that had previously contained only "Ritalin," so that the item would measure the use of either Ritalin or Adderall. In the 2010 Survey, this item was changed to "Prescription drugs (such as Ritalin, Adderall, Xanax) to get high." This wording was adopted from the CTC survey. Compared to 2010, the lifetime prevalence of prescription drug use to get

high decreased for youth in Grades 8 to 11, and monthly use decreased for youth in Grades 8, 10, and 11. Because of the change to the survey instrument, these rates are not comparable with those observed prior to 2010.

The reported lifetime use of over-the-counter drugs to get high decreased for youth in Grades 6 and 8 to 10. Because the item was modified in 2010 to include the wording "...to get high," in place of the wording "non-medical use," the 2011 rates are not comparable with those observed prior to 2010.

The lifetime and monthly prevalence rates of hallucinogen drug use increased compared to last year for Grade 12. However, this finding should be interpreted with caution. This item was consolidated in the 2010 Survey to "Hallucinogens (LSD, PCP)," from two items in previous year's surveys: "LSD (acid)" and "Other Psychedelics (psilocybin, mescaline, etc.)." Therefore, prevalence rates obtained this year (2011) are not comparable with those observed prior to 2010.

Monthly use of several other substances, including cocaine, crack, and amphetamines, held steady among youth across all grade levels.

The reported lifetime and monthly prevalence rates of heroin use held steady among youth across all grade levels. However, the prevalence of monthly heroin use among Indiana's twelfth-grade youth is three times higher than the 2010 national prevalence rate reported in *Monitoring the Future* (Johnston, O'Malley, Bachman, & Schulenberg, 2011).

The reported lifetime and monthly prevalence of ecstasy decreased or held steady among youth in all grade levels. This item was modified to read "Ecstasy (MDMA, X, XTC)," whereas in previous years it appeared as "MDMA (ecstasy, XTC, X)." As such, the 2011 rates are not comparable with those observed prior to 2010.

The lifetime and monthly prevalence of tranquilizer use held steady or decreased among students in all grade levels. This item was modified to read "Tranquilizers (downers)" in 2010, whereas it previously appeared as "Tranquilizers or Sleeping Pills (downers) (non-prescribed)," thus the 2011 rates are not comparable with those observed prior to 2010.

Lifetime and monthly prevalence of inhalant use decreased or held steady among youth in all grade levels. Among youth in Grades 6 to 8, these rates represent a continuation of a downward trend observed for at least a decade.

This was the second year that the survey contained an item about "Prescription painkillers (Vicodin, OxyContin, Percocet) to get high." Lifetime and monthly prevalence of prescription painkillers decreased or held steady among youth in all grade levels. This new item is not comparable to the item about "narcotics (opium,

morphine, codeine, oxycontin) (non-prescribed)” used in years prior to 2010.

Race and Ethnicity

We report past 30-day prevalence rates for selected drugs for three race/ethnicity categories: Non-Hispanic White, Non-Hispanic Black or African-American, and Hispanic. We provide national data for comparison. Complete prevalence rates for each of these race/ethnicity categories by grade of respondent are reported in the tables section of this monograph.

For this section of the report, we used the Youth Risk Behavior Survey (YRBS) for national comparison data. The national survey to which we usually compare prevalence measures, Monitoring the Future, was less appropriate for comparison since it provides only approximate weights for race/ethnicity subgroups; it reports data only for non-Hispanic Whites, non-Hispanic Blacks, and Hispanics, and it provides only two-year averages—rather than single-year estimates—for prevalence measures. The YRBS dataset is readily available (CDC, 2009), allowing for the calculation of the statistical significance of differences. The YRBS uses a race/ethnicity categorization similar to ours. Though YRBS questions are worded differently from ours, we chose to report in this section a concept (any use in the past 30 days) that is common to both instruments.

Table 4 shows the average prevalence of past 30-day use for respondents in Grades 9 to 12 by race/ethnicity category. YRBS national prevalence rates are provided for comparison. Statistically significant differences between Indiana prevalence rates and national prevalence rates ($p < .05$) are denoted with an asterisk (*). The substances shown were those common to both surveys for past 30-day use.

The following statistically significant differences may be observed: Prevalence of use for Indiana Whites was lower than that of US Whites for all of these drugs. Indiana Blacks’ prevalence of use was lower than US Blacks’ for alcohol and cigars, but higher for binge drinking. Indiana Hispanics’ prevalence of use was lower than US Hispanics’ for alcohol and cigars, but higher for smokeless tobacco. These comparisons are generally, but not entirely, consistent with the comparisons made in last year’s monograph. What differs from last year is that Indiana Blacks’ prevalence rates of cigarette and cocaine use are no longer significantly higher than rates for US Blacks. In addition, Indiana Hispanics’ prevalence rates of marijuana use is no longer significantly lower than the rate for US Hispanics. It is notable that the Indiana Survey and the YRBS found very different prevalence rates for alcohol use across all race/ethnic groups. While we cannot explain this difference, it has appeared in the data from the two studies for many years.

Table 4. *Prevalence of past 30-day use of substances by race/ethnicity, students in Grades 9 to 12, Indiana Survey and United States (YRBS 2009)*

| Substance | IN | US | Substance | IN | US |
|--------------------|------|------|-------------------|------|------|
| Cigarettes* | 18.8 | 19.5 | Alcohol* | 29.9 | 41.8 |
| White* | 18.9 | 22.5 | White* | 29.4 | 44.7 |
| Black | 10.9 | 9.5 | Black* | 28.2 | 33.4 |
| Hispanic | 19.5 | 18.0 | Hispanic* | 32.8 | 42.9 |
| Smokeless tobacco* | 8.3 | 8.9 | Binge drinking ** | 19.7 | 24.2 |
| White* | 8.7 | 11.9 | White* | 18.7 | 27.8 |
| Black | 2.5 | 3.3 | Black* | 19.1 | 13.7 |
| Hispanic* | 6.4 | 5.1 | Hispanic | 25.1 | 24.1 |
| Cigars* | 10.2 | 14.0 | Marijuana* | 16.7 | 20.8 |
| White* | 10.1 | 14.9 | White* | 14.9 | 20.7 |
| Black* | 7.7 | 12.8 | Black | 22.9 | 22.2 |
| Hispanic* | 10.1 | 12.7 | Hispanic | 21.3 | 21.6 |
| Cocaine* | 1.7 | 2.8 | | | |
| White* | 1.2 | 2.4 | | | |
| Black | 2.4 | 1.9 | | | |
| Hispanic | 3.9 | 4.3 | | | |

Notes.

IN = Indiana Survey by Indiana Prevention Resource Center, 2011.

US =Youth Risk Behavior Survey (YRBS) by CDC, 2009.

* $p < .05$

Binge drinking was measured during the past 2 weeks for the Indiana Survey.

Gender

This year, for the fourth time, we present an analysis of differences in substance use prevalence between males and females. For this analysis, we calculated the difference between male and female past 30-day prevalence rates for each drug (see Table 5). We analyzed differences that were statistically significant ($p < .05$). An additional table (see Table 6) provides all prevalence periods for alcohol.

For most drugs and across all grades surveyed, the majority of users were male, and the male majority was usually larger in each successive grade.

However, six drugs had majority-female use in one or more grades in one or more prevalence periods. These drugs were: cigarettes, over-the-counter drugs, prescription drugs (such as Ritalin, Adderall, Xanax), inhalants, prescription painkillers (Vicodin, OxyContin, Percocet), and alcohol. These differences were observed only between Grade 7 and Grade 10, and the extent to which the majority of users statistically are female typically peaks in Grade 8.

Reported prevalence rates for alcohol use for binge drinking (past 2 weeks), past month, and lifetime generally were mixed between male and female majorities, with no clear pattern emerging.

Table 5. *Significant differences ($p < .05$) between male and female prevalence rates for any use in the past month*

Where females' prevalence is higher, the cell is black with white text; where males' prevalence is higher, the cell is white with black text. Differences of zero and non-significant differences between genders are indicated by a dash (-).

| | Grade | | | | | | |
|--------------------------|-------|-----|-----|-----|------|------|------|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| Cigarettes | -- | -- | 1.2 | -- | 2.4 | 2.6 | 6.5 |
| Over the counter drugs | -- | 0.4 | 1.1 | 1.1 | -- | -- | 1.6 |
| Prescription drugs | -- | -- | 0.6 | 0.6 | -- | -- | 2.0 |
| Inhalants | -- | -- | 0.5 | -- | 0.7 | 1.7 | 3.4 |
| Prescription painkillers | -- | -- | 0.4 | -- | 0.5 | 0.6 | 1.4 |
| Amphetamines | -- | -- | -- | -- | -- | 0.5 | 0.8 |
| Tranquilizers | -- | 0.3 | -- | -- | 0.5 | -- | 1.3 |
| Cocaine | -- | 0.2 | -- | 0.3 | 0.7 | 1.2 | 1.0 |
| Methamphetamines | -- | 0.4 | 0.3 | 0.6 | 0.8 | 1.2 | 1.1 |
| Heroin | 0.2 | 0.3 | 0.3 | 0.4 | 0.8 | 1.1 | 1.2 |
| Crack | 0.1 | 0.2 | 0.4 | 0.7 | 1.2 | 1.7 | 2.3 |
| Ecstasy | -- | 0.2 | 0.4 | 0.5 | 0.9 | 1.2 | 1.8 |
| Hallucinogens | 0.5 | 0.8 | -- | 2.3 | 4.2 | 5.8 | 7.7 |
| Steroids | 0.2 | 0.5 | 0.5 | 1.0 | 1.1 | 1.7 | 1.7 |
| Marijuana | 0.3 | 0.5 | 0.6 | 1.7 | 2.8 | 3.2 | 5.2 |
| Pipe | 0.2 | 0.4 | 0.6 | 0.7 | 0.9 | 1.2 | 1.3 |
| Cigars | 0.5 | 1.1 | 2.2 | 5.0 | 7.9 | 10.7 | 16.7 |
| Smokeless tobacco | 1.1 | 2.4 | 4.6 | 8.2 | 11.7 | 12.9 | 16.8 |

Table 6. *Significant differences ($p < .05$) between male and female prevalence rates alcohol use*

Where females' prevalence is higher, the cell is black with white text; where males' prevalence is higher, the cell is white with black text. Differences of zero and non-significant differences between genders are indicated by a dash (-).

| | Grade | | | | | | |
|----------------------------|-------|-----|-----|-----|-----|-----|-----|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| Past 2-week binge drinking | 0.9 | -- | 1.2 | -- | 3.2 | 5.6 | 6.9 |
| Past Month | 0.9 | -- | 3.5 | 2.3 | -- | 3.0 | 4.9 |
| Lifetime | 5.1 | 2.7 | 2.3 | 3.0 | 1.6 | -- | -- |

Regional Prevalence Rates

This is the fourth year that this report presents the Indiana Survey results for Indiana's Family and Social Service Administration's sub-state planning regions (see Figure 4). Regional data provide more targeted information for local analyses. For the drugs studied, prevalence rates varied considerably among regions and between any given region and the state. This summary does not attempt to capture the extent of this variation; however, a few examples are provided.

Compared to all respondents, statewide, in the...

- Southeast Region, sixth through twelfth graders were more likely to report lifetime use of cigarettes and smokeless tobacco, and tenth through twelfth graders were more likely to report lifetime use of a variety of other substances.
- Northwest Region, sixth through tenth and twelfth graders were less likely to report lifetime use of smokeless tobacco, but students in seventh through twelfth grades were more likely to report lifetime use of a variety of other substances.
- Central Region, students in many grades were less likely to report lifetime use of cigarettes, smokeless tobacco, but students also reported higher prevalence rates for a variety of substances.
- Southwest Region, sixth through twelfth graders were less likely to report lifetime use of a variety of substances, such as marijuana.
- East Region, sixth through twelfth graders were more likely to report lifetime use of a variety of substances, including cigarettes and alcohol.

The reasons for the regional variation are not addressed in this report. Communities may gain a better understanding of their own youth drug use by comparing their prevalence rates (in their local report) to the regional rates presented in this report.




Figure 4. Indiana's Family and Social Service Administration's sub-state planning regions.

Defined Service Area Prevalence Rates

For the second time this year, the Indiana Survey presents results for Indiana's Defined Service Area's (DSAs) (see Figure 5). DSA data provide more targeted information for local analysis. For the drugs studied, prevalence rates varied considerably among DSAs and between any given DSA and the state. This summary does not attempt to capture the extent of this variation; however, a few examples are provided.

Compared to all respondents statewide, in ...

- DSA 5, sixth through twelfth grade students were less likely to report lifetime use of a variety of substances, including cigarettes.
- DSA 7, sixth through twelfth graders were more likely to report lifetime use of cigarettes, and sixth through eighth graders were more likely to report lifetime use of alcohol.
- DSA 10, sixth through ninth graders were less likely to report lifetime use of alcohol.
- DSA 12, sixth, seventh, tenth, and eleventh grade students were more likely to report lifetime use of cigarettes.




Figure 5. Indiana's Defined Service Areas

Risk and Protective Factors

Age of First Use

Research has shown that the younger a person is when she or he begins using alcohol, the more likely the person is to experience alcohol dependence and abuse (Grant, Stinson, & Harford, 2001; Warner & White, 2003). A recent study found that, compared to persons who began drinking at age 21 or older, those who began drinking before age 14 were more likely to experience alcohol dependence later in life (Hingson, Heeren, & Winter, 2006).

In the Indiana Survey data, age of first use was measured by asking, "At what age did you first use ...?" From 1993 to 2006, response options for this question consisted of two-year intervals for ages 8 through 17 (i.e., 8-9 years, 10-11 years, etc.) and open-ended ranges for extreme responses ("7 or less years" and "18 or more years"). From 1993-2006, average age of first use was calculated using the midpoints of the intervals and the upper and lower endpoints for the extreme responses. Starting in 2007, responses included open-ended ranges at the extremes ("7 or younger" and "18 or older") and one-year intervals representing discrete ages (e.g., "8 yrs," "9 yrs," etc.). Average age of first use from 2007 to 2009 was calculated, as

before, with the endpoints of the extreme responses, and with the actual age of first use given for ages 8-17 years. In 2010, response options were changed to broaden the open-ended ranges at the extremes (“10 or younger” and “17 or older”), with one-year intervals representing discrete ages (e.g., “11 yrs,” “12 yrs,” etc.).

Over the years of Indiana Survey data, the average age of first use of gateway drugs (cigarettes, alcohol, and marijuana) has ranged consistently from 12 to 14 years (see Figure 6). Reported initiation of the gateway drugs exhibits a nearly consistent pattern: cigarettes first, then alcohol, followed by marijuana. There were no significant changes between 2010 and 2011 in the reported age of initiation for cigarettes (from 13.02 to 13.04), alcohol (from 13.21 to 13.20), or marijuana (from 13.92 to 13.89).




Figure 6. Trends in average reported age of first use of cigarettes, alcohol, and marijuana.

Perceived Risk of Harm

Prior research has demonstrated that perceived risk of harm of using drugs negatively is related to prevalence of use (Millstein & Halpern-Felsher, 2002; Wild, Hinson, Cunningham, & Bacchiochi, 2001). To examine this relationship in the Indiana Survey data, respondents’ perceived harm of using gateway drugs was correlated with reported monthly use of gateway drugs (see Table 7 for definitions of monthly use of gateway drugs). Perceived risk of harm was measured for the three gateway substances using this question: “How much do you think people risk harming themselves (physically or in other ways) if they... (a) smoke one or more packs of cigarettes per day, (b) smoke marijuana regularly, (c) have five or more drinks once or twice a week.”

Table 7. Glossary of prevalence of monthly gateway drug use for all correlation analyses

| Monthly Use | |
|-------------|--|
| Cigarettes | How many times in the last month (30 days) have you used cigarettes? |
| Alcohol | How many times in the last month (30 days) have you used alcohol (beer, wine, liquor, wine coolers)? |
| Marijuana | How many times in the last month (30 days) have you used marijuana (pot, hash, weed)? |

The results show that higher perceived risk of harm was associated with lower monthly use of cigarettes, marijuana, and alcohol (see Figure 7).




Figure 7. Line of best fit for the correlation of perceived risk of harm and monthly prevalence of gateway drugs.

Perceptions of Peer Disapproval

Research has demonstrated that peer perceptions of disapproval exert an influence that lowers drug use (Butters, 2004; Chassin, Presson, & Sherman, 1984). To examine this relationship in the Indiana Survey data, respondents' perceptions of peer approval were correlated with reported use of gateway drugs (see Table 7 for a definition of monthly use of gateway drugs).

Perceived peer approval was measured for three gateway drugs with the survey items: "How do you think your close friends feel (or would feel) about you doing each of the following things... (a) smoke one or more packs of cigarettes per day; (b) smoke marijuana regularly, (c) have five or more drinks once or twice a week."

The results illustrate that when reported perception of peer disapproval for a particular behavior is stronger, the reported level of the behavior is lower (see Figure 8).




Figure 8. Line of best fit for the correlation of perceived peer approval and monthly prevalence of gateway drugs.

Perceptions of Parents' and Guardians' Approval

The Indiana Survey data measured perceived parental approval for the use of three gateway drugs with the following items: “How wrong do your parents feel it would be for you to... (a) smoke one or more packs of cigarettes per day; (b) smoke marijuana regularly, (c) have five or more drinks once or twice a week.” The results appear in the prevalence tables.

Adolescents’ perceptions of parental approval of substance use are positively related to the frequency of adolescents’ use of alcohol, tobacco and marijuana (McMaster & Wintre, 1996; Lee et al., 2000; Jessor & Donovan, 1978; Barnes & Welte, 1986). To examine this relationship in the Indiana Survey data, respondents’ perceptions of parents’ and guardians’ approval were correlated with reported use of gateway drugs in the previous month. Respondents tended to report engaging in these behaviors more often when they also reported stronger perceived parental approval for the behavior (see Figure 9). These findings are consistent with the research cited.




Figure 9. Line of best fit for the correlation of perceived parental approval and monthly prevalence of gateway drugs.

After-School Activities

Research has demonstrated that supervised after-school activities protect adolescents from using alcohol and other drugs (Grossman, et al., 2002; Riggs & Greenberg, 2004). To determine if this relationship exists in the Indiana Survey data, prevalence of monthly use of the gateway drugs was correlated with an item on student participation in after-school activities such as Afternoons R.O.C.K. in Indiana, L.E.A.D. Initiative, SADD, etc. The results illustrate that participation in after-school activities is associated with lower prevalence of gateway drug use (see Figure 10). These findings support the value of after-school programming as a strategy to prevent or reduce substance use among adolescent populations.




Figure 10. Line of best fit for the correlation of after-school activities and monthly prevalence of gateway drugs.

Perceived Availability of Drugs

Studies have shown that youth perceptions of available access to alcohol and other drugs puts them at risk for use of tobacco, alcohol, and marijuana (Kuntsche & Gmel, 2008; Gillespie, Neale, & Kendler, 2008; Lipperman-Kreda & Grube, 2009).

Perceived availability of drugs was measured by the following items: “How easy would it be for you to get... (a) Cigarettes, (b) Beer, wine or liquor (for example, vodka, whiskey or gin), (c) Marijuana, (d) A drug like cocaine, LSD or amphetamines.”

The results indicate that perceptions of ease of access were associated with higher monthly use of tobacco products, alcohol, and marijuana (see Figure 11). These findings support the value of prevention actions that take into account the social acceptance of drinking and drug use and the physical availability of these drugs in the community.




Figure 11. Line of best fit for the correlation of perceived availability of gateway drugs and monthly prevalence of gateway drugs.

Access to Alcohol

A public health strategy to reduce alcohol abuse by youth is to identify the primary sources of alcohol access and to target interventions to eliminate access at those points (Dent, et al., 2005). Adults over the age of 21 are common sources of alcohol access for youth (Wagenaar, et al., 1996; Jones-Webb, et al., 1997). However, the sources tend to vary by age. Younger adolescents tend to get alcohol from their homes and families, while older adolescents tend to get alcohol from friends and commercial sources (Harrison, et al., 2000; Hearst, et al., 2007).

The percentage of respondents who reported accessing alcohol from specific sources is shown for each grade in Table 8. The results are consistent with the research cited above. Family members were the primary source of alcohol for youth in Grades 6 to 9, excluding the “other ways” category, which potentially encompasses multiple other methods. The higher a respondent’s grade, the more likely the respondent was to report either (1) having had someone buy it for him or her or (2) having received it from a person aged 21 or older. Respondents in higher grades reported obtaining alcohol from commercial outlets at higher rates than did those in lower grades.

These results suggest that youth drinking could be reduced if parents and family members better understood the risk of harm to youth that alcohol presents, as well as ways to prevent youth from accessing alcohol kept in the home. In addition, the results suggest that persons age 21 and older may need to be informed of the laws and penalties that apply to purchasing and/or otherwise supplying alcohol to minors.

Table 8. *Main sources of alcoholic beverages in the past month (percentages)*

| | Grade | | | | | | |
|---|--------|--------|--------|--------|--------|--------|--------|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| No Answer | 6.5 | 5.7 | 5.1 | 5.7 | 6.0 | 6.2 | 7.0 |
| No drink | 88.0 | 84.3 | 78.7 | 71.9 | 67.4 | 64.4 | 57.6 |
| Restaurants/bars/clubs | 0.1 | 0.0 | 0.1 | 0.1 | 0.2 | 0.2 | 0.3 |
| Public events | 0.0 | 0.0 | 0.1 | 0.1 | 0.1 | 0.1 | 0.1 |
| Had someone else buy it | 0.3 | 0.9 | 1.9 | 4.3 | 6.9 | 9.3 | 12.6 |
| Liquor stores/convenience stores/supermarkets | 0.1 | 0.1 | 0.1 | 0.2 | 0.3 | 0.5 | 0.9 |
| Received from person 21 or older | 0.8 | 1.4 | 2.4 | 3.7 | 5.4 | 6.9 | 9.9 |
| Took it from a store | 0.1 | 0.2 | 0.3 | 0.3 | 0.3 | 0.5 | 0.3 |
| Family members | 2.0 | 3.4 | 5.0 | 5.4 | 5.0 | 4.4 | 4.1 |
| Other ways | 2.0 | 4.0 | 6.5 | 8.3 | 8.5 | 7.6 | 7.2 |
| N | 21,640 | 21,623 | 27,745 | 21,562 | 26,006 | 16,542 | 17,560 |

Reasons for Drinking Alcoholic Beverages

The percentages of respondents who reported specific reasons for drinking alcoholic beverages are shown in Table 9. Note that the majority of adolescents in Grades 6 to 9 reported no consumption of alcoholic beverages. Of those who did drink alcoholic beverages, the leading reasons reported were “to have a good time with friends,” “to experiment,” “because it tastes good,” and “to relax or relieve tension.”

Table 9. *Most important reasons for drinking (percentages; multiple responses allowed)*

| Reason | Grade | | | | | | |
|---|--------|--------|--------|--------|--------|--------|--------|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| Did not drink alcoholic beverages | 81.4 | 73.7 | 63.4 | 53.9 | 47.3 | 43.4 | 37.6 |
| To experiment | 10.8 | 16.1 | 22.1 | 23.9 | 24.6 | 22.6 | 21.8 |
| To relax or relieve tension | 1.3 | 3.4 | 7.2 | 11.3 | 15.2 | 18.6 | 21.4 |
| To feel good or get high | 1.1 | 2.7 | 5.2 | 8.9 | 12.1 | 14.6 | 17.1 |
| To seek deeper insights and understanding | 0.5 | 0.8 | 1.5 | 2.3 | 2.3 | 2.7 | 3.3 |
| To have a good time with friends | 2.0 | 5.6 | 11.5 | 20.6 | 27.8 | 33.3 | 39.7 |
| To fit in with a group I like | 0.9 | 1.6 | 2.1 | 2.4 | 2.6 | 2.8 | 2.9 |
| To get away from my problems | 1.8 | 4.1 | 7.0 | 9.4 | 10.8 | 11.4 | 11.2 |
| Because of boredom | 1.9 | 4.3 | 7.3 | 9.2 | 11.3 | 11.9 | 12.6 |
| Because of anger | 1.6 | 2.9 | 5.0 | 6.2 | 6.8 | 7.2 | 6.9 |
| To get through the day | 0.6 | 1.2 | 1.9 | 2.5 | 2.5 | 2.5 | 2.4 |
| To increase the effects of other drugs | 0.3 | 0.5 | 1.0 | 1.7 | 2.2 | 2.5 | 2.8 |
| To decrease the effects of other drugs | 0.2 | 0.2 | 0.3 | 0.4 | 0.4 | 0.4 | 0.5 |
| To get to sleep | 0.6 | 0.9 | 1.7 | 2.2 | 2.5 | 2.9 | 3.6 |
| Because it tastes good | 2.7 | 5.9 | 9.9 | 13.6 | 15.5 | 17.3 | 18.9 |
| Because I am hooked | 0.5 | 0.8 | 1.0 | 1.2 | 1.2 | 1.2 | 1.4 |
| N | 21,640 | 21,623 | 27,745 | 21,562 | 26,006 | 16,542 | 17,560 |

In the Indiana Survey, respondents were instructed to mark as many reasons for drinking alcoholic beverages as apply. Figure 12 shows that, across grade levels, greater numbers of reasons given for drinking were associated with greater frequency of drinking alcoholic beverages during their lifetime.




Figure 12. Number of reasons for drinking versus frequency of drinking during their lifetime.

To analyze the relationship between frequency of drinking and reasons given for drinking, respondents who reported any drinking during their lifetime were divided into three groups: least-frequent drinkers, intermediate-frequency drinkers, and highest-frequency drinkers. Figure 13 shows the relative frequency with which these three groups gave various reasons for their drinking.

Compared to the more frequent drinkers, the least frequent drinkers were markedly more likely to report drinking for experimentation or to fit into a group. The most frequent drinkers were the most likely to report drinking because they were hooked and to increase or reduce the effects of other drugs. Intermediate-frequency drinkers selected most reasons at about as often as did the lowest frequency drinkers. However, these two groups were distinguished from one another by their likelihood of reporting drinking to experiment (to see what it's like). The least frequent drinking group chose these options, respectively, about three times as often as the intermediate frequency drinkers.

Overall, the data suggest that infrequent drinkers tend to drink for social, recreational or coping reasons (e.g. to experiment; to fit into a group; boredom); whereas the most frequent drinkers do so to manage the physiological effects of alcohol or other drugs.




Figure 13. Lifetime alcohol use and reasons for drinking.

Communities That Care Risk and Protective Factors

Risk factors are conditions that increase the chances that children will become involved in problem behaviors in adolescence and young adulthood (Hawkins & Catalano, 2005). Measures included in the Indiana Survey from the Communities That Care (CTC) System can predict alcohol and other drug use, as well as delinquency, dropping out of school, teen pregnancy, violence, depression, and anxiety. Students with elevated risk factor scores have a higher likelihood of substance use and problem behaviors compared to those with low risk factor scores.

CTC is based on the Social Development Strategy that has as its principal focus the strengthening of protective factors. Protective factors are conditions that have a positive influence and “protect” or “buffer” against the negative influences associated with risk factors. The research-based Social Development Model has shown that positive social development in youth is promoted by the following basic factors: 1) opportunities for developmentally appropriate involvement; 2) skills; 3) recognition for effort, improvement and achievement; 4) strong social bonds; and 5) clear consistent standards for behavior. Though closely related, protective factors are not the opposite of risk factors but rather reduce the effects of existing risk factors.

A national study determined optimal cut points to define high and low risk and protective factor scores for students in Grades 6, 8, 10 and 12 (Arthur, Briney,

Hawkins, Abbott, Brooke-Weiss, et al., 2007). Although not shown, those cut points were used in Table 10 and Table 11 to calculate the percentages of Indiana students with high and low risk and protective factor scores, respectively, by grade.

CTC categorizes risk and protective factors into four domains of socialization: Community, Family, School and Peer-Individual (see Table 10 and Table 11). For six of the risk factor scales, the percentage of youth considered high risk increased by successive grades: parental attitudes favorable toward drug use, parental attitudes toward anti-social behavior, low school commitment, attitudes favorable to drug use, perceived risk of drug use, and rewards for antisocial involvement. The percentage of youth who were at high risk between grade levels was generally mixed, with no clear pattern emerging.

All the percentages in Table 10 represent youth who were at/below or above the national cutpoints in terms of risk. For instance, in the Family domain, 41.2 to 51.9 percent of Indiana students in Grades 6, 8, 10 and 12 were at high risk for family conflict. In addition, 31.3 to 44.6 percent of Indiana youth in these grades were at high risk for parental attitudes favorable towards anti-social behavior. In the School domain, 35.3 to 43.6 percent of Indiana students in Grades 6, 8, 10 and 12 were at high risk for low school commitment. In the Peer-Individual domain, 27.5 to 41.2 percent of Indiana students in these grades were at high risk for anti-social peers and 24.4 to 42.7 percent were at high risk for reward for antisocial involvement.

Table 10. *Percentage of students with CTC risk factor scores at/below (low risk) or above (high risk) the national standard[#]*

| | | Grade | | | |
|---|-----------|-------|------|------|------|
| | | 6th | 8th | 10th | 12th |
| Community Domain | | | | | |
| Law and norms favorable to drug use | Low risk | 66.3 | 68.1 | 60.0 | 64.5 |
| | High risk | 33.7 | 31.9 | 40.0 | 35.5 |
| Perceived availability of drugs | Low risk | 74.9 | 73.5 | 64.8 | 61.2 |
| | High risk | 25.1 | 26.5 | 35.2 | 38.8 |
| Family Domain | | | | | |
| Poor family management | Low risk | 75.3 | 73.8 | 75.3 | 71.5 |
| | High risk | 24.7 | 26.2 | 24.7 | 28.5 |
| Family conflict | Low risk | 58.8 | 48.1 | 57.0 | 61.4 |
| | High risk | 41.2 | 51.9 | 43.0 | 38.6 |
| Parental attitudes favorable towards drug use | Low risk | 86.3 | 75.9 | 62.8 | 57.2 |
| | High risk | 13.7 | 24.1 | 37.2 | 42.8 |
| Parental attitudes favorable towards anti-social behavior | Low risk | 68.7 | 58.8 | 56.1 | 55.4 |
| | High risk | 31.3 | 41.2 | 43.9 | 44.6 |
| School Domain | | | | | |
| School academic failure | Low risk | 70.0 | 67.0 | 66.0 | 70.0 |
| | High risk | 30.0 | 33.0 | 34.0 | 30.0 |
| Low school commitment | Low risk | 64.7 | 64.5 | 60.9 | 56.4 |
| | High risk | 35.3 | 35.5 | 39.1 | 43.6 |
| Peer-Individual Domain | | | | | |
| Rebelliousness | Low risk | 69.9 | 74.5 | 72.3 | 72.1 |
| | High risk | 30.1 | 25.5 | 27.7 | 27.9 |
| Early initiation of drug use [†] | Low risk | 80.9 | 73.4 | 74.2 | 74.9 |
| | High risk | 19.1 | 26.6 | 25.8 | 25.1 |
| Attitudes favorable towards antisocial behavior | Low risk | 72.3 | 73.3 | 69.0 | 70.7 |
| | High risk | 27.7 | 26.7 | 31.0 | 29.3 |
| Attitudes favorable towards drug use | Low risk | 84.9 | 76.7 | 67.4 | 67.0 |
| | High risk | 15.1 | 23.3 | 32.6 | 33.0 |
| Perceived risk of drug use | Low risk | 72.3 | 66.9 | 66.4 | 60.5 |
| | High risk | 27.7 | 33.1 | 33.6 | 39.5 |
| Anti-social peers | Low risk | 72.5 | 61.7 | 58.8 | 59.0 |
| | High risk | 27.5 | 38.3 | 41.2 | 41.0 |
| Rewards for antisocial involvement | Low risk | 75.6 | 64.6 | 64.6 | 57.3 |
| | High risk | 24.4 | 35.4 | 35.4 | 42.7 |

Notes. # The national standards (cut-off points) are provided by the Social Development Research Group at the University of Washington.

† The scale is missing one item out of four items from the original CTC scale.

For four scales, the percentage of youth considered at low protection increased between Grades 6 and 12: community rewards for prosocial behavior, family opportunities for prosocial behavior, school opportunities for prosocial behavior, and school rewards for prosocial involvement.

All the percentages in Table 11 represent youth who were at/below or above the national cutpoints for protective factors. For example, in the Community domain,

42.7 to 56.9 percent of Indiana students in Grades 6, 8, 10 and 12 were at or below low protection for community rewards for prosocial behavior. In addition, in the Family domain 35.2 to 40.1 percent of Indiana students in these grades were at or below low protection for family opportunities for prosocial behavior. In the School domain, 35.4 to 46.4 percent of Indiana youth were at or below low protection for school rewards for prosocial involvement. Finally, in the Peer-Individual domain, 49.8 to 53 percent of Indiana youth were at or below low protection for peer-individual interaction with prosocial peers.

Table 11. *Percentage of students with CTC protective factor scores at/below (low protection) or above (high protection) the national standard[#]*

| | | Grade | | | |
|--|-----------------|-------|------|------|------|
| | | 6th | 8th | 10th | 12th |
| Community Domain | | | | | |
| Community rewards for prosocial involvement | Low protection | 42.7 | 56.3 | 56.1 | 56.9 |
| | High protection | 57.3 | 43.7 | 43.9 | 43.1 |
| Family Domain | | | | | |
| Family opportunities for prosocial involvement | Low protection | 36.9 | 35.2 | 40.0 | 40.1 |
| | High protection | 63.1 | 64.8 | 60.0 | 59.9 |
| School Domain | | | | | |
| School opportunities for prosocial involvement | Low protection | 31.9 | 30.5 | 34.8 | 34.0 |
| | High protection | 68.1 | 69.5 | 65.2 | 66.0 |
| School rewards for prosocial involvement | Low protection | 38.0 | 40.1 | 35.4 | 46.4 |
| | High protection | 62.0 | 59.9 | 64.6 | 53.6 |
| Peer-Individual Domain | | | | | |
| Peer-individual interaction with prosocial peers | Low protection | 53.0 | 49.8 | 50.3 | 52.1 |
| | High protection | 47.0 | 50.2 | 49.7 | 47.9 |

Notes. # The national standards (cut-off points) are provided by the Social Development Research Group at the University of Washington.

Consequences of Use

Research has shown that prevention and/or reduction of alcohol and drug use among student populations leads to a decrease in related problems (Hingson, et al., 1996; Wodarski, 1988). Communities are better positioned to address substance use if they understand the relationship of that use to its consequences; these include, for example, poor performance on tests, missing school, and being hung over.

The percentages of respondents who reported specific consequences of alcohol or other drug use are shown for each grade in Table 12. The most commonly reported consequences averaged across grades were having a hangover, nausea and vomiting, and having a memory loss.

Table 12. *Consequences of alcohol, tobacco, and other drug use*

| Consequences of Use | Grade | | | | | | |
|--|--------|--------|--------|--------|--------|--------|--------|
| | 6 | 7 | 8 | 9 | 10 | 11 | 12 |
| Had a hangover | 3.4 | 8.2 | 14.9 | 22.3 | 27.9 | 32.0 | 37.5 |
| Got nauseated or vomited | 3.3 | 6.8 | 11.5 | 17.4 | 23.0 | 28.4 | 34.0 |
| Had a memory loss | 2.1 | 4.3 | 8.1 | 13.0 | 18.0 | 20.9 | 24.8 |
| Performed poorly on a test or project | 2.7 | 4.3 | 5.8 | 6.9 | 7.3 | 7.3 | 7.1 |
| Missed school | 2.5 | 3.4 | 4.8 | 5.8 | 6.2 | 6.5 | 7.7 |
| Got into a fight or argument | 4.3 | 6.4 | 9.0 | 10.7 | 12.1 | 12.5 | 14.0 |
| Damaged property, pulled fire alarms, etc. | 1.4 | 2.4 | 3.7 | 4.4 | 4.5 | 4.5 | 5.2 |
| N | 21,640 | 21,623 | 27,745 | 21,562 | 26,006 | 16,542 | 17,560 |

Substance Abuse Problems

About the CRAFFT Screening Instrument

This is the fourth year the Indiana Survey has included the items from the CRAFFT Screening Instrument. The CRAFFT may be used clinically to detect whether an individual is likely to have problem use or a substance use disorder (Knight, Shrier, Bravender, Farrell, Bilt, et al., 1999). The purpose of including the CRAFFT in the Indiana Survey is to measure the prevalence of substance use-related problems among adolescents. This has been done previously with an adolescent population in Ontario, Canada (Adlaf & Paglia-Boak, 2007).

CRAFFT is a screening instrument. Screening instruments cost little to administer but do a good job of identifying people who are at risk for having a disorder. According to the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition (DSM-IV), there are two substance use disorders: substance abuse and substance dependence. Substance abuse is characterized by recurrent or continued use accompanied by substantial problems resulting from use. Substance dependence is characterized by abuse that is accompanied by “tolerance, withdrawal, and compulsive drug-taking behavior” (American Psychiatric Association, 1994, p. 176). If someone were to screen positive on the CRAFFT in a clinical setting, the clinician would likely refer the patient for a more resource-intensive assessment to diagnose substance use disorder. In the case of substance use disorder, the criteria for such an assessment are specified in the DSM-IV.

Other validated substance use disorder screening instruments, such as the CAGE and AUDIT, ask only about alcohol and were not developed specifically for youth.

In contrast, the CRAFFT was designed for youth, and the CRAFFT asks about both alcohol and other drugs (Knight, et al., 1999).

The CRAFFT has six items; these are presented in Table 13. Response options for each are limited to “yes” and “no.” The items in the CRAFFT scale reflect consequences, patterns, and rationales of use. Affirmative responses to two or more CRAFFT items constitute a positive screen (Knight, Sherritt, Harris, Gates, & Chang, 2003). The CRAFFT was validated for use with people aged 14 to 19 (Knight, Sherritt, Shrier, Harris, & Chang, 2002).

What the CRAFFT measures

We want to emphasize strongly that **the CRAFFT *does not diagnose a substance use disorder* and that the data presented below are *not estimates of the prevalence of substance use disorders in Indiana***. However, the results do suggest problem use among those who screened positive.

The proportion of Indiana Survey respondents aged 14 to 18 who screened positive on the CRAFFT correspond to estimates of dependence and abuse that are substantially similar to those found by the 2006 National Survey on Drug Use and Health for these ages; in 2006, the rate of substance abuse or dependence among persons aged 12 to 17 in the United States was 8.0 percent (SAMHSA, 2007; predictive power estimates for the CRAFFT are described in Knight, et al., 2002).

Results

These analyses are based on data from the 109,128 students (age 14+) who responded to at least one of the CRAFFT items. Of these students, 97,412 responded to all six CRAFFT items. Of the complete responses, 48.3%, or 47,003, had a CRAFT score of zero; 22.3%, or 21,767, had a CRAFFT score of one; and 29.4%, or 28,642, had a CRAFFT score of two or more. These numbers are mostly consistent with the results reported in last year’s monograph.

Table 13 presents the percentages of all respondents age 14 to 17 and older who responded “yes” to each of the CRAFFT items (corresponding to the ages for which the CRAFFT was validated). Results for individual CRAFFT items by grade of respondent are presented in the tables section of this report.

Table 13. *Percentage of respondents, age 14 to 17 and older, indicating “yes” to each of the CRAFFT items (n = 109,128)*

| Item | % |
|---|------|
| Have you ever ridden in a car driven by someone (including yourself) who was high or had been using alcohol or drugs? | 43.8 |
| Do you use alcohol or drugs to relax, feel better about yourself, or fit in? | 18.0 |
| Do you ever use alcohol or drugs while you are alone? | 19.3 |
| Do you ever forget things you did while using alcohol or drugs? | 17.6 |
| Do your family or friends ever tell you that you should cut down on your drinking or drug use? | 8.0 |
| Have you ever gotten into trouble while you were using alcohol or drugs? | 10.9 |

Table 14 shows a breakout of CRAFFT scores by age of respondent. A Pearson’s Chi-Square test indicates significant differences among age groups for the scores ($p < .001$). The prevalence of a score of two or greater was higher for older respondents.

Table 14. *CRAFFT scores of respondents, percentages by age*

| CRAFFT Score | Age of respondent (number (n) of respondents in each group) | | | | |
|--------------|---|------------------|------------------|------------------|-------------------|
| | 14 (n=23,270) | 15 (n=20,717) | 16 (n=22,542) | 17 (n=16,691) | 18+ (n=14,192) |
| 0 | 59.2 | 51.4 | 45.1 | 40.9 | 39.4 |
| 1 | 22.6 | 22.6 | 22.7 | 22.0 | 21.3 |
| 2+ | 18.2 | 26.0 | 32.2 | 37.0 | 39.2 |

Table 15 presents the CRAFFT results by race and Hispanic ethnicity for respondents age 14 to 17 and older. Race and ethnicity data were collapsed into four categories to facilitate statistical tests of differences among categories. The construction of these categories is explained in the Data section of this monograph (see Tables 25 and 26).

A Pearson’s Chi-Square test of differences indicated significant differences among the four race/ethnicity categories ($p < .001$).

Table 15. *CRAFFT scores of respondents, age 14 to 17 and older, percentages by race and ethnic group*

| CRAFFT Score | Race or ethnicity (number (n) of respondents in each group) | | | |
|--------------|---|--|--|---------------------|
| | White or Caucasian (n=73,396) | Black or African-American (n=4,426) | Hispanic Ethnicity (any race) (n=7,447) | Other (n=11,881) |
| 0 | 49.7 | 42.3 | 42.3 | 45.1 |
| 1 | 22.3 | 25.9 | 22.3 | 21.5 |
| 2+ | 28.0 | 31.7 | 35.5 | 33.4 |

Table 16 shows the CRAFFT results by gender of respondent for respondents age 14 to 17 and older. A Pearson's Chi-Square test indicated significant differences between genders ($p < .001$).

Table 16. *CRAFFT scores of respondents, age 14 to 17 and older, percentages by gender*

| CRAFFT Score | Gender (number (n) of respondents) | |
|--------------|------------------------------------|----------------------|
| | Male (n=47,177) | Female (n=50,235) |
| 0 | 49.6 | 47.0 |
| 1 | 20.7 | 23.9 |
| 2+ | 29.7 | 29.1 |

Understanding the prevalence of problem substance use among sub-populations of youth is critical for targeting effective prevention efforts. Selective or indicated prevention strategies are likely to be more appropriate for adolescents who screen positive than are universal prevention approaches.

Gambling

As gambling options and venues in the state expand, monitoring the prevalence of gambling activities among children and adolescents is increasingly important for planning prevention efforts. This is the sixth year that the Indiana Survey has included gambling prevalence items. All gambling-related questions asked about behavior and experiences in the past year. The Indiana Survey included questions about playing cards for money, betting on games, betting on sports, buying lottery tickets, gambling in a casino, and playing online for money.

Gambling Behavior

Compared to last year, the proportion of respondents who reported any gambling behavior increased significantly in Grades 6, 7, and 9 (see Table 17). Gambling behaviors decreased significantly among adolescents in Grade 10. Gambling behaviors included playing cards for money, lottery tickets or scratch offs, at a casino or online and betting on sports teams. Twelfth graders were the most likely to report having gambled, and sixth graders were least likely to report having gambled, though there is no linear pattern; for example, eighth and ninth graders reported having gambled the second-most frequently. Significantly more males reported gambling than females (not shown; Male = 47.0%, Female = 29.6%, $\chi^2 = 4905.754$, $p < .001$).

Table 17. Any gambling behavior by Indiana adolescents in the past year, percentages

| Grade | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change 2010-2011 |
|-------|------|------|------|------|------|------|------|---------------------|
| 6 | 36.3 | 35.4 | 33.6 | 30.3 | 27.3 | 29.9 | 31.2 | 1.3 ** |
| 7 | 40.7 | 39.1 | 38.6 | 33.9 | 33.0 | 34.9 | 36.1 | 1.2 ** |
| 8 | 47.4 | 45.3 | 43.3 | 39.4 | 37.5 | 40.0 | 39.3 | -0.7 |
| 9 | 47.5 | 45.3 | 42.2 | 38.5 | 35.3 | 38.2 | 39.6 | 1.4 ** |
| 10 | 48.0 | 45.2 | 42.5 | 39.0 | 35.4 | 40.0 | 38.4 | -1.6 *** |
| 11 | 47.1 | 45.1 | 41.1 | 37.6 | 34.4 | 38.1 | 37.5 | -0.6 |
| 12 | 55.1 | 55.0 | 50.8 | 47.1 | 44.3 | 47.4 | 46.6 | -0.8 |

** p < .01 *** p < .001

Problem Gambling

Problem gambling items were “During the last 12 months, have you ever felt...? (a) bad about the amount you bet, or about what happens when you bet money, (b) that you would like to stop betting money but didn’t think you could.” These items, used with the permission of the Minnesota Department of Education, help identify gambling problems (Johnson, Hamer, & Nora, 1998; Johnson, et al., 1997). The findings indicate that, compared to 2010, there were no statistically significant changes in the proportion of students in Grades 6 to 12 who reported feeling bad about the amount of money they bet (see Table 18). Also, compared to 2010, a greater proportion of students in Grade 9 reported that they would like to stop betting money but could not (see Table 19).

Table 18. Trend in “felt bad about the amount bet” among respondents reporting any gambling in the past year, percentages

| Grade | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change 2010-2011 |
|-------|------|------|------|------|------|------|------|---------------------|
| 6 | 5.6 | 5.2 | 5.6 | 4.6 | 4.3 | 3.2 | 3.1 | -0.1 |
| 7 | 5.5 | 5.0 | 5.1 | 4.4 | 4.2 | 2.8 | 2.9 | 0.1 |
| 8 | 5.4 | 5.4 | 5.0 | 4.6 | 4.0 | 2.8 | 2.6 | -0.2 |
| 9 | 5.4 | 4.6 | 4.1 | 3.6 | 3.1 | 2.5 | 2.5 | 0.0 |
| 10 | 5.2 | 4.6 | 3.9 | 3.2 | 2.8 | 2.5 | 2.3 | -0.2 |
| 11 | 4.8 | 4.1 | 3.4 | 3.1 | 2.4 | 2.2 | 2.2 | 0.0 |
| 12 | 5.4 | 4.5 | 3.7 | 3.4 | 2.6 | 2.7 | 2.6 | -0.1 |

Table 19. *Trend in “would like to stop betting but could not” among respondents reporting any gambling in the past year, percentages*

| Grade | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change 2010-2011 |
|-------|------|------|------|------|------|------|------|---------------------|
| 6 | 4.9 | 4.5 | 4.9 | 3.8 | 3.8 | 2.5 | 2.4 | -0.1 |
| 7 | 3.7 | 3.6 | 4.0 | 3.2 | 3.1 | 1.9 | 2.0 | 0.1 |
| 8 | 3.4 | 3.2 | 3.5 | 3.0 | 2.8 | 1.9 | 2.0 | 0.1 |
| 9 | 2.6 | 2.3 | 2.6 | 2.2 | 2.1 | 1.5 | 1.7 | 0.2 * |
| 10 | 2.2 | 1.9 | 2.1 | 1.9 | 1.6 | 1.6 | 1.4 | -0.2 |
| 11 | 1.8 | 1.6 | 1.7 | 1.6 | 1.3 | 1.3 | 1.4 | 0.1 |
| 12 | 1.7 | 1.6 | 1.5 | 1.5 | 1.5 | 1.3 | 1.4 | 0.1 |

* p < .05

METHODOLOGY

The methodological history of the *Indiana Survey* can be read about in previous reports. The 2010 report can be found at <http://www.drugs.indiana.edu/indianasurvey>.

Sample

The sampling frame of the Indiana Survey is the universe of all Indiana school corporations and dioceses. The IPRC obtained from the Indiana Department of Education a list of names and addresses of principals, superintendents, and Safe and Drug-Free School Coordinators in all 2,193 public and nonpublic schools. In November 2010, recruitment materials were sent to all school superintendents, principals, and Safe and Drug-Free School Coordinators. The recruitment packet consisted of:

- an invitation letter
- a memorandum of understanding (MOU)
- a statement on parental consent
- an application form
- a list of frequently asked questions
- a stamped addressed return envelope and
- a blank survey form.

The corporations and dioceses had at least seven weeks to respond by sending back an application and a signed MOU to the IPRC. In order to participate, a school or corporation was required to:

- identify a designated coordinator at each public school corporation or multiple coordinators if schools are scattered in several locations
- describe its policy on parental consent
- identify its spring break period
- identify desired survey administration date(s) and
- convey the number of students scheduled to participate in each grade level.

Schools were responsible for obtaining parental consent for students to participate in the Indiana Survey. Information about whether a particular school or corporation participated in any year is strictly confidential. The MOU designated the results of school and corporation-level surveys as the property of the participating school or corporation. These results are released to third parties only with the explicit written permission of the school or corporation's survey coordinator or other appropriate authority.

Participation

Since 1993, participation in the Indiana Survey has been open to all Indiana school corporations, public and nonpublic. The number of usable surveys, shown in Table 20, has fluctuated from year to year. While it is not possible to explain all of the fluctuation, the causes are likely to include:

- The exclusion from participation of fifth-grade students since 1994
- The participation of some schools on biennial and triennial bases
- The arbitrary selection by some schools of the grade levels surveyed
- Variation in the interpretation of parental consent requirements and
- The growing length of the *Indiana Survey* instrument.

Table 20. *Number of usable surveys (1993-2011)*

| Year | Usable Surveys | Year | Usable Surveys |
|------|----------------|------|----------------|
| 1993 | 90,586 | 2003 | 141,342 |
| 1994 | 81,732 | 2004 | 91,577 |
| 1995 | 63,631 | 2005 | 136,782 |
| 1996 | 36,586 | 2006 | 120,914 |
| 1997 | 72,571 | 2007 | 158,632 |
| 1998 | 44,232 | 2008 | 152,732 |
| 1999 | 81,685 | 2009 | 182,496 |
| 2000 | 72,523 | 2010 | 169,059 |
| 2001 | 89,861 | 2011 | 152,678 |
| 2002 | 77,068 | | |

The rates of participation for school corporations, schools and students statewide, and for each FSSA Planning Region, are shown in Table 21. ‘Rate of participation’ is the proportion of corporations, schools and students scheduled to participate in the survey out of the entire student population. There is wide variation in rates of participation across regions. The South West Region had the highest rate of participation among corporations (75.8 percent), schools (46.7 percent) and students (48.5 percent). The East Region had the lowest rate of participation among corporations (44.7 percent), the North East Region had the lowest rate among schools (12.7 percent), and the Central Region had the lowest rate among students (22.4 percent).

This year, for the second time, we report the rate of participation for each Defined Service Area (DSA) (see Table 22). There is wide variation in rates of participation across DSAs. DSA 3 had the highest rate of participation among corporations (84.6 percent), DSA 13 had the highest rate among schools (56.2 percent), and DSA 12 had the highest rate among students (48.1 percent). DSA 4 had the lowest rate of participation among corporations (42.1 percent) and schools (8.4 percent), and DSA 9 had the lowest rate among students (11.2 percent).

Table 21. *Participation rates for students, schools, and corporations by region*

| | State | North West | North Central | North East | West | Central | East | South West | South East |
|---------------------------------|---------|------------|---------------|------------|--------|---------|--------|------------|------------|
| Students | | | | | | | | | |
| Participants | 168,801 | 20,078 | 19,503 | 16,438 | 17,465 | 35,968 | 13,146 | 19,748 | 26,455 |
| Total Population | 559,503 | 70,139 | 77,740 | 54,713 | 56,999 | 160,788 | 32,609 | 40,728 | 65,787 |
| Participation Rate ¹ | 30.2 | 28.6 | 25.1 | 30.0 | 30.6 | 22.4 | 40.3 | 48.5 | 40.2 |
| Schools | | | | | | | | | |
| Participants | 478 | 38 | 56 | 31 | 54 | 94 | 45 | 77 | 83 |
| Total Population | 1,929 | 188 | 313 | 245 | 179 | 374 | 178 | 165 | 287 |
| Participation Rate ² | 24.8 | 20.2 | 17.9 | 12.7 | 30.2 | 25.1 | 25.3 | 46.7 | 28.9 |
| Corporations | | | | | | | | | |
| Participants | 177 | 19 | 24 | 15 | 23 | 26 | 17 | 25 | 28 |
| Total Population | 308 | 33 | 49 | 27 | 39 | 47 | 38 | 33 | 42 |
| Participation Rate ³ | 57.5 | 57.6 | 49.0 | 55.6 | 59.0 | 55.3 | 44.7 | 75.8 | 66.7 |

¹ The percentage of student participants out of the total population, Grades 6 to 12.

² The percentage of schools that participated out of the total population of schools.

³ The percentage of school corporations that participated out of the total population of school corporations.

Table 22. *Participation rates for students, schools, and corporations by DSA*

| | State | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 |
|---------------------------------|---------|--------|--------|--------|--------|--------|--------|--------|
| Students | | | | | | | | |
| Participants | 168,801 | 8,768 | 14,373 | 7,110 | 6,362 | 10,025 | 6,239 | 11,273 |
| Total Population | 559,503 | 47,082 | 41,535 | 33,573 | 32,699 | 32,147 | 36,452 | 35,509 |
| Participation Rate ¹ | 30.2 | 18.6 | 30.6 | 23.0 | 19.5 | 31.2 | 17.1 | 31.8 |
| Schools | | | | | | | | |
| Participants | 478 | 18 | 35 | 28 | 14 | 15 | 27 | 34 |
| Total Population | 1,929 | 134 | 95 | 260 | 167 | 162 | 68 | 147 |
| Participation Rate ² | 24.8 | 13.4 | 36.8 | 10.8 | 8.4 | 9.3 | 39.7 | 23.1 |
| Corporations | | | | | | | | |
| Participants | 177 | 9 | 19 | 11 | 8 | 5 | 9 | 18 |
| Total Population | 308 | 16 | 33 | 13 | 19 | 6 | 20 | 37 |
| Participation Rate ³ | 57.5 | 56.3 | 57.6 | 84.6 | 42.1 | 83.3 | 45.0 | 48.6 |
| | State | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
| Students | | | | | | | | |
| Participants | 168,801 | 13,146 | 8,196 | 11,668 | 10,781 | 18,174 | 12,930 | 13,633 |
| Total Population | 559,503 | 40,005 | 73,042 | 39,658 | 33,712 | 37,779 | 34,045 | 42,265 |
| Participation Rate ¹ | 30.2 | 32.9 | 11.2 | 29.4 | 32.0 | 48.1 | 38.0 | 32.3 |
| Schools | | | | | | | | |
| Participants | 478 | 30 | 48 | 23 | 44 | 57 | 59 | 46 |
| Total Population | 1,929 | 84 | 278 | 71 | 109 | 145 | 105 | 104 |
| Participation Rate ² | 24.78 | 35.7 | 17.3 | 32.4 | 40.4 | 39.3 | 56.2 | 44.2 |
| Corporations | | | | | | | | |
| Participants | 177 | 11 | 8 | 12 | 14 | 18 | 18 | 17 |
| Total Population | 308 | 18 | 15 | 21 | 29 | 26 | 24 | 31 |
| Participation Rate ³ | 57.5 | 61.1 | 53.3 | 57.1 | 48.3 | 69.2 | 75.0 | 54.8 |

¹ The percentage of student participants out of the total population, Grades 6 to 12.

² The percentage of schools that participated out of the total population of schools.

³ The percentage of school corporations that participated out of the total population of school corporations.

Administration of the Survey

Two weeks before the designated survey date, the IPRC mailed to participant schools a packet containing:

- the machine-readable survey forms,
- return shipping labels through the FedEx Package Return Program,
- written instructions for administering the survey, and
- a 7.47 minute survey administration training DVD produced by the IPRC.

The training video describes and demonstrates for school personnel each step of administering the survey to a group of students. The video and the written instructions were provided to improve the consistency of cross-site survey administration. The video is also available for download at http://www.drugs.indiana.edu/data-survey_monograph.html.

The written and video documentation included the following instructions for school personnel who administered the survey:

1. Inform students that:
 - their participation in the survey is completely voluntary,
 - the results are confidential, and
 - there are no penalties for deciding not to participate or for not responding to specific items.
2. Students will need about 30 minutes to complete the survey.
3. Remain seated while students fill out the survey.
4. Instruct the students that, when they complete the survey, they should place their survey forms into the single envelope that the IPRC has provided and which is labeled with a school and classroom code.

No individually identifying information is collected, except gender, grade level, ethnic background and school code. Indiana Survey staff members are available to provide support and answer questions through a toll-free 800 telephone number from Monday through Friday. Schools had a ten-week window to administer the surveys between February 16 and April 24 and were asked to return them immediately to the IPRC.

Survey Instrument

Origin and Relevance of the Questions

Year-to-year changes in the Indiana Survey instrument are documented in the reports corresponding to the years in which they occur. In general, survey items have been derived from a variety of sources for purposes of comparability with national and other state survey data. All items use multiple choice response options.

Alcohol and drug use prevalence items were based on those in the MTF (Johnston, Bachman, & O'Malley, 1989), conducted by the University of Michigan. The Indiana Survey asked respondents about their use of nineteen different types of drugs or drug classifications (see the Indiana Survey instrument in appendix). In some instances, examples and/or slang descriptions appeared in parentheses next to the name of the drug or drug classification. The purpose of this was to elicit responses about the use of a single substance that might be a common ingredient in multiple products (i.e., alcohol in beer, wine, and liquor) or that might have multiple common names, some of them distributed geographically within the state. Other examples include “ecstasy,” “XTC” and “X” for MDMA, and “meth,” “crank,” and “crystal” for methamphetamine.

Prevalence of alcohol, tobacco, and a variety of other drug use is a primary focus of the Indiana Survey. Prevalence refers to the rate of total cases of a condition in a population. For the Indiana Survey, prevalence is the percentage of reported drug use among child and adolescent populations over specified time frames. Definitions for the various prevalence-of-use measures used in this report are listed below.

Lifetime prevalence: the percentage of respondents who report using a particular drug at least once in their lifetime.

Current use or monthly prevalence: the percentage of respondents who report using a particular drug at least once in the 30 days prior to the administration of the survey.

Binge drinking prevalence: the percentage of respondents who report drinking at least five (for males) / four (for females) alcoholic drinks at a sitting in the two weeks prior to administration of the survey.

The perceived risk of harm items have origins in both MTF and the National Survey on Drug Use and Health. The perceived parental approval items recently have become performance measurement requirements for SAMHSA Drug-Free Communities Support Program grantees. The item set for age of first use is similar

to that in the NHSDA. However, in the Indiana Survey these items are closed ended. The access to alcohol and the reasons for drinking items both have origins in the Youth Risk Behavior Survey.

Several risk and protective factor items are from the Communities that Care Survey (SAMHSA, 2009). These items belong to one of four categories of influence: Family, Community, Peer-Individual, School. An example of an item from the Family category is, “How wrong do your parents feel it would be for you to drink beer, wine or hard liquor (vodka, whiskey, or gin) regularly (at least once or twice each month)?” An item from the Community category is, “How easy is it to get cigarettes, smokeless tobacco, cigars, or other tobacco products?” An item from the Peer-Individual category is, “What are the chances you would be seen as cool if you smoked marijuana?” An example from the School category is, “How important do you think the things you learn in school are going to be for you later in life?”

The set of gambling items, added in 2005, were taken from the Minnesota Student Survey (Stinchfield, Kushner, & Winters, 2005; Winters & Anderson, 2000).

Data

The IPRC entered the Indiana Survey data in-house using Pearson optical scanner equipment. The data were cleaned using multiple criteria. A survey could be eliminated from the analysis for three reasons: failure of the respondent to report a grade level, failure of the respondent to answer a majority of the questions, or failure of the error check protocol. Table 23 shows the frequency and proportions of surveys eliminated for these reasons.

Table 23. *Reasons for excluding respondents from the analysis*

| | Frequency | Percent | Cumulative % |
|---|-----------|---------|--------------|
| Rejected Surveys | | | |
| Reported no gender | 3,228 | 1.9 | 1.9 |
| Reported no grade level | 921 | .5 | 2.5 |
| Refused to answer majority of questions | 5,970 | 3.5 | 6.0 |
| Failed error check protocol | 6,004 | 3.6 | 9.6 |
| Total number of usable surveys | 152,678 | 90.4 | 100.0 |
| Total | 168,801 | 100.0 | |

The error check protocol eliminated surveys that met any one of four conditions:

- Respondent provided a substantially inconsistent pattern of responses to prevalence questions (i.e., reported “never used” a particular drug in her or his lifetime and also reported using that drug in the past year or month).
- Pattern of responses was pharmacologically implausible (i.e., a combination of

drugs and frequencies of use whose cumulative effect would be lethal).

- Combination of age and grade was implausible (i.e., a ten-year-old in the eleventh grade.)
- Respondent indicated that he or she did not respond truthfully at all (Item 40). Measurement of this criterion is explained below.

The final item of the Indiana Survey asks students, “How truthfully have you answered these questions,” with response options, (a) “not truthfully at all”, (b) “somewhat truthfully”, and (c) “completely truthfully.” Those who responded “not truthfully at all” were eliminated from the analysis. The distribution of responses to this item is shown in Table 24. Variation, ranging from 1.4 to 2.0 percent (95% confidence interval), was found across regions for students who reported they did not answer truthfully at all.

Table 24. *Response to item on truthful completion of the Indiana Survey by region*

| Region | No Answer | | Not Truthfully at all | | Somewhat Truthfully | | Completely Truthfully | | Total |
|---------------|---------------|-------------|-----------------------|------------|---------------------|------------|-----------------------|-------------|----------------|
| | N | % | N | % | N | % | N | % | N |
| Northwest | 3,293 | 16.4 | 291 | 1.4 | 1,589 | 7.9 | 14,905 | 74.2 | 20,078 |
| North Central | 3,145 | 16.1 | 292 | 1.5 | 1,519 | 7.8 | 14,547 | 74.6 | 19,503 |
| Northeast | 2,058 | 12.5 | 258 | 1.6 | 1,180 | 7.2 | 12,942 | 78.7 | 16,438 |
| West | 2,569 | 14.7 | 291 | 1.7 | 1,333 | 7.6 | 13,272 | 76.0 | 17,465 |
| Central | 6,893 | 19.2 | 708 | 2.0 | 2,690 | 7.5 | 25,677 | 71.4 | 35,968 |
| East | 2,581 | 19.6 | 181 | 1.4 | 948 | 7.2 | 9,436 | 71.8 | 13,146 |
| Southwest | 2,055 | 10.4 | 308 | 1.6 | 1,625 | 8.2 | 15,760 | 79.8 | 19,748 |
| Southeast | 2,674 | 10.1 | 455 | 1.7 | 2,133 | 8.1 | 21,193 | 80.1 | 26,455 |
| Total | 25,268 | 15.0 | 2,784 | 1.6 | 13,017 | 7.7 | 127,732 | 75.7 | 168,801 |

Respondent demographics are shown in Table 25 according to gender, race/ethnicity and grade level. Respondents were asked separate questions about race and Hispanic ethnicity. To facilitate statistical tests of differences among categories, we combined these results to create four categories: Non-Hispanic White or Caucasian, Non-Hispanic Black or African-American, Hispanic Ethnicity (any race), and Other. “Other” included all other respondents, including: (1) all non-Hispanic non-Whites and non-Blacks, including the response categories of Asian, Native Hawaiian or Other Pacific Islander, American Indian or Alaska Native, More than one race, Race not know or other; (2) respondents who indicated non-Hispanic ethnicity but did not respond to the question on race (see Table 26).

Table 25. *Demographic characteristics of the participants*

| | N | % |
|-----------------------|---------|------|
| Gender | | |
| Male | 74,951 | 49.1 |
| Female | 77,727 | 50.9 |
| Grade | | |
| 6 | 21,640 | 14.2 |
| 7 | 21,623 | 14.2 |
| 8 | 27,745 | 18.2 |
| 9 | 21,562 | 14.1 |
| 10 | 26,006 | 17.0 |
| 11 | 16,542 | 10.8 |
| 12 | 17,560 | 11.5 |
| Race/Ethnicity | | |
| White | 113,067 | 74.1 |
| Black | 7,220 | 4.7 |
| Hispanic | 12,220 | 8.0 |
| Other (see Table 26) | 19,599 | 12.8 |
| No answer | 572 | 0.4 |

Table 26. *Components of race/ethnicity category “Other”*

| Race | Ethnicity | |
|---|----------------|----------------|
| | Non-Hispanic | No Answer |
| White | (see Table 25) | 6,041 |
| Black or African-American | (see Table 25) | 1,672 |
| Asian | 2,088 | 229 |
| Native Hawaiian or Other Pacific Islander | 343 | 44 |
| American Indian or Alaskan Native | 698 | 77 |
| More than one race | 5,044 | 691 |
| Race not known or other | 1,012 | 152 |
| No answer | 1,508 | (see Table 25) |

Reliability

Reliability is an estimate of how consistently survey items measure the same concept each time they are administered under the same conditions and with the same subjects. Internal reliability coefficients describe the consistency of results across sets of items (Fink, 1993). Internal reliability coefficients (i.e. Cronbach’s alpha) based on the average correlation among item sets are shown in Table 27. Alpha coefficients range in value from 0 to 1. The higher the score is, the more reliable the item sets are. Nunnally (1978) has indicated 0.7 to be an acceptable reliability coefficient, and many researchers require at least 0.8 for a “good scale.” Based

on these criteria, the internal reliability of the Indiana Survey's item sets is good to excellent, with the exception of the Parental Attitudes Favorable to Drug Use, Low School Commitment, and Early Initiation of Drug Use item sets which are acceptable. This indicates that the item sets are consistent measures of the concepts in the Indiana Survey.

Table 27. Internal reliability of item sets

| Item Set | Question Wording | Number of Items | Cronbach's Alpha |
|--|---|-----------------|------------------|
| Lifetime Use | Have you ever used...? | 19 | .908 |
| Use in the Past Month | How many times in the last month (30 days) have you used... | 19 | .846 |
| Consequences of Use | How many times have you experienced the following due to your drinking or drug use during the past year? | 7 | .865 |
| Gambling | During the last 12 months, how often have you done these activities? | 6 | .790 |
| Perceived Peer Disapproval | How do you think your close friends feel (or would feel) about you doing each of the following things? | 5 | .930 |
| Community Law and Norms Favorable to Drug Use | These questions ask about the neighborhood and community where you live. | 7 | .874 |
| Perceived Availability of Drugs | How easy would it be for you to get...? | 4 | .856 |
| Poor Family Management | The rules in my family are clear. | 8 | .851 |
| Family Conflict | We argue about the same things in my family over and over. | 3 | .837 |
| Parental Attitudes Favorable to Drug Use | How wrong do your parents feel it would be for you to... | 3 | .811 |
| Parental Attitudes Favorable to Anti-social Behavior | How wrong do your parents feel it would be for you to... | 3 | .786 |
| Low School Commitment | Now thinking back over the past year in school, how often did you... | 3 | .709 |
| Rebelliousness | I like to see how much I can get away with. | 3 | .842 |
| Early Initiation of Drug Use† | If you have ever used these drugs, at what age did you first use them? | 3 | .754 |
| Peer Attitudes Favorable to Anti-social Behavior | How wrong do you think it is for someone your age to...? | 5 | .863 |
| Peer Attitudes Favorable to Drug Use | How wrong do you think it is for someone your age to...? | 4 | .880 |
| Perceived Risk of Drugs | How much do you think people risk harming themselves (physically or in other ways) if they... | 5 | .839 |
| Anti-social Peers | In the past year (12 months), how many of your best friends have... | 6 | .852 |
| Community Rewards for Prosocial Involvement | My neighbors notice when I am doing a good job and let me know. | 3 | .929 |
| Family Opportunities for Prosocial Involvement | My parents ask me what I think before most family decisions affecting me are made. | 3 | .772 |
| School Opportunities for Prosocial Involvement | In my school, students have lots of chances to help decide things like class activities and rules. | 5 | .775 |
| School Rewards for Prosocial Involvement | My teacher(s) notices when I am doing a good job and lets me know about it. | 4 | .807 |
| Interaction with Prosocial Peers | Think of your four best four friends (the friends you feel closest to). In the past year (12 months), how many of your best friends have... | 5 | .826 |

Notes. † The scale is missing one item out of four items from the original CTC scale.

Validity

The validity of a survey is the extent to which it measures what its designers intend it to measure (Fink, 1993). The Indiana Survey is intended to measure the reported prevalence of substance use, the consequences of use, perceptions of peer approval of substance use, perceptions of personal safety, prevalence of gambling, and other concepts. The Indiana Survey measures these concepts among children and adolescents in Grades 6 through 12 located in schools across Indiana at a given point in time. The Indiana Survey uses a nonrandom sampling procedure, which presents a threat to the survey's external validity. A nonrandom procedure was used because a longstanding priority of the Indiana Survey has been to provide local level data to any and all schools or school corporations that request it. This nonrandom procedure may introduce bias, so the findings of the Indiana Survey should be generalized with caution.

To the extent that school districts have obtained information from every student in a particular grade level, the local results represent those community populations. In this respect, participating school districts can examine their own student response rates and decide how representative their survey data are for their local youth population. Formula 1.0 shows how the student response rate may be calculated from data in the local report provided to each participating school or school district.

Formula 1.0 Response Rate = Number of Respondents / Total Number of Students x 100

Where "Number of Respondents" is the number of students who completed and turned the survey in, and where "Total Number of Students" is the number of students whom the school corporation intended to survey.

If the response rate is 90% or greater, then the district level results (not shown in this report) are likely to be representative of the local student population.

At the sub-state regional and defined service area (DSA) levels, the ability to generalize the data is much lower because many school districts, and therefore many children and adolescents in the regional and DSA populations, did not participate in the Indiana Survey. As shown in Tables 21 and 22, only 22.4 to 48.5 percent of students participated in the Indiana Survey across regions, and 11.2 to 48.1 percent across DSAs, respectively. The best approach to address this limitation is to assess the extent to which the samples of student respondents compare to the entire population of students in each region and DSA. This comparison of regional respondents and the population is shown by gender, ethnicity, race, and grade level in Table 28, and of DSA respondents in Table 29. Although this comparison does not provide a basis for generalizing results at the regional or DSA level, it provides guidance on where the data do and do not coincide proportionally with demographic subgroups in the regional and DSA populations.

As shown in Table 28, the majority of samples either under- or over-represent the proportions of students by gender, ethnicity, race, or grade level in a region. For example, in the West region, females are overrepresented in the Indiana Survey sample and Whites are underrepresented. This means that the sample has proportionally more females than the West region as a whole, and this bias should be taken into consideration when interpreting the results of the Indiana Survey data. Likewise, the sample has proportionally fewer White students.

Table 28. *Regional Comparison of Samples and Population by Demographics*

| | Indiana | | Northwest | | Northcentral | | Northeast | | West | |
|-----------------------------------|---------|--------|-----------|--------|--------------|--------|-----------|--------|-----------|--------|
| | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample |
| Gender | | | | | | | | | | |
| Male | 51.0 | 49.1 * | 50.9 | 48.5 * | 51.1 | 49.0 * | 50.9 | 49.8 | 51.2 | 49.3 * |
| Female | 49.0 | 50.9 * | 49.1 | 51.5 * | 48.9 | 51.0 * | 49.1 | 50.2 | 48.8 | 50.7 * |
| Race/Ethnicity[#] | | | | | | | | | | |
| White | 75.3 | 74.3 * | 57.4 | 58.9 * | 74.2 | 73.8 | 76.6 | 74.8 * | 85.5 | 79.6 * |
| Black | 11.8 | 4.7 * | 21.0 | 9.6 * | 9.7 | 1.4 * | 10.0 | 4.6 * | 2.9 | 2.1 |
| Hispanic | 7.2 | 8.0 * | 16.1 | 16.9 * | 10.1 | 14.2 * | 7.1 | 6.4 | 6.2 | 6.5 |
| Other | 5.7 | 12.9 * | 5.5 | 14.6 * | 6.0 | 10.5 * | 6.3 | 14.2 * | 5.4 | 11.7 * |
| Grade | | | | | | | | | | |
| 6 | 14.3 | 14.2 | 14.0 | 17.7 * | 14.3 | 13.0 | 14.2 | 8.1 * | 14.1 | 13.8 |
| 7 | 14.4 | 14.2 | 14.0 | 14.9 | 14.4 | 17.6 * | 14.1 | 11.1 * | 14.1 | 11.3 * |
| 8 | 14.3 | 18.2 * | 14.2 | 18.2 * | 14.4 | 18.3 * | 14.1 | 13.5 | 14.1 | 21.0 * |
| 9 | 15.1 | 14.1 * | 15.5 | 12.8 * | 15.0 | 15.5 | 15.8 | 18.1 * | 15.2 | 12.2 * |
| 10 | 14.6 | 17.0 * | 14.7 | 14.9 | 14.4 | 14.4 | 14.4 | 20.9 * | 14.5 | 18.8 * |
| 11 | 13.9 | 10.8 * | 14.1 | 10.2 * | 13.9 | 11.2 * | 13.6 | 14.9 | 14.1 | 9.3 * |
| 12 | 13.4 | 11.5 * | 13.5 | 11.2 * | 13.6 | 10.1 * | 13.8 | 13.2 | 13.9 | 13.6 * |
| | Indiana | | Central | | East | | Southwest | | Southeast | |
| | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample |
| Gender | | | | | | | | | | |
| Male | 51.0 | 49.1 * | 50.7 | 48.2 * | 51.0 | 49.1 * | 50.6 | 49.8 | 51.5 | 49.6 * |
| Female | 49.0 | 50.9 * | 49.3 | 51.8 * | 49.0 | 50.9 * | 49.4 | 50.2 | 48.5 | 50.4 * |
| Race/Ethnicity[#] | | | | | | | | | | |
| White | 75.3 | 74.3 * | 67.4 | 68.0 | 87.2 | 73.9 * | 87.5 | 85.6 * | 91.1 | 82.2 * |
| Black | 11.8 | 4.7 * | 19.6 | 9.5 * | 5.3 | 5.5 | 4.9 | 1.0 * | 2.4 | 1.7 |
| Hispanic | 7.2 | 8.0 * | 6.7 | 7.0 | 1.9 | 4.9 * | 2.4 | 4.1 * | 2.5 | 4.7 * |
| Other | 5.7 | 12.9 * | 6.3 | 15.5 * | 5.6 | 15.7 * | 5.2 | 9.3 * | 4.0 | 11.3 * |
| Grade | | | | | | | | | | |
| 6 | 14.3 | 14.2 * | 14.7 | 16.0 * | 14.3 | 12.1 * | 14.3 | 15.2 | 14.0 | 14.1 |
| 7 | 14.4 | 14.2 * | 14.7 | 14.9 | 14.1 | 16.3 * | 14.2 | 15.7 | 14.4 | 11.7 * |
| 8 | 14.3 | 18.2 * | 14.6 | 19.3 * | 14.2 | 19.4 * | 14.1 | 16.2 * | 14.4 | 18.5 * |
| 9 | 15.1 | 14.1 * | 14.6 | 14.0 | 14.9 | 16.0 | 15.1 | 13.3 * | 15.1 | 12.7 * |
| 10 | 14.6 | 17.0 * | 15.0 | 17.0 * | 14.6 | 15.4 | 14.4 | 16.3 * | 14.4 | 18.4 * |
| 11 | 13.9 | 10.8 * | 13.6 | 9.6 * | 13.9 | 11.2 * | 14.2 | 11.3 * | 14.1 | 10.6 * |
| 12 | 13.4 | 11.5 * | 12.8 | 9.1 * | 14.0 | 9.6 * | 13.7 | 12.1 | 13.6 | 13.9 |

* The sample is significantly different from the population ($p < .05$).

No answer is not included in calculation of the percentages for the sample.

As shown in Table 29, the majority of samples either under- or over-represent the proportions of students by gender, ethnicity, race, or grade level in a DSA. For example, in DSA 2, males are underrepresented in the Indiana Survey sample and Hispanics are overrepresented. This means that the sample has proportionally fewer males than DSA 2 as a whole, and this bias should be taken into consideration when interpreting the results of the Indiana Survey data. Likewise, the sample has proportionally more Hispanic students for DSA 2. This is a known limitation to applying the Indiana Survey results to non-Hispanic groups, which are under-represented in the sample.

Table 29. DSA comparison of population and samples by demographics

| | Indiana | | DSA1 | | DSA2 | | DSA3 | | DSA4 | |
|-----------------------------------|---------|--------|------|--------|------|--------|------|--------|------|--------|
| | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample |
| Gender | | | | | | | | | | |
| Male | 51.0 | 49.1 * | 50.8 | 47.4 * | 51.3 | 49.3 * | 51.2 | 49.6 | 50.7 | 49.2 |
| Female | 49.0 | 50.9 * | 49.2 | 52.6 * | 48.7 | 50.7 * | 48.8 | 50.4 | 49.3 | 50.8 |
| Race/Ethnicity[#] | | | | | | | | | | |
| White | 75.3 | 74.3 * | 44.2 | 46.6 * | 82.6 | 72.3 * | 71.5 | 76.8 * | 76.7 | 71.9 * |
| Black | 11.8 | 4.7 * | 29.8 | 16.8 * | 4.4 | 2.3 | 12.7 | 1.4 * | 4.4 | 1.4 |
| Hispanic | 7.2 | 8.0 * | 20.0 | 19.6 | 8.5 | 14.0 * | 9.8 | 12.5 * | 14.2 | 16.0 |
| Other | 5.7 | 12.9 * | 6.0 | 17.0 * | 4.4 | 11.4 * | 6.0 | 9.3 * | 4.7 | 10.7 * |
| Grade | | | | | | | | | | |
| 6 | 14.3 | 14.2 | 14.0 | 20.2 * | 14.2 | 14.4 | 14.0 | 14.5 | 14.6 | 10.9 * |
| 7 | 14.4 | 14.2 | 14.1 | 15.5 | 14.1 | 14.7 | 14.3 | 18.5 * | 14.8 | 15.0 |
| 8 | 14.3 | 18.2 * | 14.0 | 21.3 * | 14.5 | 16.4 * | 14.4 | 21.0 * | 14.5 | 17.6 * |
| 9 | 15.1 | 14.1 * | 15.7 | 9.6 * | 15.5 | 15.1 | 14.9 | 14.7 | 14.6 | 13.3 |
| 10 | 14.6 | 17.0 * | 14.7 | 13.9 | 14.2 | 16.1 * | 14.9 | 12.8 | 14.2 | 20.5 * |
| 11 | 13.9 | 10.8 * | 14.3 | 8.1 * | 13.8 | 11.6 * | 14.0 | 9.9 * | 13.8 | 11.3 |
| 12 | 13.4 | 11.5 * | 13.2 | 11.4 | 13.7 | 11.7 * | 13.6 | 8.6 * | 13.5 | 11.3 |
| | DSA5 | | DSA6 | | DSA7 | | DSA8 | | DSA9 | |
| | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample |
| Gender | | | | | | | | | | |
| Male | 50.8 | 49.8 | 50.8 | 50.1 | 50.8 | 48.7 * | 50.5 | 48.1 * | 50.7 | 48.3 * |
| Female | 49.2 | 50.2 | 49.2 | 49.9 | 49.2 | 51.3 * | 49.5 | 51.9 * | 49.3 | 51.7 * |
| Race/Ethnicity[#] | | | | | | | | | | |
| White | 66.8 | 70.4 * | 85.9 | 82.4 * | 88.6 | 74.0 * | 82.9 | 74.9 * | 44.0 | 38.0 * |
| Black | 16.7 | 6.4 * | 3.6 | 2.6 | 4.2 | 5.0 | 5.0 | 3.7 | 38.2 | 27.0 * |
| Hispanic | 7.9 | 6.9 | 5.0 | 3.5 | 2.4 | 5.9 * | 4.3 | 7.8 * | 10.7 | 13.1 * |
| Other | 8.6 | 16.4 * | 5.5 | 11.5 * | 4.8 | 15.0 * | 7.8 | 13.5 * | 7.1 | 21.9 * |
| Grade | | | | | | | | | | |
| 6 | 14.3 | 5.3 * | 14.1 | 25.1 * | 14.0 | 12.7 | 14.7 | 13.9 | 14.9 | 20.2 * |
| 7 | 14.1 | 11.6 * | 14.1 | 5.5 * | 13.8 | 16.3 * | 14.9 | 21.2 * | 14.6 | 4.0 * |
| 8 | 13.9 | 10.9 * | 14.1 | 23.9 * | 14.0 | 18.0 * | 14.6 | 21.7 * | 14.5 | 18.0 * |
| 9 | 16.5 | 19.5 * | 15.2 | 5.5 * | 14.9 | 15.4 | 14.6 | 12.3 * | 14.5 | 15.8 |
| 10 | 14.2 | 21.7 * | 14.7 | 21.5 * | 14.8 | 15.3 | 14.4 | 13.9 | 15.7 | 22.4 * |
| 11 | 13.4 | 16.0 * | 14.1 | 4.9 * | 14.4 | 12.1 * | 13.7 | 8.1 * | 13.5 | 10.6 * |
| 12 | 13.7 | 15.0 | 13.7 | 13.5 | 14.1 | 10.3 * | 13.1 | 8.9 * | 12.3 | 8.9 * |

Table 29. *DSA comparison of population and samples by demographics (continued)*

| | DSA10 | | DSA11 | | DSA12 | | DSA13 | | DSA14 | |
|-----------------------------------|-------|--------|-------|--------|-------|--------|-------|--------|-------|--------|
| | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample | Pop. | Sample |
| Gender | | | | | | | | | | |
| Male | 50.9 | 48.5 * | 51.4 | 49.4 * | 51.8 | 49.7 * | 50.9 | 50.1 | 50.9 | 48.9 * |
| Female | 49.1 | 51.5 * | 48.6 | 50.6 * | 48.2 | 50.3 * | 49.1 | 49.9 | 49.1 | 51.1 * |
| Race/Ethnicity[#] | | | | | | | | | | |
| White | 89.9 | 81.5 * | 88.7 | 82.4 * | 90.6 | 81.2 * | 86.4 | 85.3 * | 90.1 | 82.8 * |
| Black | 2.7 | 2.3 | 4.9 | 1.8 | 1.4 | 1.6 | 5.8 | 1.2 * | 3.2 | 2.0 |
| Hispanic | 3.0 | 4.2 | 2.2 | 3.7 | 2.9 | 5.1 * | 1.8 | 3.2 | 3.0 | 5.2 * |
| Other | 4.4 | 12.0 * | 4.2 | 12.1 * | 5.1 | 12.1 * | 6.0 | 10.2 * | 3.7 | 10.0 * |
| Grade | | | | | | | | | | |
| 6 | 14.5 | 13.7 | 14.3 | 12.3 | 13.9 | 11.3 * | 14.4 | 15.6 | 14.1 | 15.3 |
| 7 | 14.4 | 13.9 | 14.7 | 14.4 | 14.5 | 14.3 | 14.2 | 15.5 | 14.2 | 12.2 * |
| 8 | 14.5 | 17.6 * | 14.4 | 17.4 * | 14.3 | 17.0 * | 14.2 | 16.6 * | 14.4 | 20.5 * |
| 9 | 15.0 | 15.3 | 14.9 | 18.2 * | 15.2 | 15.3 | 15.1 | 13.1 * | 15.0 | 10.8 * |
| 10 | 14.5 | 17.0 * | 14.2 | 15.2 | 14.3 | 17.0 * | 14.3 | 16.3 * | 14.5 | 18.6 * |
| 11 | 13.6 | 11.9 | 13.6 | 12.1 | 14.1 | 12.2 | 14.2 | 11.3 * | 14.0 | 8.5 * |
| 12 | 13.5 | 10.6 * | 13.9 | 10.4 * | 13.7 | 13.0 | 13.6 | 11.8 | 13.8 | 14.1 |

Notes. Pop. = Population; Sam. = Sample

* The sample is significantly different from the population ($p < .05$)

No answer is not included in calculation of the percentages for the sample.

The strength of the Indiana Survey is that it collects and reports valid and reliable data for local school corporations. At the state level, the data has the same degree of reliability. However, according to sampling theory, the state-level data is less valid (Levy & Lemeshow, 2003). This tradeoff reflects the state's current priorities for serving the substance use prevention needs of local communities. At the same time, statewide results from the Indiana Survey have a longstanding and high degree of concurrence with state-level surveys that use probability sampling, such as the Youth Risk Behavior Survey and the Monitoring the Future survey. This high degree of concurrence is due to the large percentage of school corporations (57.5 percent in 2011) and students who participate in the Indiana Survey. Due to this level of agreement, it is reasonable to compare the statewide Indiana Survey data with national survey results. Furthermore, the results are consistent with findings from other prevention studies and this reinforces our confidence in the validity of Indiana Survey data at the state level.

HOW TO INTERPRET THE RESULTS

Margin of Error

The Indiana Survey has a margin of error plus or minus less than 1 percent with a 95 percent confidence interval. This means that if we selected a group of students 100 times and asked a given question from this survey each time, then in 95 of those instances, the percentage of students giving a particular answer would be within one percentage point of the percentage who gave that same answer in this year's results of the Indiana Survey. Margin of error and confidence intervals, however, have limited applicability to the Indiana Survey because respondents were chosen nonrandomly.

2010-2011 Changes

For the statewide data tables, values shown in the 'change' column refer to statistically significant observed differences in prevalence of use between years 2010 and 2011. Statistical significance describes a mathematical measure of difference between groups (Daniel, 1991). The difference is said to be statistically significant if it is greater than what might be expected to happen by chance alone. "P" is an estimate of the probability that the result has occurred by statistical accident. In this report, the p-value is set at 0.05, which means that the difference or change described has a 95 percent chance of not having been due to chance alone. The plus or minus sign next to the change value indicates whether the 2011 percentage is an increase or decrease, respectively, from the previous year.

For the regional data tables, the symbols ▲ or ▼ are used to indicate statistically significant regional differences in prevalence of use compared to the entire state. The ▲ symbol indicates that the regional rate is higher than the state rate, and the ▼ symbol indicates that the regional rate is lower than the state rate.

Generalizing Local *Indiana Survey* Data to Local Populations

The IPRC provides each participant school corporation with a local report of the Indiana Survey results from its district. This section describes the method for estimating how well local data applies to local populations. The IPRC strongly encourages school corporations to share the results of this analysis with citizens and organizations engaged in local surveillance and planning.

The Response Rate

The response rate estimates how well the local results of the Indiana Survey describe all the youth in a grade or grades, in a school or schools, or in an entire school corporation.

If the response rate is 90 or greater, then the results are likely to be representative of the population defined. If the response rate is lower than 90%, results should be interpreted cautiously. In this case, the results might not be representative of students who were absent the day the survey was administered or of those who did not turn in a survey.

In any case, results do not generalize to those whose responses the IPRC systematically excluded from tabulation (see the “Data” section, above, for details).

The IPRC is unable to calculate the response rate for individual localities because one of the components of the response rate depends on information not collected along with the survey. This information, the Total Number of Students, is explained below.

Two pieces of information are required to calculate the response rate: “the total number of students” and “the total number of respondents.”

The Total Number of Students

The total number of students can only be determined by each school district. The total number of students is the number of students that the school district intended to survey. For instance, if the intention was to survey all ninth and tenth-grade students in all schools then the total enrollment number in Grades 9 through 10 would constitute the total number of students.

Total Number of Respondents

Local reports of the Indiana Survey issued to participating school corporations include a table similar to Table 30. The asterisk (*) in Table 30 illustrates where to locate the total number of respondents.

Table 30. *Sample Table from Local Report*

| | Frequency | Percent |
|---|-----------|---------|
| No gender reported | | |
| No grade reported | | |
| Unusable Surveys/Refused to Participate | | |
| Surveys Rejected Due to Error Check | | |
| Total Number of Usable Surveys | | |
| Total | ##### * | 100.0 |

Calculating the Response Rate

To determine the response rate for a locality, divide the Total Number of Respondents by the Total Number of Students and multiply the result by 100.

$$\text{Response Rate} = \left(\frac{\text{Number of Respondents}}{\text{Total Number of Students}} \right) \times 100$$

Identifying Significant Differences between Local and State Prevalence Rates

In the local report, the first set of tables, titled “Significant Difference between Local and State Drug Use Rates, 2011,” show where a school corporation had statistically significant higher or lower rates compared to the state. Positive numbers indicate that a school corporation had rates higher than those of the state. Negative numbers indicate that a school corporation had rates lower than those of the state. Dash marks appear where there were no statistically significant differences between the school corporation and the state.

If a school corporation had prevalence rates higher than those of the state on particular drugs, the community might want to consider targeting prevention efforts at those drugs.

Practical Significance

The next set of tables in a local report shows the percentage of student participants who reported that they used specific drugs daily, during the past month and during

their lifetime. These rates are shown for the school district over a five year period if such data are available and compared to the state and most recently available national rates.

Targeting prevention efforts at particular drugs might be appropriate even though local prevalence rates are not higher than those of the state. For instance, if sixth-grade students in a corporation had a prevalence rate similar to that of the state for a specific drug, but the rate itself was high, prevention or treatment programs might be needed for sixth graders.

Key points for interpreting prevalence rate results of the Indiana Survey:

- Monthly prevalence is a general indicator of frequent or regular use of a drug.
- Lifetime prevalence is a general indicator of occasional or experimental use of a drug.
- “Prevalence” indicates any use of a drug within the specified time frame, regardless of its severity or frequency within that time frame (for example, any use of marijuana in the last month could be one time or two hundred times). Use monthly and lifetime prevalence rates in conjunction with detailed frequencies of use (for example, 1-5 times, 6-19 times, 20-40 times, more than 40 times). These are provided in tables in the back of a local report.

Interpreting Local Prevalence Rates of Gateway Drug Use

- Cigarette smoking is the best predictor of future drug use.
- Binge drinking is a key indicator of future substance abuse.
- Alcohol use indicates a need for student assistance programs.
- Marijuana use is a key indicator of future drug use.
- Grades with the most rapid increases indicate appropriate timing of secondary prevention strategies.

If a school corporation has participated more than three times in the Indiana Survey, it has sufficient data to examine changes over time. If enrollment is relatively stable over time, trends and patterns may appear. A commonly observed pattern is the “cohort effect.” For certain groups of youth, prevalence rates are inconsistent with the average use rates. These differences often persist over time. For instance, eighth-grade students this year may have used particular drugs 10% more than eighth-grade students two years ago. This pattern may continue from year to year until the students graduate.

References

- Adlaf, E. M., & Paglia-Boak, A. (2007). Drug use among Ontario students, 1977-2007: Detailed OSDUS findings [CAMH Research Document series No. 20]. Toronto: Center for Addiction and Mental Health. Retrieved July 9, 2008 from http://www.camh.net/Research/Areas_of_research/Population_Life_Course_Studies/OSDUS/OSDUHS2007_DrugDetailed_final.pdf.
- American Psychiatric Association. (1994). Diagnostic and statistical manual of mental disorders (4th ed.). Washington, DC.
- Arthur, M. W., Briney, J. S., Hawkins, J. D., Abbott, R.D. Brooke-Weiss, B. L., & Catalano, R. F. (2007). Measuring risk and protection in communities using the Communities That Care Youth Survey. *Evaluation and Program Planning*, 20, 197-211.
- Bailey, S. L. (1992). Adolescents' multisubstance use patterns: The role of heavy alcohol and cigarette use. *American Journal of Public Health*, 82(9), 1220-1224.
- Barnes, G. M., & Welte, J. W. (1986). Patterns and predictors of alcohol use among 7-12th grade students in New York state. *Journal of Studies on Alcohol*, 47(1), 53-62.
- Butters, J. E. (2004). The impact of peers and social disapproval on high-risk cannabis use: Gender differences and implications for drug education. *Drugs Education Prevention Policy*, 11(5), 381-390.
- Chassin, L., Presson, C. C., & Sherman, S. J. (1984). Cognitive and social influence factors in adolescent smoking cessation. *Addictive Behaviors*, 9(4), 383-390.
- Daniel, W. W. (1991). *Biostatistics: A foundation for analysis in the health sciences* (5th ed.). New York, NY: Wiley.
- Dent, C. W., Grube, J. W., & Biglan, A. (2005). Community level alcohol availability and enforcement of possession laws as predictors of youth drinking. *Preventive Medicine*, 40(3), 355-362.
- Donovan, J. E., & Jessor, R. (1983). Problem drinking and the dimension of involvement with drugs: A Guttman scalogram analysis of adolescent drug use. *American Journal of Public Health*, 73(5), 543-552.
- Fink, A. (1993). *Evaluation fundamentals*. Newbury, CA: Sage Publication, Inc.
- Fleming, R., Leventhal, H., Glynn, K., & Ershler, J. (1989). The role of cigarettes in the initiation and progression of early substance use. *Addictive Behaviors*, 14(3), 261-272.
- Gillespie, N. A., Neale, M. C., & Kendler, K. S. (2008). Pathways to cannabis abuse: A multi-stage model from cannabis availability, cannabis initiation and progress to abuse. *Addiction*, 104, 430-438.
- Golub, A., & Johnson, B. D. (2001). Variation in youthful risks of progression from alcohol and tobacco to marijuana and to hard drugs across generations. *American Journal of Public Health*, 91(2), 225-232.
- Grant, B. F., Stinson, F. S., & Harford, T. C. (2001). Age at onset of alcohol use and DSM-IV alcohol abuse and dependence: A 12-year follow-up. *Journal of Substance Abuse*, 13(4), 493-504.
- Grossman, J. B., Price, M. L., Fellerath, V., Jucovy, L. Z., Kotloff, L. J., Raley, R., et al.

- (2002). Multiple choices after school: Findings from the extended-service schools initiative. Philadelphia, PA. : Public/Private Ventures.
- Harrison, P. A., Fulkerson, J. A., & Park, E. (2000). The relative importance of social versus commercial sources in youth access to tobacco, alcohol, and other drugs. *Preventive Medicine*, 31(1), 39-48.
- Hawkins, J. D., & Catalano, R. F. (2005). Investing in your community's youth: An introduction to the Communities that Care System. Retrieved July 1, 2010 from <http://download.ncadi.samhsa.gov/Prevline/pdfs/ctc/Investing%20in%20Your%20Community's%20Youth.pdf>.
- Hearst, M. O., Fulkerson, J. A., Maldonado-Moline, M. M., Perry, C. L., & Komro, K. A. (2007). Who needs liquor stores when parents will do? The importance of social sources of alcohol among young urban teens. *Preventive Medicine*, 44, 471-476.
- Hingson, R., McGovern, T., Howland, J., Heeren, T., Winter, M., & Zakocs, R. (1996). Reducing alcohol-impaired driving in Massachusetts: The saving lives program. *American Journal of Public Health*, 86(6), p791-791.
- Hingson, R. W., Heeren, T., & Winter, M. R. (2006). Age at drinking onset and alcohol dependence: Age at onset, duration, and severity. *Archives of Pediatrics & Adolescent Medicine*, 160(7), 739-746.
- Johnson, E. E., Hamer, R., Nora, R. M., Tan, B., Eisenstein, N., & Engelhart, C. (1997). The Lie/Bet questionnaire for screening pathological gamblers. *Psychological Reports* 80, 83-88.
- Johnson, E. E., Hamer, R., & Nora, R. M. (1998). The Lie/Bet questionnaire for screening pathological gamblers: A follow-up study. *Psychological Reports* 83, 1219-1224.
- Johnston, L. D., Bachman, J. G., & O'Malley, P. M. (1989). Drug use, drinking, and smoking: National survey results from high school, college, and young adult populations, 1975-1989. Rockville, MD: National Institute on Drug Abuse.
- Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2009). Teen marijuana use tilts up, while some drugs decline in use. University of Michigan News Service, Ann Arbor. Retrieved June 1, 2010, from <http://www.monitoringthefuture.org>.
- Johnston, L. D., O'Malley, P. M., Bachman, J. G., & Schulenberg, J. E. (2011). Monitoring the Future national results on adolescent drug use: Overview of key findings, 2010. Ann Arbor: Institute for Social Research, The University of Michigan, Retrieved June 1, 2011, from <http://www.monitoringthefuture.org>.
- Jones-Webb, R., Toomey, T., Miner, K., Wagenaar, AC., Wolfson, M., & Poon, R. (1997). Why and in what context adolescents obtain alcohol from adults: A pilot study. *Substance Use and Misuse*, 32(2), 219-228.
- Jessor, R., Chase, J. A., & Donovan, J. E. (1980). Psychosocial correlates of marijuana use and problem drinking in a national sample of adolescents. *American Journal of Public Health*, 70(6), 604-613.
- Kandel, D., & Yamaguchi, K. (1993). From beer to crack: Developmental patterns of drug involvement. *American Journal of Public Health*, 83(6), 851-855.
- Knight, J. R., Shrier, L. A., Bravender, T. D., Farrell, M., Vander Bilt, J., & Shaffer, H. J. (1999). A new brief screen for adolescent substance abuse. *Archives of Pediatrics & Adolescent Medicine* 153, 591-596.

- Knight, J. R., Sherritt, L., Shrier, L. A., Harris, S. K., & Chang, G. (2002). Validity of the CRAFFT substance abuse screening test among adolescent clinic patients. *Archives Of Pediatrics & Adolescent Medicine*, 156 (6), 607-614.
- Knight, J. R., Sherritt, L., Harris, K. S., Gates, E. C., & Chang, G. (2003). Validity of brief alcohol screening tests among adolescents: a comparison of the AUDIT, POSIT, CAGE, and CRAFFT. *Alcoholism: Clinical And Experimental Research*, 27 (1), 67-73.
- Kuntsche, E., Kuendig, H., & Gmel, G. (2008). Alcohol outlet density, perceived availability and adolescent alcohol use: A multilevel structural equation model. *Journal of Epidemiology and Community Health*, 62, 811-816.
- Lee, C. Y., Eyre, S. L., & Millstein, S. G. (2000). Gender and ethnic differences in adolescent alcohol use: Problem behavior theory approach. *Journal of Adolescent Health*, 26(2), 143.
- Levy, P. S., & Lemeshow, S. (2003). *Sampling of populations: Methods and applications* (3rd ed.). New York, NY: John Wiley & Sons.
- Lippman-Kredam S., & Grube, J. W. (2009). Students' perception of community disapproval, perceived enforcement of school antismoking policies, personal beliefs, and their cigarette smoking behaviors: Results from a structural equation analysis. *Nicotine and Tobacco Research*, 11(5), 531-539.
- McMaster, L. E., & Wintre, M. G. (1996). Relations between perceived parental reciprocity, perceived parental approval, and adolescent substance use. *Journal of Adolescent Research*, 11(4), 440-460.
- Millstein, S. G., & Halpern-Felsher, B. L. (2002). Perceptions of risk and vulnerability. *The Journal of Adolescent Health*, 31(S), 10-27.
- Nunnally, J. C. (1978). *Psychometric theory* (2nd ed.). New York, NY: McGraw Hill.
- Riggs, N. R., & Greenberg, M. T. (2004). After-school youth development programs: A developmental-ecological model of current research. *Clinical Child & Family Psychology Review*, 7(3), 177-190.
- Stinchfield, R., Kushner, M. G., & Winters, K. C. (2005). Alcohol use and prior substance abuse treatment in relation to gambling problem severity and gambling treatment outcome. *Journal of Gambling Studies*, 21(3), 273-297.
- Substance Abuse and Mental Health Services Administration. (2007). Results from the 2006 National Survey on Drug Use and Health: National Findings (Office of Applied Studies, NSDUH Series H-32, DHHS Publication No. SMA 07-4293). Rockville, MD. Estimates for abuse and dependence by age retrieved July 9, 2008 from <http://www.oas.samhsa.gov/NSDUH/2k6NSDUH/tabs/Sect5peTabs1to56.htm#Tab5.2B>.
- Substance Abuse and Mental Health Services Administration. (2008). National Outcome Measures. Retrieved July 17, 2008 from http://www.nationaloutcomemeasures.samhsa.gov/outcome/index_2007.asp.
- Substance Abuse and Mental Health Services Administration. (2009). Communities That Care Youth Survey. Retrieved October 15, 2009 from http://download.ncadi.samhsa.gov/Prevline/pdfs/ctc/CTC_Youth_Survey_2006.pdf.
- Wagenaar, A. C., & Toomey, T. L. (1996). Sources of alcohol for underage drinkers. *Journal of Studies on Alcohol*, 57(3), 325-333.
- Warner, L. A., & White, H. R. (2003). Longitudinal effects of age at onset and first drinking

situations on problem drinking. *Substance Use & Misuse*, 38(14), 1983-2016.

Wild, T. C., Hinson, R., Cunningham, J., & Bacchiochi, J. (2001). Perceived vulnerability to alcohol-related harm in young adults: Independent effects of risky alcohol use and drinking motives. *Experimental and Clinical Psychopharmacology*, 9(1), 117-125.

Winters, K. C., & Anderson, N. (2000). Gambling involvement and drug use among adolescents. *Journal Of Gambling Studies*, 16(2-3), 175-198.

Wodarski, J. S. (1988). Teams-Games-Tournaments: Teaching adolescents about alcohol and driving. *Journal of Alcohol and Drug Education*, 33(3), 46-57. ■



II. STATEWIDE PREVALENCE OF USE

Statewide Prevalence Rates by Grade: Lifetime and Monthly Use

| | |
|------------------|----|
| • Grade 6 | 62 |
| • Grade 7 | 64 |
| • Grade 8 | 66 |
| • Grade 9 | 68 |
| • Grade 10 | 70 |
| • Grade 11 | 72 |
| • Grade 12 | 74 |

| | |
|--|-----------|
| Mean Age of First Time Use of Alcohol, Tobacco, and Other Drugs | 76 |
|--|-----------|

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 29.2 | 28.8 | 27.6 | 27.7 | 27.3 | 26.3 | 23.3 | 21.8 | 20.6 | 17.7 | 17.6 | 17.7 | 14.6 | 11.9 | 11.1 | 9.8 | 8.9 | 8.1 | 7.3 | -0.8 |
| Smokeless tobacco | 8.5 | 9.0 | 7.7 | 7.1 | 6.7 | 4.6 | 3.7 | 4.0 | 3.5 | 2.9 | 3.2 | 3.0 | 2.9 | 2.7 | 2.4 | 2.5 | 2.0 | 2.8 | 2.5 | -0.3 |
| Cigars | -- | -- | -- | -- | -- | 9.7 | 8.4 | 7.7 | 10.5 | 6.9 | 6.8 | 7.0 | 6.0 | 5.4 | 4.8 | 4.5 | 3.6 | 2.2 | 1.9 | -0.3 |
| Pipe | -- | -- | -- | -- | -- | 3.2 | 3.1 | 2.9 | -- | 2.1 | 2.2 | 2.4 | 2.0 | 1.8 | 1.3 | 1.2 | 1.0 | 1.3 | 1.3 | 1.3 |
| Alcohol | 37.8 | 36.1 | 32.4 | 29.3 | 32.3 | 25.7 | 26.9 | 25.8 | 29.6 | 29.4 | 29.4 | 29.7 | 26.1 | 25.2 | 22.5 | 21.2 | 19.2 | 18.3 | 17.4 | -0.9 |
| Marijuana | 2.2 | 2.9 | 3.6 | 5.1 | 4.7 | 5.0 | 4.2 | 3.8 | 4.5 | 4.8 | 4.6 | 4.5 | 3.9 | 2.9 | 3.0 | 2.5 | 2.4 | 2.8 | 3.1 | 0.3 |
| Cocaine | 0.1 | 0.5 | 0.6 | 0.8 | 0.8 | 0.9 | 0.7 | 0.7 | 0.8 | 0.7 | 0.8 | 0.7 | 0.8 | 0.6 | 0.5 | 0.4 | 0.4 | 0.6 | 0.6 | 0.6 |
| Crack | 0.1 | 0.6 | 0.6 | 1.0 | 0.9 | 1.0 | 0.9 | 0.8 | 1.1 | 1.0 | 1.0 | 1.0 | 1.1 | 0.8 | 0.7 | 0.5 | 0.6 | 0.6 | 0.6 | 0.6 |
| Inhalants | 7.8 | 8.4 | 8.1 | 8.1 | 10.1 | 7.9 | 7.3 | 6.4 | 6.9 | 6.1 | 7.2 | 8.1 | 7.3 | 6.5 | 6.2 | 5.6 | 4.8 | 1.9 | 1.8 | 1.8 |
| Amphetamines | 1.5 | 1.6 | 1.3 | 1.5 | 1.5 | 1.3 | 0.9 | 0.9 | 1.0 | 0.8 | 0.7 | 0.7 | 0.6 | 0.5 | 0.4 | 0.4 | 0.3 | 0.5 | 0.5 | 0.4 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.7 | 0.5 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | 0.5 |
| Heroin | 0.1 | 0.5 | 0.4 | 0.8 | 0.7 | 0.9 | 0.6 | 0.6 | 0.8 | 0.7 | 0.7 | 0.7 | 0.6 | 0.5 | 0.4 | 0.3 | 0.4 | 0.5 | 0.4 | 0.4 |
| Steroids | 1.2 | 1.5 | 1.0 | 1.4 | 1.4 | 1.3 | 1.4 | 1.2 | 1.1 | 0.9 | 1.0 | 1.1 | 0.9 | 0.8 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | 0.6 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.7 | 0.6 | 0.7 | 0.5 | 0.4 | 0.3 | 0.2 | 0.2 | 0.5 | 0.4 | 0.4 |
| Tranquilizers | 4.3 | 4.4 | 4.0 | 4.2 | 5.1 | 4.7 | 4.1 | 3.7 | 4.0 | 4.3 | 4.4 | 4.4 | 4.1 | 3.9 | 4.0 | 3.6 | 3.4 | 0.5 | 0.5 | 0.5 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.8 | 0.7 | 0.7 | 0.5 | 0.5 | 0.3 | 0.3 | 0.3 | 0.4 | 0.4 | 0.4 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.2 | 1.1 | 1.1 |
| Prescription drugs | -- | -- | -- | -- | -- | 1.4 | 1.0 | 1.2 | 1.3 | 1.1 | 1.1 | 1.1 | 1.1 | 0.9 | 1.0 | 0.9 | 0.9 | 1.1 | 1.0 | 1.0 |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 4.7 | 4.1 | 3.6 | 1.4 | 1.1 | -0.3 |

NOTES: -- Data not available
* Statistically significant changes between 2010 and 2011 prevalence rates (p<.05).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* | |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|------|
| Cigarettes | 8.5 | 10.0 | 9.3 | 9.7 | 9.7 | 9.5 | 8.2 | 7.0 | 5.7 | 6.0 | 4.8 | 5.1 | 5.4 | 4.2 | 3.3 | 2.8 | 2.6 | 2.3 | 3.3 | 2.6 | -0.7 |
| Smokeless tobacco | 3.5 | 4.0 | 3.3 | 3.1 | 3.1 | 3.1 | 1.9 | 1.4 | 1.4 | 1.4 | 1.2 | 1.4 | 1.3 | 1.1 | 1.1 | 0.9 | 1.0 | 0.8 | 1.2 | 1.0 | -0.2 |
| Cigars | -- | -- | -- | -- | -- | -- | 3.9 | 3.3 | 2.9 | 3.8 | 2.4 | 2.5 | 2.5 | 2.2 | 1.8 | 1.6 | 1.5 | 1.2 | 0.9 | 0.7 | -0.2 |
| Pipe | -- | -- | -- | -- | -- | -- | 1.5 | 1.0 | 0.8 | -- | 0.8 | 0.9 | 0.8 | 0.7 | 0.7 | 0.5 | 0.6 | 0.5 | 0.7 | 0.6 | |
| Alcohol | 13.6 | 13.7 | 11.4 | 10.7 | 11.5 | 9.5 | 9.5 | 9.9 | 9.2 | 9.5 | 9.5 | 9.9 | 9.8 | 8.1 | 7.6 | 6.9 | 6.3 | 5.6 | 6.2 | 5.3 | -0.9 |
| Binge drinking# | 8.0 | 8.1 | 7.2 | 7.6 | 7.6 | 7.6 | 8.7 | 8.2 | 7.9 | 6.7 | 6.6 | 5.3 | 5.9 | 4.5 | 4.2 | 4.9 | 4.2 | 4.2 | 5.5 | 5.8 | |
| Marijuana | 1.2 | 1.7 | 1.9 | 3.0 | 2.7 | 2.2 | 2.2 | 2.2 | 1.8 | 2.4 | 2.2 | 2.3 | 2.5 | 1.8 | 1.4 | 1.4 | 1.3 | 1.1 | 1.5 | 1.6 | |
| Cocaine | 0.1 | 0.3 | 0.3 | 0.6 | 0.5 | 0.5 | 0.3 | 0.4 | 0.4 | 0.4 | 0.3 | 0.4 | 0.4 | 0.4 | 0.2 | 0.2 | 0.2 | 0.2 | 0.4 | 0.3 | |
| Crack | 0.1 | 0.3 | 0.3 | 0.6 | 0.5 | 0.6 | 0.3 | 0.4 | 0.4 | 0.4 | 0.4 | 0.5 | 0.5 | 0.5 | 0.3 | 0.3 | 0.3 | 0.2 | 0.4 | 0.3 | |
| Inhalants | 3.8 | 4.1 | 3.6 | 3.8 | 4.9 | 3.6 | 3.4 | 2.7 | 3.1 | 2.5 | 3.2 | 3.6 | 3.1 | 2.9 | 2.3 | 2.3 | 2.3 | 1.8 | 1.0 | 0.9 | |
| Amphetamines | 0.1 | 0.9 | 0.6 | 1.0 | 0.9 | 0.6 | 0.5 | 0.3 | 0.4 | 0.4 | 0.4 | 0.3 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.2 | 0.3 | 0.2 | |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.3 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.3 | |
| Heroin | 0.1 | 0.3 | 0.2 | 0.5 | 0.4 | 0.4 | 0.3 | 0.3 | 0.3 | 0.4 | 0.3 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.2 | 0.1 | 0.3 | 0.2 | |
| Steroids | 0.1 | 0.6 | 0.4 | 0.7 | 0.7 | 0.6 | 0.5 | 0.4 | 0.5 | 0.4 | 0.4 | 0.4 | 0.5 | 0.4 | 0.3 | 0.3 | 0.2 | 0.2 | 0.3 | 0.2 | |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.4 | 0.3 | 0.3 | 0.3 | 0.1 | 0.1 | 0.1 | 0.1 | 0.3 | 0.2 | |
| Tranquilizers | 2.0 | 2.2 | 1.6 | 1.9 | 2.4 | 2.0 | 1.8 | 1.6 | 1.7 | 1.7 | 1.7 | 1.8 | 1.9 | 1.7 | 1.5 | 1.5 | 1.4 | 1.3 | 0.3 | 0.3 | |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.4 | 0.3 | 0.3 | 0.3 | 0.2 | 0.1 | 0.2 | 0.1 | 0.2 | 0.3 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.7 | 0.6 | |
| Prescription drugs | -- | -- | -- | -- | -- | -- | 0.7 | 0.5 | 0.5 | 0.6 | 0.4 | 0.4 | 0.5 | 0.4 | 0.3 | 0.4 | 0.4 | 0.3 | 0.7 | 0.6 | |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.4 | 2.3 | 1.9 | 0.8 | 0.7 | |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 38.6 | 40.1 | 39.8 | 40.2 | 38.5 | 35.1 | 34.0 | 31.2 | 29.1 | 26.3 | 23.4 | 24.6 | 22.0 | 19.7 | 17.9 | 16.4 | 16.1 | 13.9 | 13.2 | -0.7 |
| Smokeless tobacco | 13.9 | 14.9 | 14.6 | 12.7 | 11.5 | 8.1 | 7.3 | 5.9 | 5.2 | 5.3 | 4.8 | 6.1 | 4.8 | 5.0 | 4.3 | 4.4 | 4.9 | 4.9 | 5.1 | |
| Cigars | -- | -- | -- | -- | -- | 15.1 | 14.3 | 12.3 | 14.9 | 11.5 | 9.8 | 10.6 | 9.6 | 8.3 | 8.6 | 8.3 | 8.1 | 4.5 | 4.3 | |
| Pipe | -- | -- | -- | -- | -- | 5.1 | 5.0 | 5.0 | -- | 3.6 | 3.6 | 3.8 | 3.2 | 2.9 | 3.0 | 2.8 | 2.7 | 2.8 | 3.2 | 0.4 |
| Alcohol | 48.7 | 47.4 | 44.4 | 44.0 | 42.4 | 36.2 | 37.5 | 36.5 | 39.5 | 39.4 | 37.7 | 38.3 | 34.0 | 33.3 | 32.7 | 29.8 | 29.3 | 28.0 | 26.1 | -1.9 |
| Marijuana | 5.3 | 7.7 | 9.5 | 12.8 | 10.2 | 9.9 | 9.4 | 8.8 | 9.8 | 9.3 | 8.7 | 8.7 | 8.0 | 6.8 | 7.7 | 6.4 | 6.9 | 6.8 | 7.3 | 0.5 |
| Cocaine | 1.0 | 1.2 | 1.3 | 2.1 | 1.6 | 1.6 | 1.6 | 1.2 | 1.2 | 1.5 | 1.6 | 1.6 | 1.3 | 1.5 | 1.2 | 1.2 | 1.3 | 1.0 | 1.1 | |
| Crack | 0.1 | 1.3 | 1.3 | 1.9 | 1.4 | 1.7 | 1.6 | 1.4 | 1.5 | 1.8 | 1.6 | 1.9 | 1.6 | 1.5 | 1.3 | 1.3 | 1.4 | 1.0 | 1.1 | |
| Inhalants | 9.8 | 12.0 | 12.3 | 13.5 | 12.9 | 10.6 | 8.8 | 7.9 | 7.8 | 7.6 | 8.8 | 10.3 | 8.9 | 9.0 | 7.9 | 7.5 | 7.2 | 3.6 | 3.2 | -0.4 |
| Amphetamines | 3.9 | 4.5 | 4.6 | 5.7 | 3.5 | 3.0 | 2.5 | 2.4 | 2.1 | 2.4 | 2.1 | 2.0 | 1.5 | 1.4 | 1.2 | 1.1 | 1.2 | 0.9 | 1.0 | |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.2 | 1.2 | 1.0 | 0.9 | 0.9 | 0.9 | 1.0 | |
| Heroin | 1.0 | 1.2 | 1.1 | 1.2 | 1.3 | 1.1 | 1.0 | 0.9 | 1.0 | 1.1 | 1.2 | 1.2 | 1.0 | 1.1 | 0.9 | 0.9 | 0.9 | 0.8 | 0.8 | |
| Steroids | 2.2 | 2.4 | 1.7 | 2.0 | 1.8 | 1.6 | 2.1 | 1.5 | 1.3 | 1.6 | 1.5 | 1.6 | 1.3 | 1.2 | 1.2 | 1.1 | 1.2 | 0.8 | 1.0 | |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.8 | 1.8 | 1.5 | 1.2 | 1.2 | 1.0 | 0.9 | 0.9 | 1.0 | 1.1 | |
| Tranquilizers | 6.5 | 6.8 | 6.7 | 8.5 | 7.4 | 7.1 | 6.2 | 6.4 | 6.0 | 6.8 | 6.5 | 7.2 | 6.0 | 6.0 | 5.6 | 5.5 | 5.9 | 0.9 | 0.9 | |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.7 | 1.6 | 1.5 | 1.2 | 1.1 | 1.0 | 1.0 | 1.0 | 0.9 | 0.9 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.6 | 2.7 |
| Prescription drugs | -- | -- | -- | -- | -- | 2.5 | 2.1 | 2.4 | 1.9 | 2.0 | 1.9 | 1.9 | 1.9 | 1.8 | 2.2 | 1.9 | 1.9 | 2.4 | 2.4 | |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.5 | 6.4 | 6.1 | 2.7 | 2.8 | |

NOTES: -- Data not available
* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 14.5 | 16.9 | 17.7 | 19.0 | 16.3 | 14.0 | 12.3 | 11.0 | 9.9 | 9.1 | 8.1 | 8.9 | 7.5 | 6.7 | 6.2 | 5.4 | 5.6 | 6.4 | 6.0 | |
| Smokeless tobacco | 6.7 | 7.5 | 7.1 | 6.7 | 5.2 | 3.9 | 2.7 | 2.2 | 2.1 | 2.3 | 2.1 | 2.8 | 2.2 | 2.3 | 2.0 | 2.0 | 2.3 | 2.1 | 2.2 | |
| Cigars | -- | -- | -- | -- | -- | 6.9 | 5.8 | 5.3 | 6.0 | 4.6 | 3.7 | 4.5 | 3.7 | 3.2 | 3.6 | 3.3 | 3.4 | 1.9 | 1.6 | -0.3 |
| Pipe | -- | -- | -- | -- | -- | 2.4 | 1.8 | 2.1 | -- | 1.5 | 1.5 | 1.6 | 1.2 | 1.2 | 1.5 | 1.3 | 1.4 | 1.6 | 1.5 | |
| Alcohol | 21.8 | 21.5 | 19.6 | 21.3 | 18.4 | 16.8 | 16.8 | 16.5 | 16.1 | 15.2 | 14.7 | 15.3 | 12.7 | 12.9 | 12.2 | 11.2 | 10.8 | 11.6 | 10.2 | -1.4 |
| Binge drinking# | 12.7 | 13.0 | 12.3 | 13.5 | 12.0 | 12.3 | 12.4 | 12.4 | 9.7 | 10.0 | 8.0 | 8.9 | 6.8 | 7.0 | 8.3 | 7.5 | 7.7 | 7.7 | 7.9 | |
| Marijuana | 2.7 | 4.8 | 5.4 | 7.9 | 5.7 | 5.5 | 5.1 | 5.2 | 5.2 | 4.7 | 4.6 | 4.5 | 4.1 | 3.6 | 3.9 | 3.2 | 3.5 | 3.8 | 4.1 | |
| Cocaine | 0.1 | 0.8 | 0.7 | 1.3 | 1.0 | 0.9 | 0.7 | 0.7 | 0.6 | 0.8 | 0.9 | 0.8 | 0.7 | 0.8 | 0.7 | 0.6 | 0.6 | 0.6 | 0.6 | |
| Crack | 0.1 | 0.7 | 0.7 | 1.2 | 0.9 | 0.9 | 0.8 | 0.7 | 0.7 | 0.7 | 0.9 | 0.9 | 0.8 | 0.8 | 0.6 | 0.6 | 0.7 | 0.6 | 0.7 | |
| Inhalants | 4.7 | 6.1 | 5.6 | 7.0 | 6.6 | 4.8 | 3.7 | 3.3 | 3.1 | 3.0 | 3.7 | 4.8 | 3.7 | 3.6 | 3.1 | 3.0 | 2.7 | 1.7 | 1.4 | -0.3 |
| Amphetamines | 2.2 | 2.7 | 2.4 | 3.3 | 2.3 | 1.7 | 1.2 | 1.2 | 1.1 | 1.1 | 1.0 | 1.1 | 0.7 | 0.8 | 0.6 | 0.6 | 0.6 | 0.5 | 0.5 | |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.6 | 0.6 | 0.5 | 0.4 | 0.5 | 0.5 | 0.5 | |
| Heroin | 0.1 | 0.8 | 0.6 | 0.9 | 0.8 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.6 | 0.6 | 0.5 | 0.5 | 0.4 | 0.5 | 0.4 | 0.4 | 0.4 | |
| Steroids | 0.1 | 1.3 | 0.9 | 1.0 | 0.8 | 0.6 | 1.0 | 0.7 | 0.7 | 0.8 | 0.7 | 0.8 | 0.6 | 0.6 | 0.6 | 0.5 | 0.5 | 0.5 | 0.5 | |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.0 | 0.9 | 0.8 | 0.6 | 0.6 | 0.6 | 0.4 | 0.4 | 0.5 | 0.5 | |
| Tranquilizers | 3.1 | 3.7 | 3.3 | 5.0 | 4.1 | 4.0 | 3.2 | 3.3 | 3.0 | 3.2 | 2.9 | 3.3 | 2.7 | 2.8 | 2.5 | 2.3 | 2.6 | 0.5 | 0.5 | |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 0.8 | 0.8 | 0.8 | 0.7 | 0.5 | 0.5 | 0.4 | 0.5 | 0.4 | 0.5 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.5 | 1.5 | |
| Prescription drugs | -- | -- | -- | -- | -- | 1.4 | 1.0 | 1.2 | 0.8 | 0.8 | 0.9 | 0.8 | 0.8 | 0.8 | 1.0 | 0.8 | 0.8 | 1.5 | 1.4 | |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 3.4 | 3.5 | 3.0 | 1.6 | 1.6 | |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* 2010 | National ^{###} Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--------------|------------------------------------|
| Cigarettes | 50.0 | 51.3 | 52.4 | 53.7 | 52.6 | 49.5 | 45.7 | 43.7 | 39.9 | 37.4 | 35.1 | 34.7 | 32.1 | 30.2 | 28.1 | 25.5 | 25.0 | 22.4 | 20.6 | -1.8 | 20.0 |
| Smokeless tobacco | 18.9 | 20.1 | 19.2 | 17.9 | 16.7 | 12.4 | 10.9 | 10.9 | 8.2 | 7.7 | 7.3 | 8.3 | 7.3 | 8.3 | 7.2 | 7.9 | 8.4 | 9.1 | 8.6 | -0.5 | 9.9 |
| Cigars | -- | -- | -- | -- | -- | 24.4 | 22.5 | 21.3 | 22.1 | 17.4 | 16.4 | 16.1 | 15.8 | 14.6 | 14.7 | 14.2 | 13.7 | 9.7 | 8.1 | -1.6 | -- |
| Pipe | -- | -- | -- | -- | -- | 8.0 | 8.6 | 7.8 | -- | 5.5 | 5.5 | 5.6 | 5.2 | 4.9 | 5.8 | 5.5 | 5.6 | 6.3 | 6.0 | -- | -- |
| Alcohol | 60.9 | 61.7 | 60.2 | 59.1 | 59.4 | 51.5 | 53.0 | 51.0 | 53.3 | 52.6 | 51.7 | 52.0 | 49.0 | 46.5 | 45.4 | 44.0 | 41.6 | 41.7 | 37.7 | -4.0 | 35.8 |
| Marijuana | 12.0 | 16.0 | 20.2 | 24.3 | 23.0 | 22.1 | 19.7 | 19.0 | 19.5 | 20.0 | 19.1 | 18.6 | 17.6 | 15.6 | 16.1 | 14.4 | 15.0 | 15.3 | 14.6 | -0.7 | 17.3 |
| Cocaine | 2.0 | 2.5 | 2.9 | 3.6 | 3.5 | 3.2 | 2.6 | 2.6 | 2.7 | 2.6 | 2.9 | 2.7 | 2.9 | 2.5 | 2.4 | 2.1 | 2.2 | 2.0 | 2.0 | -- | 2.6 |
| Crack | 1.5 | 2.0 | 2.4 | 3.0 | 2.9 | 2.9 | 2.4 | 2.4 | 2.5 | 2.3 | 2.7 | 2.4 | 2.6 | 2.2 | 2.0 | 1.8 | 1.9 | 1.7 | 1.7 | -- | 1.5 |
| Inhalants | 14.5 | 15.3 | 18.1 | 17.4 | 17.6 | 13.9 | 12.0 | 11.2 | 10.3 | 9.3 | 10.7 | 11.0 | 11.3 | 10.4 | 10.5 | 9.5 | 9.0 | 6.5 | 5.4 | -1.1 | 14.5 |
| Amphetamines | 9.0 | 9.1 | 10.6 | 10.0 | 9.1 | 6.6 | 5.8 | 5.6 | 5.5 | 5.0 | 4.7 | 4.3 | 3.7 | 3.2 | 3.0 | 2.6 | 2.7 | 2.5 | 2.3 | -- | 5.7 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.4 | 2.0 | 1.6 | 1.5 | 1.5 | 1.5 | 1.5 | -- | 1.8 |
| Heroin | 1.4 | 1.7 | 1.8 | 2.2 | 1.8 | 1.9 | 1.3 | 1.6 | 1.7 | 1.4 | 1.8 | 1.5 | 1.6 | 1.5 | 1.3 | 1.4 | 1.5 | 1.3 | 1.2 | -- | 1.3 |
| Steroids | 2.6 | 2.7 | 2.6 | 2.1 | 2.3 | 2.5 | 2.1 | 2.5 | 2.2 | 2.1 | 2.2 | 2.2 | 1.8 | 1.5 | 1.5 | 1.5 | 1.4 | 1.3 | 1.3 | -- | 1.1 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 3.4 | 3.3 | 2.7 | 2.5 | 2.3 | 2.3 | 2.0 | 2.1 | 2.3 | 2.2 | -- | 3.3 |
| Tranquilizers | 10.3 | 10.7 | 11.4 | 11.9 | 12.5 | 11.5 | 10.1 | 10.3 | 10.2 | 10.2 | 10.7 | 10.2 | 9.8 | 8.9 | 9.1 | 8.6 | 8.5 | 2.1 | 1.9 | -- | 4.4 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 3.8 | 3.2 | 2.6 | 2.7 | 2.3 | 2.2 | 2.0 | 2.1 | 2.3 | 2.3 | -- | 3.4 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.3 | 5.6 | -0.7 | -- |
| Prescription drugs | -- | -- | -- | -- | -- | 4.2 | 3.9 | 4.2 | 4.0 | 3.5 | 3.6 | 3.1 | 3.5 | 3.1 | 4.5 | 4.1 | 4.2 | 5.3 | 4.8 | -0.5 | -- |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 10.2 | 9.7 | 9.4 | 5.4 | 5.0 | -0.4 | -- |

NOTES: -- Data not available
 * Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).
 ### Monitoring the Future Study, the University of Michigan, 2011

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* 2010 | National ^{###} Change* 2010 |
|-----------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--------------|---|
| Cigarettes | 21.1 | 24.2 | 26.3 | 27.1 | 25.8 | 24.1 | 19.7 | 18.2 | 16.1 | 14.7 | 14.0 | 13.4 | 12.5 | 11.6 | 10.8 | 9.7 | 9.9 | 10.8 | 10.0 | -0.8 | 7.1 |
| Smokeless tobacco | 9.0 | 10.6 | 9.3 | 8.6 | 7.2 | 5.6 | 4.6 | 4.5 | 3.5 | 3.4 | 3.2 | 3.8 | 3.3 | 3.8 | 3.3 | 3.7 | 4.1 | 3.9 | 3.9 | 4.1 | 4.1 |
| Cigars | -- | -- | -- | -- | -- | -- | 12.0 | 9.6 | 9.7 | 10.0 | 6.6 | 6.8 | 6.7 | 6.5 | 5.9 | 6.4 | 6.2 | 5.9 | 3.9 | 3.2 | -0.7 |
| Pipe | -- | -- | -- | -- | -- | -- | 3.5 | 3.1 | -- | 2.2 | 2.2 | 2.3 | 2.1 | 2.0 | 3.1 | 2.7 | 2.8 | 3.1 | 3.0 | 3.0 | -- |
| Alcohol | 30.9 | 32.0 | 31.3 | 31.3 | 30.3 | 27.2 | 27.7 | 26.4 | 24.7 | 24.9 | 24.3 | 23.5 | 21.1 | 20.9 | 19.9 | 18.8 | 17.5 | 19.3 | 16.7 | -2.6 | 13.8 |
| Binge drinking [#] | 18.5 | 18.6 | 18.4 | 19.0 | 18.9 | 19.7 | 19.5 | 18.6 | 14.8 | 15.5 | 13.4 | 13.4 | 11.6 | 11.0 | 13.2 | 12.2 | 12.5 | 12.0 | 11.1 | -0.9 | 7.2 |
| Marijuana | 6.5 | 9.9 | 13.1 | 15.2 | 13.8 | 12.8 | 11.3 | 11.3 | 11.3 | 11.1 | 10.6 | 9.8 | 9.3 | 8.2 | 8.3 | 7.1 | 7.8 | 8.9 | 8.3 | -0.6 | 8.0 |
| Cocaine | 0.1 | 1.3 | 1.4 | 1.7 | 1.9 | 1.5 | 1.1 | 1.1 | 1.3 | 1.2 | 1.3 | 1.2 | 1.4 | 1.1 | 1.1 | 0.9 | 0.9 | 1.1 | 1.0 | 0.6 | 0.6 |
| Crack | 0.1 | 1.0 | 1.1 | 1.4 | 1.4 | 1.3 | 1.1 | 1.0 | 1.2 | 0.9 | 1.2 | 1.0 | 1.2 | 1.0 | 0.9 | 0.8 | 0.7 | 0.9 | 0.9 | 0.4 | 0.4 |
| Inhalants | 6.5 | 7.5 | 8.4 | 8.0 | 8.2 | 5.7 | 4.8 | 4.4 | 4.0 | 3.4 | 4.2 | 4.7 | 4.3 | 4.1 | 3.7 | 3.3 | 3.2 | 2.7 | 2.3 | -0.4 | 3.6 |
| Amphetamines | 5.1 | 5.2 | 5.9 | 5.6 | 5.3 | 3.3 | 2.8 | 2.8 | 2.8 | 2.4 | 2.4 | 2.1 | 1.9 | 1.4 | 1.4 | 1.2 | 1.2 | 1.2 | 1.2 | 1.8 | 1.8 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.1 | 0.9 | 0.7 | 0.7 | 0.6 | 0.8 | 0.9 | 0.7 | 0.7 |
| Heroin | 0.1 | 1.0 | 0.9 | 1.1 | 1.1 | 1.0 | 0.6 | 0.7 | 0.8 | 0.7 | 0.9 | 0.8 | 0.8 | 0.7 | 0.6 | 0.6 | 0.6 | 0.7 | 0.7 | 0.4 | 0.4 |
| Steroids | 1.2 | 1.3 | 1.0 | 1.0 | 1.0 | 1.1 | 0.9 | 0.9 | 1.0 | 0.9 | 1.0 | 0.9 | 0.8 | 0.7 | 0.7 | 0.6 | 0.7 | 0.7 | 0.7 | 0.3 | 0.3 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.6 | 1.5 | 1.3 | 1.1 | 1.0 | 0.9 | 0.8 | 0.9 | 1.1 | 1.1 | 1.1 | 1.1 |
| Tranquilizers | 5.6 | 5.7 | 5.8 | 6.0 | 6.5 | 5.7 | 5.2 | 5.1 | 5.2 | 4.8 | 5.1 | 4.9 | 4.7 | 4.1 | 3.9 | 3.7 | 3.6 | 1.1 | 1.0 | 1.2 | 1.2 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.5 | 1.3 | 1.2 | 1.2 | 0.9 | 0.9 | 0.9 | 0.9 | 1.1 | 1.2 | 1.0 | 1.0 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 3.4 | 3.0 | -0.4 | -- |
| Prescription drugs | -- | -- | -- | -- | -- | -- | 2.0 | 1.7 | 2.0 | 1.7 | 1.3 | 1.4 | 1.3 | 1.5 | 1.3 | 1.9 | 1.7 | 1.8 | 3.0 | 2.7 | -0.3 |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 5.2 | 5.1 | 4.9 | 2.9 | 2.7 | -- | -- |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Monitoring the Future Study, the University of Michigan, 2011

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 56.5 | 57.4 | 58.7 | 61.4 | 57.3 | 55.8 | 53.7 | 50.6 | 47.2 | 43.6 | 39.8 | 40.7 | 36.8 | 36.3 | 33.5 | 32.0 | 31.1 | 28.6 | 27.2 | -1.4 |
| Smokeless tobacco | 24.3 | 25.8 | 26.2 | 24.2 | 21.3 | 18.7 | 15.0 | 14.2 | 11.9 | 12.4 | 10.5 | 12.2 | 10.7 | 11.6 | 10.8 | 11.6 | 12.0 | 13.7 | 13.1 | |
| Cigars | -- | -- | -- | -- | -- | 31.0 | 28.7 | 26.8 | 29.8 | 23.1 | 20.7 | 21.3 | 20.6 | 20.4 | 19.4 | 19.3 | 19.0 | 15.6 | 14.9 | |
| Pipe | -- | -- | -- | -- | -- | 10.9 | 10.1 | 9.7 | -- | 7.6 | 7.1 | 7.3 | 6.5 | 6.9 | 9.1 | 9.1 | 9.3 | 10.4 | 10.4 | |
| Alcohol | 70.2 | 69.9 | 68.7 | 69.7 | 65.9 | 61.8 | 61.4 | 59.8 | 62.6 | 59.9 | 58.4 | 58.4 | 55.2 | 55.5 | 51.7 | 50.6 | 49.7 | 48.8 | 47.7 | -1.1 |
| Marijuana | 18.4 | 23.9 | 26.1 | 35.0 | 28.9 | 30.7 | 27.8 | 26.7 | 28.4 | 28.1 | 25.2 | 24.0 | 22.9 | 22.5 | 21.2 | 21.3 | 21.0 | 22.4 | 23.0 | |
| Cocaine | 2.5 | 3.7 | 3.7 | 4.9 | 4.8 | 5.3 | 4.2 | 3.9 | 3.8 | 4.3 | 4.3 | 4.7 | 4.2 | 4.4 | 3.7 | 3.6 | 3.1 | 3.0 | 2.9 | |
| Crack | 1.8 | 2.5 | 2.7 | 4.0 | 3.5 | 4.2 | 3.3 | 3.0 | 3.0 | 3.1 | 3.2 | 3.3 | 3.0 | 3.0 | 2.4 | 2.5 | 2.2 | 2.2 | 1.9 | |
| Inhalants | 13.3 | 17.4 | 17.3 | 18.3 | 16.7 | 15.7 | 12.3 | 11.6 | 11.1 | 9.9 | 9.8 | 11.3 | 10.9 | 11.4 | 9.8 | 9.7 | 9.1 | 8.2 | 6.9 | -1.3 |
| Amphetamines | 13.5 | 14.8 | 14.3 | 16.7 | 13.3 | 12.0 | 9.9 | 10.1 | 10.0 | 9.7 | 8.6 | 8.6 | 7.0 | 6.8 | 5.3 | 5.1 | 4.8 | 4.9 | 4.3 | -0.6 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 3.5 | 3.0 | 2.2 | 2.0 | 2.1 | 2.0 | 1.9 | |
| Heroin | 1.5 | 2.0 | 1.9 | 2.4 | 2.1 | 2.1 | 1.7 | 1.6 | 1.8 | 1.8 | 2.0 | 2.0 | 2.1 | 1.9 | 1.5 | 1.8 | 1.8 | 1.8 | 1.6 | |
| Steroids | 2.8 | 3.0 | 2.6 | 2.5 | 2.3 | 2.5 | 2.4 | 2.8 | 2.5 | 2.4 | 2.6 | 2.7 | 2.3 | 1.8 | 1.7 | 1.7 | 1.6 | 1.6 | 1.5 | |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 5.4 | 4.2 | 4.1 | 3.6 | 3.7 | 3.2 | 3.3 | 3.3 | 3.7 | 3.4 | |
| Tranquilizers | 11.1 | 13.8 | 13.4 | 15.3 | 15.0 | 14.9 | 13.7 | 14.0 | 14.2 | 14.4 | 13.2 | 13.6 | 12.3 | 12.4 | 10.9 | 10.4 | 10.2 | 3.6 | 3.3 | -0.3 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.3 | 4.5 | 4.3 | 3.9 | 3.4 | 2.9 | 3.4 | 3.2 | 3.9 | 3.6 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 9.9 | 9.4 | |
| Prescription drugs | -- | -- | -- | -- | -- | 6.8 | 5.0 | 6.2 | 6.3 | 5.3 | 5.1 | 4.9 | 5.4 | 5.4 | 7.3 | 7.4 | 6.9 | 8.8 | 8.2 | -0.6 |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 12.3 | 11.7 | 11.1 | 7.9 | 7.2 | -0.7 |

NOTES: -- Data not available
* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 26.2 | 29.4 | 30.8 | 34.4 | 30.5 | 28.9 | 25.4 | 24.9 | 21.8 | 19.1 | 17.9 | 18.2 | 16.4 | 16.5 | 14.7 | 14.7 | 13.8 | 15.2 | 14.4 | -0.8 |
| Smokeless tobacco | 12.2 | 13.3 | 13.2 | 12.5 | 9.8 | 8.3 | 6.5 | 6.5 | 5.4 | 5.2 | 4.9 | 5.6 | 5.2 | 5.7 | 5.3 | 6.0 | 5.9 | 6.6 | 6.2 | -0.4 |
| Cigars | -- | -- | -- | -- | -- | 15.5 | 13.2 | 12.6 | 14.1 | 9.2 | 8.3 | 9.1 | 8.6 | 8.8 | 8.7 | 8.6 | 8.6 | 6.3 | 6.0 | |
| Pipe | -- | -- | -- | -- | -- | 4.5 | 4.1 | 4.2 | -- | 2.9 | 2.7 | 3.2 | 2.7 | 3.0 | 4.4 | 4.4 | 4.4 | 5.2 | 4.8 | -0.4 |
| Alcohol | 38.2 | 39.6 | 38.5 | 40.8 | 36.0 | 36.7 | 36.2 | 35.3 | 33.2 | 30.5 | 30.8 | 29.4 | 26.6 | 28.5 | 24.8 | 24.5 | 22.1 | 24.4 | 23.8 | |
| Binge drinking# | 23.3 | 23.6 | 23.3 | 25.5 | 23.1 | 25.7 | 24.3 | 24.3 | 19.7 | 19.8 | 17.9 | 17.3 | 14.8 | 15.6 | 16.9 | 16.2 | 15.5 | 15.2 | 15.3 | |
| Marijuana | 10.6 | 15.2 | 15.9 | 22.8 | 17.2 | 18.4 | 16.2 | 15.6 | 16.1 | 14.6 | 13.9 | 12.1 | 11.5 | 11.4 | 10.3 | 10.4 | 10.5 | 12.7 | 13.0 | |
| Cocaine | 0.1 | 1.6 | 1.7 | 2.4 | 2.2 | 2.4 | 1.4 | 1.6 | 1.6 | 1.6 | 1.8 | 2.0 | 1.7 | 1.7 | 1.4 | 1.4 | 1.2 | 1.4 | 1.4 | |
| Crack | 0.1 | 1.4 | 1.3 | 2.1 | 1.5 | 1.9 | 1.2 | 1.3 | 1.2 | 1.1 | 1.2 | 1.4 | 1.3 | 1.1 | 0.9 | 0.9 | 0.9 | 1.1 | 1.0 | |
| Inhalants | 5.2 | 7.3 | 6.9 | 7.2 | 6.1 | 5.8 | 4.4 | 4.2 | 3.7 | 3.0 | 3.0 | 3.9 | 3.6 | 3.6 | 2.9 | 3.0 | 2.6 | 2.8 | 2.2 | -0.6 |
| Amphetamines | 6.4 | 7.8 | 7.2 | 8.9 | 6.8 | 5.9 | 4.8 | 5.1 | 5.2 | 4.3 | 3.6 | 3.8 | 3.0 | 3.0 | 2.0 | 2.1 | 1.8 | 2.1 | 1.9 | |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.5 | 1.2 | 0.8 | 0.9 | 0.8 | 0.9 | 0.9 | |
| Heroin | 0.1 | 1.1 | 0.9 | 1.3 | 1.0 | 0.9 | 0.6 | 0.7 | 0.8 | 0.7 | 0.9 | 0.9 | 0.9 | 0.8 | 0.7 | 0.8 | 0.8 | 0.9 | 0.8 | |
| Steroids | 1.3 | 1.4 | 1.2 | 1.2 | 1.1 | 1.1 | 0.9 | 1.0 | 1.1 | 1.0 | 1.2 | 1.1 | 0.9 | 0.9 | 0.8 | 0.8 | 0.8 | 0.9 | 0.8 | -0.1 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.9 | 1.5 | 1.6 | 1.3 | 1.5 | 1.2 | 1.3 | 1.1 | 1.7 | 1.4 | |
| Tranquilizers | 5.2 | 7.1 | 6.2 | 7.7 | 7.9 | 7.5 | 7.2 | 7.4 | 7.3 | 6.7 | 5.9 | 6.3 | 5.1 | 5.7 | 4.4 | 4.3 | 4.0 | 1.7 | 1.6 | |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.0 | 1.5 | 1.6 | 1.4 | 1.3 | 1.1 | 1.3 | 1.1 | 1.7 | 1.5 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 5.0 | 4.7 | |
| Prescription drugs | -- | -- | -- | -- | -- | 3.0 | 2.0 | 2.5 | 2.4 | 1.9 | 1.8 | 1.9 | 2.0 | 2.0 | 2.8 | 3.0 | 2.6 | 4.6 | 4.3 | |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 5.6 | 5.7 | 5.0 | 4.0 | 3.5 | -0.5 |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | National ^{###} Change* 2010 |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---|
| Cigarettes | 60.8 | 61.5 | 61.8 | 63.6 | 64.1 | 61.4 | 60.5 | 57.4 | 55.5 | 51.5 | 47.2 | 46.7 | 43.9 | 43.0 | 40.6 | 39.1 | 37.6 | 34.8 | 32.8 | -2.0 |
| Smokeless tobacco | 27.9 | 28.0 | 28.7 | 30.0 | 30.0 | 26.2 | 22.1 | 19.6 | 17.3 | 16.0 | 13.8 | 14.1 | 14.3 | 15.4 | 15.0 | 15.4 | 15.6 | 17.5 | 17.2 | 16.8 |
| Cigars | -- | -- | -- | -- | -- | -- | 37.5 | 36.4 | 32.4 | 36.4 | 27.2 | 27.8 | 26.9 | 26.8 | 26.7 | 25.9 | 25.0 | 22.2 | 20.9 | -1.3 |
| Pipe | -- | -- | -- | -- | -- | -- | 12.8 | 12.1 | 10.6 | -- | 8.3 | 7.7 | 7.8 | 8.6 | 13.3 | 12.6 | 12.9 | 15.2 | 15.0 | -- |
| Alcohol | 76.5 | 76.2 | 73.2 | 74.3 | 75.2 | 68.4 | 69.2 | 66.6 | 71.3 | 68.4 | 67.2 | 65.5 | 63.5 | 62.3 | 61.0 | 58.7 | 56.9 | 58.5 | 55.5 | -3.0 |
| Marijuana | 25.6 | 30.5 | 34.2 | 40.2 | 40.1 | 38.2 | 37.4 | 34.1 | 37.8 | 36.9 | 34.8 | 33.5 | 31.6 | 30.1 | 29.9 | 28.3 | 29.1 | 30.9 | 30.3 | 33.4 |
| Cocaine | 3.4 | 4.3 | 4.9 | 6.9 | 6.7 | 6.6 | 6.0 | 5.1 | 6.1 | 6.5 | 6.0 | 6.0 | 6.1 | 5.9 | 5.8 | 4.9 | 4.2 | 4.1 | 3.8 | 3.7 |
| Crack | 1.9 | 2.9 | 3.6 | 4.8 | 4.7 | 5.1 | 4.1 | 3.4 | 3.9 | 3.5 | 3.7 | 3.8 | 3.8 | 3.4 | 3.1 | 2.7 | 2.4 | 2.3 | 2.0 | -0.3 |
| Inhalants | 15.0 | 16.1 | 17.5 | 17.8 | 17.5 | 15.9 | 12.8 | 11.4 | 11.9 | 10.7 | 10.9 | 11.3 | 11.2 | 11.1 | 10.6 | 9.9 | 9.2 | 8.7 | 7.4 | -1.3 |
| Amphetamines | 17.3 | 17.2 | 17.7 | 18.8 | 17.6 | 15.1 | 13.5 | 12.2 | 13.7 | 13.2 | 12.3 | 11.3 | 10.2 | 9.0 | 8.2 | 7.3 | 6.7 | 6.6 | 5.9 | -0.7 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 4.5 | 3.5 | 3.0 | 2.5 | 2.3 | 2.2 | 2.2 | 2.5 |
| Heroin | 1.4 | 2.0 | 1.9 | 2.8 | 2.2 | 2.5 | 2.0 | 1.9 | 2.1 | 2.2 | 2.2 | 2.1 | 2.1 | 2.0 | 2.0 | 2.1 | 1.8 | 1.8 | 1.9 | 1.3 |
| Steroids | 2.4 | 2.8 | 2.2 | 2.9 | 2.4 | 3.1 | 2.3 | 2.7 | 2.5 | 3.0 | 2.8 | 2.7 | 2.2 | 2.2 | 2.1 | 2.0 | 1.6 | 1.7 | 1.7 | 1.6 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.9 | 5.7 | 4.7 | 4.6 | 4.4 | 4.3 | 4.4 | 4.4 | 4.9 | 4.4 | -0.5 |
| Tranquilizers | 13.0 | 14.6 | 13.5 | 16.3 | 16.3 | 16.1 | 15.5 | 14.9 | 16.4 | 17.4 | 16.5 | 16.0 | 15.0 | 14.3 | 13.6 | 12.1 | 12.0 | 4.8 | 4.3 | -0.5 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 8.6 | 6.0 | 5.1 | 4.7 | 4.3 | 3.9 | 4.3 | 4.1 | 5.2 | 5.1 | 6.1 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 13.7 | 12.6 | -1.1 |
| Prescription drugs | -- | -- | -- | -- | -- | -- | 6.8 | 6.6 | 7.5 | 7.5 | 6.5 | 6.1 | 6.5 | 6.5 | 10.6 | 10.0 | 9.6 | 12.2 | 11.2 | -1.0 |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 14.4 | 13.9 | 13.1 | 9.8 | 8.8 | -1.0 |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Monitoring the Future Study, the University of Michigan, 2011

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | National ^{###} Change* 2010 |
|-----------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---|
| Cigarettes | 30.4 | 33.0 | 34.4 | 36.7 | 37.2 | 33.9 | 31.5 | 28.9 | 27.0 | 23.8 | 22.2 | 22.1 | 20.8 | 20.6 | 19.3 | 18.7 | 18.1 | 18.6 | 17.2 | -1.4 13.6 |
| Smokeless tobacco | 13.2 | 13.6 | 13.8 | 14.6 | 12.3 | 9.5 | 8.2 | 7.4 | 6.9 | 7.1 | 6.0 | 6.4 | 7.0 | 7.8 | 7.3 | 7.5 | 7.6 | 8.1 | 7.9 | 7.5 |
| Cigars | -- | -- | -- | -- | -- | -- | 19.0 | 16.5 | 14.6 | 17.8 | 11.9 | 11.5 | 12.4 | 11.8 | 12.3 | 11.8 | 11.8 | 9.3 | 8.7 | -0.6 -- |
| Pipe | -- | -- | -- | -- | -- | -- | 4.8 | 4.3 | 3.6 | -- | 3.2 | 3.3 | 2.9 | 3.2 | 3.4 | 5.6 | 5.8 | 6.8 | 6.6 | -- |
| Alcohol | 43.8 | 44.9 | 42.7 | 43.7 | 43.9 | 41.3 | 41.6 | 39.7 | 39.7 | 38.1 | 36.9 | 34.1 | 33.0 | 33.0 | 31.1 | 28.4 | 27.3 | 30.3 | 28.4 | -1.9 28.9 |
| Binge drinking [#] | 27.3 | 26.8 | 26.1 | 27.3 | 27.8 | 29.0 | 28.8 | 27.1 | 24.2 | 24.9 | 21.8 | 20.2 | 19.3 | 19.9 | 21.7 | 19.5 | 19.2 | 18.7 | 18.1 | 16.3 |
| Marijuana | 14.3 | 18.9 | 21.5 | 24.9 | 23.7 | 21.8 | 21.1 | 19.0 | 21.4 | 19.2 | 18.2 | 17.2 | 16.0 | 14.6 | 14.4 | 13.5 | 14.6 | 16.8 | 16.4 | 16.7 |
| Cocaine | 1.4 | 1.6 | 2.2 | 2.6 | 2.6 | 2.9 | 2.3 | 1.7 | 2.4 | 2.4 | 2.3 | 2.4 | 2.2 | 2.1 | 1.9 | 1.7 | 1.4 | 1.6 | 1.6 | 0.9 |
| Crack | 0.1 | 1.2 | 1.6 | 1.8 | 1.9 | 2.1 | 1.5 | 1.1 | 1.5 | 1.1 | 1.4 | 1.4 | 1.3 | 1.2 | 1.0 | 1.1 | 0.8 | 1.1 | 1.0 | 0.5 |
| Inhalants | 5.0 | 5.6 | 5.8 | 5.1 | 5.3 | 4.9 | 3.4 | 2.9 | 3.3 | 2.9 | 2.9 | 3.3 | 3.0 | 3.0 | 2.5 | 2.6 | 2.2 | 2.5 | 2.1 | -0.4 2.0 |
| Amphetamines | 7.5 | 8.3 | 7.9 | 7.8 | 7.8 | 6.5 | 5.6 | 5.4 | 6.5 | 5.9 | 5.2 | 4.8 | 4.0 | 3.7 | 2.9 | 2.7 | 2.4 | 2.6 | 2.4 | 3.3 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.6 | 1.2 | 1.0 | 1.0 | 0.9 | 1.0 | 0.9 | 0.7 |
| Heroin | 0.1 | 0.9 | 1.0 | 1.3 | 1.0 | 1.3 | 0.9 | 0.7 | 1.0 | 0.8 | 0.9 | 0.9 | 0.9 | 0.8 | 0.8 | 1.0 | 0.8 | 0.9 | 0.9 | 0.4 |
| Steroids | 1.0 | 1.3 | 1.1 | 1.4 | 0.9 | 1.4 | 0.9 | 0.9 | 1.0 | 1.1 | 1.0 | 1.1 | 0.9 | 1.1 | 0.9 | 0.9 | 0.7 | 0.9 | 0.8 | 0.5 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.4 | 1.8 | 1.5 | 1.5 | 1.5 | 1.4 | 1.6 | 1.5 | 2.0 | 1.7 | -0.3 1.9 |
| Tranquilizers | 5.7 | 6.9 | 6.2 | 7.0 | 7.4 | 7.5 | 7.1 | 7.0 | 8.0 | 7.6 | 7.1 | 7.0 | 6.4 | 5.9 | 5.0 | 4.5 | 4.3 | 1.9 | 2.0 | 2.2 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.2 | 1.7 | 1.5 | 1.4 | 1.5 | 1.2 | 1.5 | 1.1 | 2.0 | 1.9 | 1.6 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.2 | 5.8 | -- |
| Prescription drugs | -- | -- | -- | -- | -- | -- | 2.5 | 2.1 | 2.4 | 2.6 | 2.2 | 2.1 | 2.0 | 2.1 | 2.2 | 3.7 | 3.5 | 5.9 | 5.2 | -0.7 -- |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 5.9 | 5.9 | 5.2 | 4.1 | 3.6 | -0.5 -- |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Monitoring the Future Study, the University of Michigan, 2011

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 64.7 | 64.5 | 66.5 | 67.7 | 66.9 | 66.9 | 65.3 | 62.3 | 58.7 | 54.8 | 51.2 | 50.1 | 46.9 | 46.8 | 44.6 | 43.2 | 42.0 | 39.8 | 38.1 | -1.7 |
| Smokeless tobacco | 31.9 | 31.0 | 32.2 | 31.9 | 29.1 | 25.6 | 23.0 | 22.9 | 17.4 | 17.6 | 16.3 | 18.8 | 15.8 | 17.2 | 16.9 | 17.3 | 17.9 | 20.7 | 20.0 | |
| Cigars | -- | -- | -- | -- | -- | 43.9 | 42.1 | 39.3 | 40.7 | 34.2 | 31.5 | 31.8 | 30.9 | 30.7 | 31.0 | 30.6 | 29.9 | 28.1 | 27.0 | -1.1 |
| Pipe | -- | -- | -- | -- | -- | 14.5 | 13.1 | 12.4 | -- | 9.4 | 8.8 | 9.3 | 8.1 | 9.2 | 16.2 | 16.7 | 16.9 | 19.9 | 20.1 | |
| Alcohol | 82.2 | 79.7 | 78.5 | 78.1 | 77.5 | 74.9 | 73.3 | 71.9 | 74.3 | 72.9 | 70.2 | 68.7 | 66.6 | 66.7 | 64.3 | 62.4 | 60.6 | 61.9 | 60.6 | -1.3 |
| Marijuana | 31.1 | 33.2 | 36.6 | 44.1 | 42.3 | 43.7 | 42.1 | 39.8 | 41.0 | 41.1 | 38.4 | 35.9 | 34.3 | 34.4 | 32.4 | 32.4 | 32.7 | 34.6 | 35.4 | |
| Cocaine | 4.2 | 4.9 | 5.2 | 7.5 | 7.8 | 8.1 | 7.6 | 7.5 | 7.1 | 7.8 | 8.0 | 7.8 | 7.1 | 7.8 | 6.9 | 6.3 | 5.4 | 5.2 | 5.1 | |
| Crack | 2.0 | 2.7 | 3.2 | 4.1 | 4.5 | 5.3 | 4.2 | 3.8 | 3.8 | 4.0 | 3.8 | 4.6 | 3.7 | 3.8 | 3.3 | 3.0 | 2.8 | 2.7 | 2.5 | |
| Inhalants | 14.0 | 15.7 | 16.0 | 17.6 | 15.4 | 16.2 | 12.3 | 13.0 | 11.3 | 10.5 | 9.9 | 11.1 | 9.8 | 10.1 | 9.7 | 9.3 | 8.6 | 9.1 | 7.9 | -1.2 |
| Amphetamines | 18.8 | 18.5 | 18.2 | 18.3 | 18.5 | 16.8 | 15.7 | 15.5 | 15.1 | 14.8 | 14.3 | 13.6 | 11.1 | 10.8 | 9.1 | 8.7 | 8.1 | 8.2 | 7.8 | |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 4.6 | 4.2 | 3.3 | 2.8 | 2.6 | 2.6 | 2.7 | |
| Heroin | 1.2 | 1.5 | 1.7 | 1.9 | 1.9 | 2.4 | 2.0 | 2.5 | 2.1 | 2.1 | 2.1 | 2.4 | 2.1 | 2.3 | 2.0 | 2.0 | 2.2 | 2.3 | 2.2 | |
| Steroids | 2.0 | 2.6 | 2.5 | 2.3 | 2.4 | 2.7 | 2.2 | 2.7 | 2.3 | 2.5 | 2.5 | 2.3 | 2.1 | 1.9 | 1.8 | 1.8 | 1.8 | 1.9 | 2.0 | |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 8.1 | 6.5 | 5.7 | 4.9 | 5.4 | 4.8 | 5.0 | 5.3 | 6.2 | 5.7 | -0.5 |
| Tranquilizers | 12.9 | 13.5 | 13.9 | 16.6 | 16.3 | 17.5 | 16.0 | 16.7 | 17.0 | 18.6 | 16.7 | 15.9 | 14.6 | 14.5 | 13.4 | 12.6 | 11.8 | 5.7 | 5.2 | -0.5 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 10.7 | 7.9 | 6.4 | 4.9 | 4.9 | 4.4 | 4.8 | 4.8 | 6.3 | 6.5 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 15.8 | 14.9 | -0.9 |
| Prescription drugs | -- | -- | -- | -- | -- | 7.5 | 7.3 | 8.0 | 7.3 | 7.6 | 7.0 | 5.9 | 6.6 | 6.7 | 10.7 | 11.6 | 11.1 | 14.1 | 13.3 | -0.8 |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 13.5 | 13.3 | 12.7 | 10.0 | 9.5 | |

NOTES: -- Data not available
* Statistically significant changes between 2010 and 2011 prevalence rates (p<.05).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---------|
| Cigarettes | 33.7 | 34.4 | 39.3 | 40.1 | 39.2 | 39.8 | 37.4 | 34.1 | 30.2 | 27.1 | 25.2 | 25.1 | 23.5 | 23.8 | 22.5 | 21.3 | 21.4 | 21.9 | 20.7 | -1.2 |
| Smokeless tobacco | 14.4 | 14.6 | 15.4 | 14.2 | 12.0 | 10.2 | 8.9 | 9.7 | 7.0 | 7.5 | 7.2 | 8.1 | 7.3 | 8.0 | 8.1 | 8.2 | 8.8 | 9.2 | 8.8 | |
| Cigars | -- | -- | -- | -- | -- | 21.0 | 18.6 | 16.8 | 19.8 | 13.4 | 13.1 | 13.9 | 13.0 | 12.8 | 13.8 | 13.6 | 13.9 | 11.8 | 11.1 | |
| Pipe | -- | -- | -- | -- | -- | 4.5 | 4.3 | 3.8 | -- | 3.2 | 3.2 | 3.2 | 3.0 | 3.5 | 7.4 | 7.0 | 7.5 | 8.4 | 8.3 | |
| Alcohol | 49.4 | 47.0 | 47.7 | 48.2 | 45.8 | 47.8 | 45.6 | 43.2 | 42.1 | 41.4 | 38.1 | 36.5 | 34.5 | 35.8 | 33.8 | 31.7 | 29.4 | 32.3 | 31.7 | |
| Binge drinking# | 30.6 | 28.9 | 29.6 | 30.1 | 29.8 | 32.3 | 31.3 | 30.0 | 25.7 | 26.5 | 24.0 | 23.3 | 21.2 | 21.8 | 23.2 | 22.5 | 21.4 | 20.9 | 20.8 | |
| Marijuana | 16.7 | 19.6 | 21.8 | 25.7 | 22.3 | 24.6 | 22.0 | 20.5 | 21.7 | 21.2 | 18.4 | 17.2 | 15.3 | 15.7 | 14.6 | 14.6 | 15.3 | 17.9 | 18.6 | |
| Cocaine | 1.2 | 1.6 | 2.0 | 3.0 | 2.9 | 3.3 | 2.4 | 2.2 | 2.6 | 2.7 | 2.5 | 3.1 | 2.1 | 2.7 | 2.1 | 1.8 | 1.4 | 1.9 | 1.8 | |
| Crack | 0.1 | 1.0 | 1.3 | 1.6 | 1.6 | 1.9 | 1.0 | 1.0 | 1.2 | 1.1 | 1.1 | 1.8 | 1.1 | 1.3 | 1.1 | 1.0 | 0.9 | 1.1 | 1.2 | |
| Inhalants | 3.8 | 4.2 | 4.3 | 4.3 | 3.6 | 3.9 | 2.7 | 2.8 | 2.3 | 2.1 | 2.2 | 2.7 | 2.1 | 2.3 | 2.2 | 1.9 | 1.9 | 2.1 | 1.8 | -0.3 |
| Amphetamines | 7.2 | 7.0 | 7.0 | 6.4 | 7.0 | 6.0 | 5.8 | 6.6 | 6.4 | 5.6 | 5.3 | 5.4 | 3.5 | 3.9 | 2.8 | 2.8 | 2.5 | 2.8 | 2.6 | |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.5 | 1.5 | 1.1 | 0.9 | 0.9 | 1.2 | 1.3 | |
| Heroin | 0.1 | 0.6 | 0.8 | 0.8 | 0.8 | 1.0 | 0.7 | 0.9 | 0.7 | 0.6 | 0.8 | 1.1 | 0.8 | 1.0 | 0.9 | 0.7 | 0.8 | 1.1 | 1.1 | |
| Steroids | 0.1 | 1.1 | 1.0 | 0.8 | 0.8 | 1.0 | 0.6 | 0.9 | 0.7 | 0.8 | 1.0 | 1.0 | 0.9 | 0.8 | 0.7 | 0.7 | 0.8 | 0.9 | 1.0 | |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.3 | 1.7 | 1.8 | 1.3 | 1.8 | 1.3 | 1.5 | 1.5 | 2.1 | 1.8 | -0.3 |
| Tranquilizers | 4.6 | 5.3 | 5.7 | 6.9 | 7.0 | 7.1 | 6.9 | 7.3 | 7.7 | 7.6 | 6.7 | 6.5 | 5.1 | 5.6 | 4.3 | 4.2 | 3.8 | 2.1 | 2.0 | |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.1 | 1.7 | 1.7 | 1.3 | 1.6 | 1.3 | 1.4 | 1.3 | 2.1 | 2.2 | |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.7 | 5.9 | -0.8 |
| Prescription drugs | -- | -- | -- | -- | -- | 2.4 | 1.9 | 2.2 | 2.0 | 2.0 | 1.8 | 1.6 | 1.8 | 2.0 | 3.2 | 3.4 | 3.4 | 6.2 | 5.6 | -0.6 |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 4.7 | 4.6 | 4.4 | 3.6 | 3.3 | |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | Change* National ^{###} 2010 | 2010 |
|--------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|--|------|
| Cigarettes | 66.4 | 65.6 | 67.9 | 66.4 | 68.6 | 67.2 | 68.1 | 65.8 | 62.9 | 60.5 | 55.9 | 53.9 | 51.9 | 50.9 | 48.4 | 48.2 | 46.9 | 43.7 | 42.1 | -1.6 | 42.2 |
| Smokeless tobacco | 34.0 | 33.9 | 35.8 | 31.3 | 31.4 | 29.4 | 26.9 | 26.3 | 22.9 | 21.2 | 18.7 | 19.1 | 18.8 | 20.4 | 19.4 | 20.1 | 20.7 | 22.2 | 23.6 | 1.4 | 17.6 |
| Cigars | -- | -- | -- | -- | -- | 50.1 | 48.9 | 45.3 | 46.7 | 41.2 | 39.4 | 37.3 | 38.9 | 37.7 | 37.9 | 37.8 | 37.5 | 36.0 | 35.9 | -- | -- |
| Pipe | -- | -- | -- | -- | -- | 15.9 | 15.6 | 14.0 | -- | 10.9 | 9.7 | 10.0 | 10.5 | 10.7 | 20.0 | 22.6 | 22.9 | 27.5 | 27.1 | -- | -- |
| Alcohol | 85.0 | 82.4 | 81.4 | 80.6 | 80.8 | 76.1 | 78.5 | 75.9 | 78.6 | 77.3 | 74.8 | 72.4 | 72.3 | 70.2 | 69.2 | 68.5 | 66.5 | 67.8 | 65.4 | -2.4 | 71.0 |
| Marijuana | 35.3 | 38.1 | 40.0 | 43.4 | 45.6 | 46.5 | 45.7 | 44.4 | 45.5 | 44.8 | 42.3 | 40.5 | 40.1 | 37.1 | 36.5 | 36.5 | 36.8 | 38.6 | 38.6 | -- | 43.8 |
| Cocaine | 5.0 | 6.1 | 7.1 | 7.1 | 8.9 | 9.3 | 8.4 | 8.7 | 9.0 | 8.6 | 8.7 | 8.8 | 8.8 | 8.6 | 7.8 | 7.4 | 6.6 | 6.0 | 5.9 | -- | 5.5 |
| Crack | 2.0 | 2.8 | 4.3 | 4.1 | 5.6 | 5.6 | 4.5 | 4.2 | 4.1 | 4.1 | 4.0 | 4.4 | 3.9 | 4.0 | 3.2 | 3.1 | 2.9 | 2.7 | 2.6 | -- | 2.4 |
| Inhalants | 14.0 | 15.1 | 15.5 | 13.7 | 15.5 | 15.3 | 12.8 | 12.3 | 11.6 | 10.3 | 9.6 | 9.5 | 9.6 | 10.1 | 8.5 | 9.0 | 8.0 | 8.2 | 7.8 | -- | 9.0 |
| Amphetamines | 19.8 | 18.8 | 18.0 | 16.5 | 19.2 | 16.6 | 15.6 | 15.4 | 15.5 | 14.9 | 14.4 | 13.3 | 12.5 | 11.3 | 9.0 | 9.3 | 8.5 | 8.3 | 8.3 | -- | 11.1 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 5.5 | 5.0 | 3.4 | 2.7 | 2.7 | 2.5 | 2.8 | -- | 2.3 |
| Heroin | 1.3 | 1.8 | 2.0 | 2.1 | 2.3 | 2.7 | 2.1 | 2.3 | 2.3 | 1.9 | 2.1 | 2.3 | 2.0 | 2.3 | 1.9 | 2.0 | 2.4 | 2.3 | 2.4 | -- | 1.6 |
| Steroids | 2.4 | 2.8 | 2.6 | 2.1 | 2.5 | 2.6 | 2.5 | 2.5 | 2.3 | 2.4 | 2.3 | 2.3 | 2.1 | 2.3 | 1.9 | 1.9 | 2.1 | 2.0 | 2.1 | -- | 2.0 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 9.5 | 7.4 | 6.4 | 5.9 | 5.6 | 5.1 | 5.5 | 5.9 | 6.5 | 6.7 | -- | 7.3 |
| Tranquilizers | 12.6 | 13.2 | 13.2 | 14.2 | 16.2 | 16.4 | 15.8 | 15.7 | 16.5 | 17.3 | 16.6 | 15.6 | 15.5 | 14.5 | 12.9 | 12.4 | 12.0 | 5.2 | 5.4 | -- | 8.5 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 11.9 | 9.3 | 6.6 | 5.5 | 5.3 | 4.5 | 4.8 | 5.0 | 6.6 | 7.2 | 0.6 | 8.6 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 16.4 | 16.0 | -- | -- |
| Prescription drugs | -- | -- | -- | -- | -- | 5.9 | 7.4 | 7.4 | 7.8 | 7.2 | 6.9 | 6.3 | 7.1 | 7.0 | 11.3 | 11.8 | 11.6 | 14.6 | 14.6 | -- | -- |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 13.5 | 12.9 | 12.5 | 10.0 | 9.7 | -- | -- |

NOTES: -- Data not available
 * Statistically significant changes between 2010 and 2011 prevalence rates (p<.05).
 ### Monitoring the Future Study, the University of Michigan, 2011

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 1993-2011

| DRUG | 1993 | 1994 | 1995 | 1996 | 1997 | 1998 | 1999 | 2000 | 2001 | 2002 | 2003 | 2004 | 2005 | 2006 | 2007 | 2008 | 2009 | 2010 | 2011 | National ^{###} Change* 2010 |
|-----------------------------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|------|---|
| Cigarettes | 35.6 | 37.3 | 40.6 | 39.8 | 41.6 | 41.6 | 40.5 | 38.9 | 35.1 | 30.5 | 28.8 | 27.4 | 26.5 | 26.9 | 24.3 | 24.8 | 24.4 | 24.9 | 24.6 | 19.2 |
| Smokeless tobacco | 15.4 | 15.5 | 16.5 | 14.4 | 13.5 | 12.1 | 11.2 | 11.3 | 9.4 | 8.3 | 7.9 | 8.7 | 8.6 | 9.8 | 9.1 | 9.8 | 10.5 | 10.5 | 10.7 | 8.5 |
| Cigars | -- | -- | -- | -- | -- | 25.8 | 23.0 | 20.2 | 23.9 | 16.8 | 17.0 | 16.8 | 17.6 | 16.6 | 17.5 | 17.7 | 17.9 | 16.8 | 16.7 | -- |
| Pipe | -- | -- | -- | -- | -- | 4.8 | 5.2 | 4.4 | -- | 3.3 | 3.3 | 3.4 | 3.7 | 4.6 | 9.3 | 10.1 | 10.3 | 13.0 | 12.5 | -- |
| Alcohol | 53.2 | 53.7 | 51.2 | 52.5 | 50.1 | 52.0 | 51.7 | 50.1 | 49.9 | 48.1 | 46.1 | 42.2 | 41.8 | 42.2 | 39.7 | 38.4 | 35.7 | 39.4 | 37.8 | -1.6 41.2 |
| Binge drinking [#] | 34.3 | 34.8 | 33.3 | 35.6 | 33.2 | 37.5 | 36.3 | 35.4 | 31.9 | 31.5 | 29.3 | 27.0 | 25.9 | 27.3 | 28.6 | 26.9 | 26.4 | 26.2 | 26.4 | 23.2 |
| Marijuana | 17.8 | 21.6 | 24.0 | 25.4 | 23.5 | 24.5 | 23.5 | 22.7 | 23.5 | 20.5 | 19.8 | 18.3 | 17.8 | 17.2 | 15.8 | 16.2 | 16.7 | 19.2 | 19.8 | 21.4 |
| Cocaine | 1.5 | 2.0 | 2.5 | 3.0 | 3.3 | 3.5 | 2.4 | 2.7 | 2.8 | 2.9 | 2.5 | 2.9 | 2.9 | 3.1 | 2.4 | 2.0 | 1.6 | 1.9 | 2.0 | 1.3 |
| Crack | 0.1 | 0.9 | 1.7 | 1.5 | 1.8 | 2.0 | 1.1 | 1.1 | 1.1 | 1.1 | 1.1 | 1.4 | 1.2 | 1.4 | 1.1 | 1.0 | 0.9 | 1.0 | 1.2 | 0.7 |
| Inhalants | 2.9 | 3.8 | 3.8 | 3.1 | 3.3 | 3.2 | 2.6 | 2.4 | 2.0 | 1.9 | 1.7 | 1.8 | 1.9 | 2.2 | 1.5 | 1.6 | 1.4 | 1.8 | 1.7 | 1.4 |
| Amphetamines | 6.5 | 7.2 | 6.5 | 6.1 | 6.7 | 5.8 | 5.1 | 5.9 | 5.7 | 5.3 | 4.9 | 4.5 | 3.8 | 4.1 | 2.5 | 2.6 | 2.3 | 2.8 | 2.7 | 3.3 |
| Methamphetamines | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.7 | 1.5 | 1.0 | 0.9 | 0.9 | 1.0 | 1.3 | 0.3 0.5 |
| Heroin | 0.1 | 0.7 | 0.9 | 1.1 | 1.0 | 1.1 | 0.7 | 0.9 | 0.8 | 0.6 | 0.6 | 0.8 | 0.7 | 1.0 | 0.8 | 0.8 | 0.8 | 1.1 | 1.2 | 0.4 |
| Steroids | 0.1 | 0.9 | 0.8 | 0.8 | 1.0 | 1.1 | 0.9 | 0.6 | 0.7 | 0.6 | 0.7 | 0.8 | 0.8 | 1.0 | 0.7 | 0.9 | 0.8 | 0.9 | 1.1 | 0.2 1.1 |
| Ecstasy | -- | -- | -- | -- | -- | -- | -- | -- | -- | 2.1 | 1.6 | 1.4 | 1.3 | 1.7 | 1.4 | 1.6 | 1.5 | 1.9 | 2.1 | 1.4 |
| Tranquilizers | 4.5 | 4.6 | 4.9 | 6.0 | 6.2 | 6.6 | 5.9 | 6.5 | 6.7 | 6.0 | 5.9 | 5.4 | 5.2 | 5.3 | 4.0 | 4.0 | 3.7 | 1.8 | 2.0 | 2.5 |
| Hallucinogens | -- | -- | -- | -- | -- | -- | -- | -- | -- | 1.5 | 1.5 | 1.4 | 1.2 | 1.3 | 1.1 | 1.3 | 1.2 | 2.0 | 2.4 | 0.4 1.9 |
| Prescription painkillers | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 6.3 | 6.6 | -- |
| Prescription drugs | -- | -- | -- | -- | -- | 1.7 | 1.5 | 1.9 | 2.0 | 1.4 | 1.5 | 1.4 | 1.7 | 1.8 | 2.9 | 3.3 | 3.1 | 5.9 | 5.9 | -- |
| Over the counter drugs | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | -- | 4.3 | 4.3 | 4.0 | 3.2 | 3.2 | -- |

NOTES: -- Data not available

* Statistically significant changes between 2010 and 2011 prevalence rates ($p < .05$).

Binge drinking during the past two weeks.

Monitoring the Future Study, the University of Michigan, 2011

Mean Age of First Time Use of Alcohol, Tobacco, and Other Drugs, 2011

| DRUG | 6th | 7th | 8th | 9th | 10th | 11th | 12th | Total |
|--------------------------|------|------|------|------|------|------|------|-------|
| Cigarettes | 10.7 | 11.3 | 11.9 | 12.6 | 13.2 | 13.8 | 14.6 | 13.0 |
| Smokeless Tobacco | 10.9 | 11.5 | 12.2 | 13.0 | 13.8 | 14.4 | 15.0 | 13.6 |
| Cigars | 10.9 | 11.5 | 12.3 | 13.2 | 14.0 | 14.6 | 15.4 | 14.1 |
| Pipe | 11.0 | 11.6 | 12.4 | 13.3 | 14.1 | 14.8 | 15.7 | 14.3 |
| Alcohol | 10.7 | 11.3 | 12.0 | 12.9 | 13.6 | 14.2 | 14.8 | 13.2 |
| Marijuana | 11.2 | 11.8 | 12.5 | 13.3 | 14.0 | 14.6 | 15.1 | 13.9 |
| Cocaine | 10.9 | 11.7 | 12.5 | 13.2 | 14.1 | 14.7 | 15.2 | 14.0 |
| Crack | 10.8 | 11.6 | 12.3 | 13.0 | 13.5 | 13.8 | 14.2 | 13.1 |
| Inhalants | 10.8 | 11.5 | 12.3 | 13.0 | 13.7 | 13.9 | 14.5 | 13.2 |
| Amphetamines | 10.8 | 11.8 | 12.5 | 13.4 | 14.1 | 14.5 | 15.0 | 14.0 |
| Methamphetamines | 10.8 | 11.7 | 12.3 | 13.0 | 13.9 | 14.3 | 14.9 | 13.5 |
| Heroin | 10.9 | 11.6 | 12.4 | 13.0 | 13.7 | 14.0 | 14.6 | 13.4 |
| Steroids | 10.9 | 11.7 | 12.0 | 13.0 | 13.6 | 13.9 | 14.5 | 13.2 |
| Ecstasy | 11.0 | 11.8 | 12.7 | 13.4 | 14.3 | 14.9 | 15.5 | 14.2 |
| Tranquilizers | 10.9 | 11.8 | 12.3 | 13.3 | 13.8 | 14.3 | 14.8 | 13.8 |
| Hallucinogens | 11.2 | 11.7 | 12.7 | 13.4 | 14.2 | 14.9 | 15.5 | 14.2 |
| Prescription painkillers | 10.9 | 11.8 | 12.6 | 13.5 | 14.2 | 14.8 | 15.3 | 14.1 |
| Prescription drugs | 10.8 | 11.8 | 12.6 | 13.5 | 14.3 | 14.8 | 15.3 | 14.1 |
| Over the counter drugs | 11.2 | 11.8 | 12.6 | 13.3 | 14.0 | 14.6 | 15.0 | 13.7 |



III. PREVALENCE OF USE BY RACE/ETHNICITY

Prevalence Rates by Race/Ethnicity: Lifetime and Monthly Use

| | |
|------------------|----|
| • Grade 6 | 78 |
| • Grade 7 | 80 |
| • Grade 8 | 82 |
| • Grade 9 | 84 |
| • Grade 10 | 86 |
| • Grade 11 | 88 |
| • Grade 12 | 90 |

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

6th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|-------|---------|
| Cigarettes | 6.5 * | 10.2 * | 9.6 * | 9.4 * | 7.3 |
| Smokeless tobacco | 2.3 | 1.7 | 3.2 * | 3.4 * | 2.5 |
| Cigars | 1.4 * | 2.8 * | 3.1 * | 3.0 * | 1.9 |
| Pipe | 0.9 * | 1.9 | 3.0 * | 2.2 * | 1.3 |
| Alcohol | 16.4 * | 21.2 * | 21.8 * | 18.8 | 17.4 |
| Marijuana | 2.1 * | 8.7 * | 6.0 * | 4.3 * | 3.1 |
| Cocaine | 0.2 * | 1.2 * | 2.1 * | 1.1 * | 0.6 |
| Crack | 0.3 * | 1.1 * | 1.7 * | 0.9 * | 0.6 |
| Inhalants | 1.5 * | 2.4 | 2.8 * | 2.4 * | 1.8 |
| Amphetamines | 0.2 * | 1.0 * | 1.4 * | 0.6 | 0.4 |
| Methamphetamines | 0.3 * | 1.1 * | 1.9 * | 0.8 | 0.5 |
| Heroin | 0.2 * | 1.0 * | 1.5 * | 0.8 * | 0.4 |
| Steroids | 0.3 * | 1.2 * | 1.5 * | 0.9 * | 0.6 |
| Ecstasy | 0.2 * | 1.0 * | 1.4 * | 0.7 * | 0.4 |
| Tranquilizers | 0.2 * | 1.2 * | 1.5 * | 0.7 | 0.5 |
| Hallucinogens | 0.2 * | 1.1 * | 1.1 * | 0.9 * | 0.4 |
| Prescription painkillers | 0.8 * | 1.5 | 2.5 * | 1.3 | 1.1 |
| Prescription drugs | 0.8 * | 1.7 * | 2.4 * | 1.2 | 1.0 |
| Over the counter drugs | 0.8 * | 1.7 * | 2.5 * | 1.6 * | 1.1 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

6th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|-------|-------|----------|-------|---------|
| Cigarettes | 2.3 * | 3.6 * | 3.5 * | 3.6 * | 2.6 |
| Smokeless tobacco | 0.8 | 0.7 | 1.5 * | 1.5 * | 1.0 |
| Cigars | 0.5 * | 1.2 * | 1.2 * | 1.2 * | 0.7 |
| Pipe | 0.5 | 0.9 | 1.3 * | 0.9 | 0.6 |
| Alcohol | 4.5 * | 8.6 * | 8.3 * | 6.7 * | 5.3 |
| Binge drinking [#] | 4.8 * | 9.4 * | 10.4 * | 7.2 * | 5.8 |
| Marijuana | 1.1 * | 5.0 * | 3.4 * | 2.2 * | 1.6 |
| Cocaine | 0.1 * | 0.4 | 1.1 * | 0.6 * | 0.3 |
| Crack | 0.1 * | 0.4 | 1.1 * | 0.4 | 0.3 |
| Inhalants | 0.7 | 1.1 | 1.6 * | 1.1 | 0.9 |
| Amphetamines | 0.1 * | 0.6 * | 1.1 * | 0.3 | 0.2 |
| Methamphetamines | 0.1 * | 0.7 * | 1.1 * | 0.4 | 0.3 |
| Heroin | 0.1 * | 0.5 * | 0.9 * | 0.4 | 0.2 |
| Steroids | 0.1 * | 0.5 | 0.8 * | 0.3 | 0.2 |
| Ecstasy | 0.1 * | 0.5 | 0.9 * | 0.5 * | 0.2 |
| Tranquilizers | 0.1 * | 0.5 | 0.9 * | 0.4 | 0.3 |
| Hallucinogens | 0.1 * | 0.4 | 0.9 * | 0.5 * | 0.3 |
| Prescription painkillers | 0.5 | 0.5 | 1.3 * | 1.0 * | 0.6 |
| Prescription drugs | 0.5 | 1.1 | 1.1 * | 0.8 | 0.6 |
| Over the counter drugs | 0.5 * | 1.0 | 1.7 * | 1.0 | 0.7 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

7th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 12.0 * | 17.6 * | 17.6 * | 15.6 * | 13.2 |
| Smokeless tobacco | 5.0 | 2.9 * | 5.0 | 6.5 * | 5.1 |
| Cigars | 3.7 * | 4.5 | 6.3 * | 6.0 * | 4.3 |
| Pipe | 2.8 * | 2.7 | 5.3 * | 4.2 * | 3.2 |
| Alcohol | 24.7 * | 33.0 * | 32.7 * | 27.0 | 26.1 |
| Marijuana | 5.8 * | 14.7 * | 12.5 * | 9.9 * | 7.3 |
| Cocaine | 0.8 * | 1.4 | 3.1 * | 1.2 | 1.1 |
| Crack | 0.8 * | 1.4 | 3.2 * | 1.4 | 1.1 |
| Inhalants | 2.9 | 3.5 | 4.5 * | 3.6 | 3.2 |
| Amphetamines | 0.9 | 1.3 | 1.7 * | 1.3 | 1.0 |
| Methamphetamines | 0.7 * | 1.3 | 2.5 * | 1.5 * | 1.0 |
| Heroin | 0.6 * | 1.2 | 2.4 * | 0.8 | 0.8 |
| Steroids | 0.7 * | 1.2 | 2.5 * | 1.4 | 1.0 |
| Ecstasy | 0.9 * | 1.2 | 2.4 * | 1.2 | 1.1 |
| Tranquilizers | 0.8 | 0.9 | 2.1 * | 1.2 | 0.9 |
| Hallucinogens | 0.8 | 1.2 | 2.1 * | 0.9 | 0.9 |
| Prescription painkillers | 2.3 * | 2.9 | 4.7 * | 3.0 | 2.7 |
| Prescription drugs | 2.0 * | 2.5 | 4.4 * | 2.6 | 2.4 |
| Over the counter drugs | 2.4 * | 2.9 | 5.3 * | 3.1 | 2.8 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

7th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|-------|--------|----------|-------|---------|
| Cigarettes | 5.6 | 7.2 | 8.1 * | 6.5 | 6.0 |
| Smokeless tobacco | 2.2 | 1.9 | 2.4 | 2.3 | 2.2 |
| Cigars | 1.3 | 1.8 | 2.7 * | 2.2 * | 1.6 |
| Pipe | 1.2 * | 1.7 | 2.6 * | 2.0 * | 1.5 |
| Alcohol | 9.4 * | 13.5 * | 14.1 * | 11.3 | 10.2 |
| Binge drinking [#] | 6.7 * | 13.2 * | 13.1 * | 9.5 * | 7.9 |
| Marijuana | 3.3 * | 9.2 * | 7.2 * | 5.1 * | 4.1 |
| Cocaine | 0.4 * | 1.0 | 1.7 * | 0.8 | 0.6 |
| Crack | 0.5 * | 0.8 | 2.1 * | 0.7 | 0.7 |
| Inhalants | 1.3 | 2.2 | 1.9 | 1.7 | 1.4 |
| Amphetamines | 0.5 | 0.8 | 1.1 * | 0.4 | 0.5 |
| Methamphetamines | 0.3 * | 0.9 | 1.9 * | 0.5 | 0.5 |
| Heroin | 0.3 * | 0.4 | 1.5 * | 0.5 | 0.4 |
| Steroids | 0.4 * | 0.9 | 1.3 * | 0.6 | 0.5 |
| Ecstasy | 0.4 | 0.3 | 1.0 * | 0.7 | 0.5 |
| Tranquilizers | 0.4 | 0.3 | 1.2 * | 0.6 | 0.5 |
| Hallucinogens | 0.4 | 0.4 | 1.2 * | 0.5 | 0.5 |
| Prescription painkillers | 1.4 | 1.0 | 2.3 * | 1.5 | 1.5 |
| Prescription drugs | 1.2 | 1.3 | 2.6 * | 1.7 | 1.4 |
| Over the counter drugs | 1.4 | 1.8 | 3.2 * | 1.8 | 1.6 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

8th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 19.2 * | 21.8 | 25.2 * | 25.1 * | 20.6 |
| Smokeless tobacco | 8.9 | 4.2 * | 8.0 | 9.2 | 8.6 |
| Cigars | 7.5 * | 8.3 | 10.2 * | 10.3 * | 8.1 |
| Pipe | 5.4 * | 4.2 * | 8.8 * | 7.8 * | 6.0 |
| Alcohol | 36.0 * | 44.8 * | 45.5 * | 39.4 * | 37.7 |
| Marijuana | 12.1 * | 24.8 * | 21.1 * | 20.1 * | 14.6 |
| Cocaine | 1.5 * | 2.8 | 5.1 * | 2.9 * | 2.0 |
| Crack | 1.2 * | 2.7 * | 4.3 * | 2.1 | 1.7 |
| Inhalants | 5.0 * | 5.5 | 7.1 * | 6.6 * | 5.4 |
| Amphetamines | 2.1 | 2.1 | 3.5 * | 2.8 | 2.3 |
| Methamphetamines | 1.2 * | 1.7 | 3.3 * | 2.2 * | 1.5 |
| Heroin | 0.8 * | 2.0 * | 3.0 * | 1.6 * | 1.2 |
| Steroids | 0.9 * | 1.9 | 2.7 * | 2.3 * | 1.3 |
| Ecstasy | 1.7 * | 2.8 | 4.0 * | 3.3 * | 2.2 |
| Tranquilizers | 1.7 * | 1.5 | 3.9 * | 2.2 | 1.9 |
| Hallucinogens | 1.9 * | 2.8 | 4.0 * | 3.3 * | 2.3 |
| Prescription painkillers | 5.0 * | 5.2 | 7.9 * | 7.1 * | 5.6 |
| Prescription drugs | 4.3 * | 4.5 | 6.8 * | 6.1 * | 4.8 |
| Over the counter drugs | 4.5 * | 4.7 | 7.3 * | 5.9 * | 5.0 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

8th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 9.4 * | 8.8 | 12.2 * | 12.5 * | 10.0 |
| Smokeless tobacco | 3.9 | 2.1 * | 3.6 | 4.5 | 3.9 |
| Cigars | 2.8 * | 4.2 * | 4.5 * | 4.4 * | 3.2 |
| Pipe | 2.6 * | 2.6 | 4.4 * | 3.9 * | 3.0 |
| Alcohol | 15.7 * | 19.5 * | 21.6 * | 18.2 * | 16.7 |
| Binge drinking [#] | 9.6 * | 15.3 * | 17.1 * | 14.1 * | 11.1 |
| Marijuana | 6.8 * | 15.2 * | 12.1 * | 11.4 * | 8.3 |
| Cocaine | 0.7 * | 1.5 | 3.0 * | 1.5 * | 1.0 |
| Crack | 0.6 * | 1.5 * | 2.3 * | 1.2 * | 0.9 |
| Inhalants | 2.0 * | 2.7 | 3.2 * | 3.0 * | 2.3 |
| Amphetamines | 1.1 | 1.1 | 2.1 * | 1.3 | 1.2 |
| Methamphetamines | 0.6 * | 1.0 | 2.2 * | 1.4 * | 0.9 |
| Heroin | 0.5 * | 1.0 | 2.0 * | 0.9 | 0.7 |
| Steroids | 0.5 * | 1.1 | 1.5 * | 1.3 * | 0.7 |
| Ecstasy | 0.9 * | 1.1 | 2.3 * | 1.5 * | 1.1 |
| Tranquilizers | 0.8 * | 0.7 | 2.2 * | 1.5 * | 1.0 |
| Hallucinogens | 0.9 * | 1.3 | 2.3 * | 1.8 * | 1.2 |
| Prescription painkillers | 2.7 | 3.1 | 4.3 * | 3.9 * | 3.0 |
| Prescription drugs | 2.5 | 2.7 | 3.8 * | 3.4 * | 2.7 |
| Over the counter drugs | 2.4 * | 2.2 | 3.9 * | 3.7 * | 2.7 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

9th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 26.2 * | 21.6 * | 33.4 * | 30.8 * | 27.2 |
| Smokeless tobacco | 13.8 | 4.9 * | 10.1 * | 14.2 | 13.1 |
| Cigars | 14.5 | 10.8 * | 16.5 | 17.5 * | 14.9 |
| Pipe | 9.8 | 6.2 * | 13.8 * | 13.0 * | 10.4 |
| Alcohol | 46.5 * | 46.6 | 55.9 * | 50.1 * | 47.7 |
| Marijuana | 20.4 * | 31.6 * | 30.8 * | 29.9 * | 23.0 |
| Cocaine | 2.2 * | 2.2 | 6.7 * | 4.8 * | 2.9 |
| Crack | 1.4 * | 1.6 | 3.9 * | 3.8 * | 1.9 |
| Inhalants | 6.4 | 5.1 * | 9.1 * | 8.7 * | 6.9 |
| Amphetamines | 4.0 | 1.9 * | 5.3 * | 5.7 * | 4.3 |
| Methamphetamines | 1.4 * | 1.2 | 3.9 * | 3.3 * | 1.9 |
| Heroin | 1.2 * | 1.9 | 3.2 * | 3.0 * | 1.6 |
| Steroids | 1.2 * | 1.6 | 2.8 * | 2.7 * | 1.5 |
| Ecstasy | 2.8 * | 3.1 | 6.2 * | 5.3 * | 3.4 |
| Tranquilizers | 2.9 * | 1.9 * | 5.2 * | 4.7 * | 3.3 |
| Hallucinogens | 3.1 * | 2.5 | 5.2 * | 5.4 * | 3.6 |
| Prescription painkillers | 9.0 | 5.3 * | 12.2 * | 11.4 * | 9.4 |
| Prescription drugs | 7.9 | 4.9 * | 10.6 * | 9.7 * | 8.2 |
| Over the counter drugs | 6.8 | 4.3 * | 9.3 * | 9.0 * | 7.2 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

9th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 14.0 | 8.2 * | 17.2 * | 16.4 * | 14.4 |
| Smokeless tobacco | 6.4 | 1.7 * | 5.1 | 7.3 * | 6.2 |
| Cigars | 5.6 | 5.5 | 7.2 * | 7.6 * | 6.0 |
| Pipe | 4.2 * | 3.2 * | 7.0 * | 6.9 * | 4.8 |
| Alcohol | 22.7 * | 23.6 | 30.4 * | 26.2 * | 23.8 |
| Binge drinking [#] | 13.7 * | 16.0 | 23.7 * | 18.7 * | 15.3 |
| Marijuana | 11.2 * | 18.4 * | 18.3 * | 17.4 * | 13.0 |
| Cocaine | 0.9 * | 1.4 | 3.2 * | 2.7 * | 1.4 |
| Crack | 0.6 * | 1.4 | 2.3 * | 2.2 * | 1.0 |
| Inhalants | 2.0 | 2.1 | 3.0 * | 3.2 * | 2.2 |
| Amphetamines | 1.6 | 1.1 | 2.5 | 3.2 * | 1.9 |
| Methamphetamines | 0.6 * | 0.9 | 1.9 * | 1.7 * | 0.9 |
| Heroin | 0.5 * | 1.0 | 1.8 * | 1.7 * | 0.8 |
| Steroids | 0.5 * | 0.9 | 1.5 * | 1.6 * | 0.8 |
| Ecstasy | 1.1 * | 1.1 | 2.5 * | 2.6 * | 1.4 |
| Tranquilizers | 1.3 * | 1.1 | 2.8 * | 2.7 * | 1.6 |
| Hallucinogens | 1.2 * | 1.4 | 2.5 * | 2.7 * | 1.5 |
| Prescription painkillers | 4.4 | 2.1 * | 6.8 * | 5.4 | 4.7 |
| Prescription drugs | 4.1 | 2.3 * | 5.8 * | 4.9 | 4.3 |
| Over the counter drugs | 3.2 | 2.1 * | 4.8 * | 4.9 * | 3.5 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

10th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 32.3 | 25.5 * | 37.6 * | 35.4 * | 32.8 |
| Smokeless tobacco | 18.4 * | 4.8 * | 13.3 * | 17.7 | 17.2 |
| Cigars | 21.1 | 13.6 * | 20.2 | 22.7 * | 20.9 |
| Pipe | 14.5 | 9.8 * | 18.7 * | 17.4 * | 15.0 |
| Alcohol | 54.7 | 55.1 | 61.6 * | 57.0 | 55.5 |
| Marijuana | 27.5 * | 40.4 * | 39.9 * | 36.7 * | 30.3 |
| Cocaine | 3.1 * | 3.0 | 8.5 * | 5.5 * | 3.8 |
| Crack | 1.5 * | 2.0 | 4.7 * | 3.3 * | 2.0 |
| Inhalants | 7.1 | 5.2 * | 9.6 * | 9.4 * | 7.4 |
| Amphetamines | 5.9 | 2.8 * | 7.1 * | 7.0 * | 5.9 |
| Methamphetamines | 1.8 * | 2.2 | 3.8 * | 3.4 * | 2.2 |
| Heroin | 1.4 * | 1.8 | 4.4 * | 3.1 * | 1.9 |
| Steroids | 1.2 * | 1.6 | 3.5 * | 3.0 * | 1.7 |
| Ecstasy | 3.7 * | 3.7 | 7.8 * | 6.8 * | 4.4 |
| Tranquilizers | 3.8 * | 3.2 | 7.1 * | 5.7 * | 4.3 |
| Hallucinogens | 4.7 | 3.3 * | 7.1 * | 6.9 * | 5.1 |
| Prescription painkillers | 12.5 | 7.5 * | 13.9 | 14.2 * | 12.6 |
| Prescription drugs | 11.1 | 5.7 * | 12.8 * | 12.6 * | 11.2 |
| Over the counter drugs | 8.4 | 6.1 * | 11.0 * | 10.7 * | 8.8 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

10th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 17.1 | 10.1 * | 18.0 | 20.4 * | 17.2 |
| Smokeless tobacco | 8.3 | 2.1 * | 6.0 * | 9.2 * | 7.9 |
| Cigars | 8.5 | 5.9 * | 9.2 | 10.7 * | 8.7 |
| Pipe | 6.0 * | 4.8 * | 9.8 * | 8.6 * | 6.6 |
| Alcohol | 27.5 * | 27.9 | 32.3 * | 31.2 * | 28.4 |
| Binge drinking [#] | 16.9 * | 17.6 | 24.2 * | 21.8 * | 18.1 |
| Marijuana | 14.5 * | 21.2 * | 23.3 * | 21.3 * | 16.4 |
| Cocaine | 1.1 * | 2.4 * | 3.8 * | 2.6 * | 1.6 |
| Crack | 0.7 * | 1.7 * | 2.7 * | 1.6 * | 1.0 |
| Inhalants | 1.8 * | 1.7 | 3.8 * | 3.1 * | 2.1 |
| Amphetamines | 2.2 | 1.7 | 3.5 * | 2.9 | 2.4 |
| Methamphetamines | 0.7 * | 1.2 | 2.4 * | 1.3 * | 0.9 |
| Heroin | 0.6 * | 1.2 | 2.2 * | 1.5 * | 0.9 |
| Steroids | 0.6 * | 1.2 | 1.9 * | 1.3 * | 0.8 |
| Ecstasy | 1.3 * | 1.8 | 4.0 * | 2.9 * | 1.7 |
| Tranquilizers | 1.7 | 1.7 | 3.7 * | 2.4 | 2.0 |
| Hallucinogens | 1.6 * | 1.7 | 3.1 * | 2.9 * | 1.9 |
| Prescription painkillers | 5.6 | 3.8 * | 6.9 | 7.3 * | 5.8 |
| Prescription drugs | 5.1 | 3.4 * | 6.4 * | 6.1 * | 5.2 |
| Over the counter drugs | 3.3 | 2.8 | 5.0 * | 4.6 * | 3.6 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

11th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 38.2 | 28.3 * | 41.2 * | 39.1 | 38.1 |
| Smokeless tobacco | 21.5 * | 6.1 * | 15.5 * | 19.3 | 20.0 |
| Cigars | 27.8 | 16.8 * | 23.5 * | 27.8 | 27.0 |
| Pipe | 20.1 | 11.7 * | 21.6 | 22.6 * | 20.1 |
| Alcohol | 60.3 | 63.2 | 62.9 | 60.1 | 60.6 |
| Marijuana | 33.3 * | 46.6 * | 40.3 * | 41.6 * | 35.4 |
| Cocaine | 4.2 * | 6.1 | 9.2 * | 7.9 * | 5.1 |
| Crack | 1.8 * | 3.9 * | 4.9 * | 4.9 * | 2.5 |
| Inhalants | 7.4 | 6.2 | 9.2 | 11.3 * | 7.9 |
| Amphetamines | 7.2 | 4.3 * | 9.6 * | 11.5 * | 7.8 |
| Methamphetamines | 2.1 * | 3.0 | 4.7 * | 5.1 * | 2.7 |
| Heroin | 1.7 * | 3.4 * | 4.2 * | 4.2 * | 2.2 |
| Steroids | 1.5 * | 2.3 | 4.2 * | 3.8 * | 2.0 |
| Ecstasy | 4.9 * | 5.5 | 7.9 * | 9.3 * | 5.7 |
| Tranquilizers | 4.7 | 2.8 * | 7.0 * | 7.6 * | 5.2 |
| Hallucinogens | 5.9 * | 4.9 | 8.2 * | 9.7 * | 6.5 |
| Prescription painkillers | 14.6 | 10.2 * | 15.0 | 18.7 * | 14.9 |
| Prescription drugs | 13.0 | 8.1 * | 14.5 | 16.1 * | 13.3 |
| Over the counter drugs | 8.9 | 7.3 * | 10.8 | 12.8 * | 9.5 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

11th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 21.1 | 13.1 * | 20.4 | 21.6 | 20.7 |
| Smokeless tobacco | 9.4 | 2.8 * | 6.5 * | 8.8 | 8.8 |
| Cigars | 11.0 | 8.8 * | 11.0 | 13.1 * | 11.1 |
| Pipe | 7.9 | 8.0 | 10.4 * | 9.8 * | 8.3 |
| Alcohol | 31.5 | 29.5 | 34.1 | 32.4 | 31.7 |
| Binge drinking [#] | 19.9 * | 20.2 | 25.1 * | 24.3 * | 20.8 |
| Marijuana | 16.8 * | 26.7 * | 22.7 * | 23.9 * | 18.6 |
| Cocaine | 1.2 * | 3.5 * | 4.3 * | 3.6 * | 1.8 |
| Crack | 0.7 * | 2.4 * | 3.0 * | 2.6 * | 1.2 |
| Inhalants | 1.5 * | 2.7 | 2.7 * | 3.2 * | 1.8 |
| Amphetamines | 2.3 | 2.4 | 3.4 | 4.0 * | 2.6 |
| Methamphetamines | 0.8 * | 2.0 | 2.7 * | 2.8 * | 1.3 |
| Heroin | 0.8 * | 1.9 | 2.4 * | 2.2 * | 1.1 |
| Steroids | 0.6 * | 1.9 * | 2.2 * | 2.1 * | 1.0 |
| Ecstasy | 1.3 * | 2.4 | 3.5 * | 3.1 * | 1.8 |
| Tranquilizers | 1.6 * | 1.6 | 3.0 * | 3.5 * | 2.0 |
| Hallucinogens | 1.8 * | 2.6 | 3.8 * | 4.0 * | 2.2 |
| Prescription painkillers | 5.5 | 4.4 | 6.5 | 8.4 * | 5.9 |
| Prescription drugs | 5.3 | 4.2 | 6.4 | 7.5 * | 5.6 |
| Over the counter drugs | 3.0 | 3.5 | 3.9 | 5.2 * | 3.3 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

12th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|--------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 42.4 | 28.0 * | 45.1 * | 43.5 | 42.1 |
| Smokeless tobacco | 24.9 * | 7.9 * | 21.1 | 22.0 | 23.6 |
| Cigars | 37.2 * | 20.8 * | 31.8 * | 34.8 | 35.9 |
| Pipe | 27.3 | 16.8 * | 27.7 | 29.7 * | 27.1 |
| Alcohol | 65.7 | 61.0 * | 66.1 | 64.7 | 65.4 |
| Marijuana | 36.9 * | 48.5 * | 40.9 | 45.6 * | 38.6 |
| Cocaine | 5.2 * | 5.1 | 11.6 * | 7.9 * | 5.9 |
| Crack | 1.8 * | 3.3 | 7.3 * | 5.0 * | 2.6 |
| Inhalants | 7.4 | 6.1 | 10.6 * | 9.5 * | 7.8 |
| Amphetamines | 8.1 | 4.5 * | 10.3 * | 10.7 * | 8.3 |
| Methamphetamines | 2.2 * | 3.3 | 6.7 * | 4.4 * | 2.8 |
| Heroin | 1.9 * | 2.8 | 6.0 * | 4.2 * | 2.4 |
| Steroids | 1.6 * | 2.8 | 5.2 * | 3.7 * | 2.1 |
| Ecstasy | 5.9 * | 6.8 | 11.4 * | 9.2 * | 6.7 |
| Tranquilizers | 5.0 | 4.3 | 9.1 * | 7.2 * | 5.4 |
| Hallucinogens | 6.7 | 4.7 * | 10.5 * | 9.7 * | 7.2 |
| Prescription painkillers | 16.0 | 10.1 * | 16.6 | 18.3 * | 16.0 |
| Prescription drugs | 14.6 | 8.9 * | 15.1 | 16.5 * | 14.6 |
| Over the counter drugs | 9.4 | 6.5 * | 12.0 * | 12.1 * | 9.7 |

NOTES: * Statistically significant differences from the Indiana rates (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011

(Values expressed as percentages)

12th Grade

| DRUG | White | Black | Hispanic | Other | Indiana |
|-----------------------------|--------|--------|----------|--------|---------|
| Cigarettes | 25.1 | 13.3 * | 24.5 | 25.2 | 24.6 |
| Smokeless tobacco | 11.2 | 4.1 * | 8.9 * | 11.4 | 10.7 |
| Cigars | 16.7 | 12.8 * | 15.1 | 19.0 * | 16.7 |
| Pipe | 12.1 | 8.9 * | 14.7 * | 14.8 * | 12.5 |
| Alcohol | 38.1 | 33.2 * | 35.7 | 39.6 | 37.8 |
| Binge drinking [#] | 25.8 | 24.8 | 28.6 | 30.6 * | 26.4 |
| Marijuana | 18.2 * | 27.8 * | 21.2 | 27.5 * | 19.8 |
| Cocaine | 1.5 * | 2.7 | 5.0 * | 3.9 * | 2.0 |
| Crack | 0.6 * | 2.3 * | 4.1 * | 2.5 * | 1.2 |
| Inhalants | 1.3 * | 1.7 | 4.2 * | 3.0 * | 1.7 |
| Amphetamines | 2.4 | 1.6 | 4.9 * | 4.1 * | 2.7 |
| Methamphetamines | 0.8 * | 1.9 | 3.9 * | 2.6 * | 1.3 |
| Heroin | 0.8 * | 1.7 | 3.7 * | 2.3 * | 1.2 |
| Steroids | 0.6 * | 1.7 | 3.7 * | 2.3 * | 1.1 |
| Ecstasy | 1.6 * | 2.8 | 5.4 * | 3.3 * | 2.1 |
| Tranquilizers | 1.5 * | 2.0 | 4.9 * | 3.5 * | 2.0 |
| Hallucinogens | 2.0 * | 2.5 | 6.0 * | 3.7 * | 2.4 |
| Prescription painkillers | 6.3 | 5.2 | 8.5 * | 7.8 * | 6.6 |
| Prescription drugs | 5.7 | 3.9 * | 8.1 * | 7.2 * | 5.9 |
| Over the counter drugs | 2.8 | 2.4 | 5.3 * | 4.8 * | 3.2 |

NOTES: * Statistically significant differences from the Indiana rates ($p < .05$).

[#] Binge drinking during the past two weeks.



VI. PREVALENCE OF USE BY GENDER

Prevalence Rates by Gender: Lifetime and Monthly Use

| | |
|------------------|-----|
| • Grade 6 | 94 |
| • Grade 7 | 95 |
| • Grade 8 | 96 |
| • Grade 9 | 97 |
| • Grade 10 | 98 |
| • Grade 11 | 99 |
| • Grade 12 | 100 |

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

6th Grade

| DRUG | Male | Female | Indiana |
|--------------------------|------|--------|---------|
| Cigarettes* | 8.3 | 6.3 | 7.3 |
| Smokeless tobacco* | 4.0 | 1.0 | 2.5 |
| Cigars* | 2.9 | 0.9 | 1.9 |
| Pipe* | 1.7 | 0.9 | 1.3 |
| Alcohol* | 19.9 | 14.8 | 17.4 |
| Marijuana* | 3.9 | 2.3 | 3.1 |
| Cocaine* | 0.7 | 0.4 | 0.6 |
| Crack | 0.6 | 0.5 | 0.6 |
| Inhalants* | 2.2 | 1.4 | 1.8 |
| Amphetamines | 0.5 | 0.3 | 0.4 |
| Methamphetamines | 0.6 | 0.4 | 0.5 |
| Heroin | 0.5 | 0.3 | 0.4 |
| Steroids* | 0.8 | 0.4 | 0.6 |
| Ecstasy* | 0.5 | 0.3 | 0.4 |
| Tranquilizers | 0.6 | 0.4 | 0.5 |
| Hallucinogens | 0.5 | 0.4 | 0.4 |
| Prescription painkillers | 1.2 | 1.0 | 1.1 |
| Prescription drugs* | 1.2 | 0.9 | 1.0 |
| Over the counter drugs | 1.2 | 1.0 | 1.1 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

6th Grade

| DRUG | Male | Female | Indiana |
|--------------------------|------|--------|---------|
| Cigarettes | 2.7 | 2.6 | 2.6 |
| Smokeless tobacco* | 1.5 | 0.4 | 1.0 |
| Cigars* | 0.9 | 0.4 | 0.7 |
| Pipe* | 0.8 | 0.5 | 0.6 |
| Alcohol* | 5.8 | 4.9 | 5.3 |
| Binge drinking# | 6.3 | 5.4 | 5.8 |
| Marijuana* | 1.9 | 1.4 | 1.6 |
| Cocaine* | 0.4 | 0.2 | 0.3 |
| Crack* | 0.3 | 0.2 | 0.3 |
| Inhalants | 0.9 | 0.8 | 0.9 |
| Amphetamines | 0.3 | 0.2 | 0.2 |
| Methamphetamines | 0.3 | 0.2 | 0.3 |
| Heroin | 0.3 | 0.1 | 0.2 |
| Steroids* | 0.3 | 0.1 | 0.2 |
| Ecstasy* | 0.3 | 0.2 | 0.2 |
| Tranquilizers* | 0.3 | 0.2 | 0.3 |
| Hallucinogens | 0.3 | 0.2 | 0.3 |
| Prescription painkillers | 0.6 | 0.6 | 0.6 |
| Prescription drugs | 0.6 | 0.6 | 0.6 |
| Over the counter drugs | 0.7 | 0.7 | 0.7 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).
Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011 (Values expressed as percentages)

7th Grade

| DRUG | Male | Female | Indiana |
|--------------------------|------|--------|---------|
| Cigarettes | 13.7 | 12.7 | 13.2 |
| Smokeless tobacco* | 7.7 | 2.5 | 5.1 |
| Cigars* | 6.1 | 2.5 | 4.3 |
| Pipe* | 4.2 | 2.3 | 3.2 |
| Alcohol | 27.4 | 24.7 | 26.1 |
| Marijuana* | 8.3 | 6.3 | 7.3 |
| Cocaine* | 1.5 | 0.7 | 1.1 |
| Crack | 1.4 | 0.9 | 1.1 |
| Inhalants | 3.2 | 3.2 | 3.2 |
| Amphetamines | 1.3 | 0.8 | 1.0 |
| Methamphetamines* | 1.2 | 0.8 | 1.0 |
| Heroin | 1.0 | 0.6 | 0.8 |
| Steroids* | 1.4 | 0.6 | 1.0 |
| Ecstasy* | 1.3 | 0.8 | 1.1 |
| Tranquilizers | 1.1 | 0.8 | 0.9 |
| Hallucinogens* | 1.2 | 0.6 | 0.9 |
| Prescription painkillers | 2.8 | 2.5 | 2.7 |
| Prescription drugs | 2.4 | 2.3 | 2.4 |
| Over the counter drugs* | 2.6 | 2.9 | 2.8 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011 (Values expressed as percentages)

7th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 5.9 | 6.1 | 6.0 |
| Smokeless tobacco* | 3.4 | 1.0 | 2.2 |
| Cigars* | 2.1 | 1.0 | 1.6 |
| Pipe* | 1.7 | 1.2 | 1.5 |
| Alcohol* | 9.9 | 10.5 | 10.2 |
| Binge drinking# | 7.8 | 8.0 | 7.9 |
| Marijuana* | 4.5 | 3.7 | 4.1 |
| Cocaine* | 0.9 | 0.4 | 0.6 |
| Crack | 0.9 | 0.5 | 0.7 |
| Inhalants | 1.4 | 1.5 | 1.4 |
| Amphetamines | 0.7 | 0.4 | 0.5 |
| Methamphetamines* | 0.6 | 0.4 | 0.5 |
| Heroin | 0.6 | 0.3 | 0.4 |
| Steroids* | 0.7 | 0.3 | 0.5 |
| Ecstasy* | 0.6 | 0.4 | 0.5 |
| Tranquilizers | 0.5 | 0.4 | 0.5 |
| Hallucinogens | 0.6 | 0.4 | 0.5 |
| Prescription painkillers* | 1.5 | 1.4 | 1.5 |
| Prescription drugs | 1.4 | 1.5 | 1.4 |
| Over the counter drugs* | 1.4 | 1.8 | 1.6 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

8th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 20.3 | 20.9 | 20.6 |
| Smokeless tobacco* | 13.0 | 4.4 | 8.6 |
| Cigars* | 11.1 | 5.3 | 8.1 |
| Pipe* | 7.2 | 4.8 | 6.0 |
| Alcohol* | 36.5 | 38.8 | 37.7 |
| Marijuana* | 15.4 | 13.9 | 14.6 |
| Cocaine* | 2.3 | 1.8 | 2.0 |
| Crack* | 1.7 | 1.6 | 1.7 |
| Inhalants* | 5.2 | 5.7 | 5.4 |
| Amphetamines | 2.4 | 2.3 | 2.3 |
| Methamphetamines | 1.6 | 1.4 | 1.5 |
| Heroin* | 1.4 | 1.0 | 1.2 |
| Steroids* | 1.8 | 0.8 | 1.3 |
| Ecstasy* | 2.5 | 1.9 | 2.2 |
| Tranquilizers | 1.9 | 1.9 | 1.9 |
| Hallucinogens* | 2.6 | 2.1 | 2.3 |
| Prescription painkillers* | 5.3 | 5.9 | 5.6 |
| Prescription drugs* | 4.4 | 5.1 | 4.8 |
| Over the counter drugs* | 4.2 | 5.7 | 5.0 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

8th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 9.4 | 10.6 | 10.0 |
| Smokeless tobacco* | 6.2 | 1.6 | 3.9 |
| Cigars* | 4.3 | 2.1 | 3.2 |
| Pipe* | 3.3 | 2.7 | 3.0 |
| Alcohol* | 14.9 | 18.4 | 16.7 |
| Binge drinking# | 10.5 | 11.7 | 11.1 |
| Marijuana* | 8.6 | 8.1 | 8.3 |
| Cocaine | 1.3 | 0.8 | 1.0 |
| Crack* | 1.0 | 0.7 | 0.9 |
| Inhalants* | 2.1 | 2.5 | 2.3 |
| Amphetamines | 1.3 | 1.1 | 1.2 |
| Methamphetamines | 1.0 | 0.8 | 0.9 |
| Heroin | 0.8 | 0.5 | 0.7 |
| Steroids* | 1.0 | 0.4 | 0.7 |
| Ecstasy* | 1.3 | 0.9 | 1.1 |
| Tranquilizers | 1.0 | 1.0 | 1.0 |
| Hallucinogens* | 1.4 | 1.0 | 1.2 |
| Prescription painkillers* | 2.8 | 3.3 | 3.0 |
| Prescription drugs* | 2.4 | 3.0 | 2.7 |
| Over the counter drugs* | 2.1 | 3.2 | 2.7 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).
Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011 (Values expressed as percentages)

9th Grade

| DRUG | Male | Female | Indiana |
|--------------------------|------|--------|---------|
| Cigarettes | 27.3 | 27.1 | 27.2 |
| Smokeless tobacco* | 19.8 | 6.7 | 13.1 |
| Cigars* | 20.6 | 9.4 | 14.9 |
| Pipe* | 12.6 | 8.3 | 10.4 |
| Alcohol* | 46.2 | 49.2 | 47.7 |
| Marijuana* | 24.9 | 21.1 | 23.0 |
| Cocaine* | 3.7 | 2.2 | 2.9 |
| Crack* | 2.4 | 1.5 | 1.9 |
| Inhalants | 7.0 | 6.8 | 6.9 |
| Amphetamines | 4.4 | 4.1 | 4.3 |
| Methamphetamines* | 2.2 | 1.5 | 1.9 |
| Heroin* | 2.0 | 1.2 | 1.6 |
| Steroids* | 2.2 | 0.8 | 1.5 |
| Ecstasy* | 3.8 | 3.0 | 3.4 |
| Tranquilizers | 3.4 | 3.2 | 3.3 |
| Hallucinogens* | 4.3 | 2.9 | 3.6 |
| Prescription painkillers | 9.6 | 9.2 | 9.4 |
| Prescription drugs* | 8.1 | 8.3 | 8.2 |
| Over the counter drugs* | 6.5 | 7.9 | 7.2 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011 (Values expressed as percentages)

9th Grade

| DRUG | Male | Female | Indiana |
|--------------------------|------|--------|---------|
| Cigarettes | 14.1 | 14.6 | 14.4 |
| Smokeless tobacco* | 10.3 | 2.1 | 6.2 |
| Cigars* | 8.5 | 3.5 | 6.0 |
| Pipe* | 5.6 | 3.9 | 4.8 |
| Alcohol* | 22.7 | 25.0 | 23.8 |
| Binge drinking# | 15.5 | 15.1 | 15.3 |
| Marijuana* | 14.1 | 11.8 | 13.0 |
| Cocaine* | 1.9 | 0.9 | 1.4 |
| Crack* | 1.3 | 0.7 | 1.0 |
| Inhalants | 2.2 | 2.3 | 2.2 |
| Amphetamines | 1.9 | 1.9 | 1.9 |
| Methamphetamines* | 1.0 | 0.7 | 0.9 |
| Heroin* | 1.0 | 0.6 | 0.8 |
| Steroids* | 1.1 | 0.4 | 0.8 |
| Ecstasy* | 1.7 | 1.2 | 1.4 |
| Tranquilizers | 1.6 | 1.6 | 1.6 |
| Hallucinogens* | 1.9 | 1.2 | 1.5 |
| Prescription painkillers | 4.5 | 4.9 | 4.7 |
| Prescription drugs | 4.0 | 4.6 | 4.3 |
| Over the counter drugs* | 3.0 | 4.1 | 3.5 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

10th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 34.3 | 31.4 | 32.8 |
| Smokeless tobacco* | 26.7 | 8.3 | 17.2 |
| Cigars* | 28.5 | 13.8 | 20.9 |
| Pipe* | 18.3 | 11.9 | 15.0 |
| Alcohol* | 54.7 | 56.3 | 55.5 |
| Marijuana* | 32.8 | 28.0 | 30.3 |
| Cocaine* | 5.0 | 2.7 | 3.8 |
| Crack* | 2.9 | 1.3 | 2.0 |
| Inhalants* | 8.0 | 7.0 | 7.4 |
| Amphetamines* | 6.4 | 5.5 | 5.9 |
| Methamphetamines* | 2.8 | 1.6 | 2.2 |
| Heroin* | 2.6 | 1.2 | 1.9 |
| Steroids* | 2.5 | 0.8 | 1.7 |
| Ecstasy* | 5.4 | 3.5 | 4.4 |
| Tranquilizers | 4.6 | 4.0 | 4.3 |
| Hallucinogens* | 6.5 | 3.7 | 5.1 |
| Prescription painkillers* | 13.3 | 11.9 | 12.6 |
| Prescription drugs | 11.5 | 10.9 | 11.2 |
| Over the counter drugs | 9.0 | 8.7 | 8.8 |

NOTES: * Differences between males and females are statistically significant (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

10th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 18.5 | 16.1 | 17.2 |
| Smokeless tobacco* | 14.0 | 2.3 | 7.9 |
| Cigars* | 12.8 | 4.9 | 8.7 |
| Pipe* | 8.0 | 5.2 | 6.6 |
| Alcohol | 28.9 | 27.9 | 28.4 |
| Binge drinking# | 19.8 | 16.6 | 18.1 |
| Marijuana* | 18.6 | 14.4 | 16.4 |
| Cocaine* | 2.1 | 1.0 | 1.6 |
| Crack* | 1.4 | 0.6 | 1.0 |
| Inhalants* | 2.4 | 1.9 | 2.1 |
| Amphetamines* | 2.6 | 2.1 | 2.4 |
| Methamphetamines* | 1.3 | 0.6 | 0.9 |
| Heroin* | 1.3 | 0.5 | 0.9 |
| Steroids* | 1.3 | 0.4 | 0.8 |
| Ecstasy* | 2.2 | 1.3 | 1.7 |
| Tranquilizers | 2.1 | 1.8 | 2.0 |
| Hallucinogens* | 2.5 | 1.3 | 1.9 |
| Prescription painkillers* | 6.2 | 5.5 | 5.8 |
| Prescription drugs | 5.4 | 5.1 | 5.2 |
| Over the counter drugs | 3.6 | 3.6 | 3.6 |

NOTES: * Differences between males and females are statistically significant (p < .05).
Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011 (Values expressed as percentages)

11th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 40.0 | 36.3 | 38.1 |
| Smokeless tobacco* | 31.2 | 9.9 | 20.0 |
| Cigars* | 37.0 | 18.0 | 27.0 |
| Pipe* | 24.3 | 16.3 | 20.1 |
| Alcohol* | 60.7 | 60.5 | 60.6 |
| Marijuana* | 38.8 | 32.3 | 35.4 |
| Cocaine* | 7.1 | 3.3 | 5.1 |
| Crack* | 3.5 | 1.5 | 2.5 |
| Inhalants* | 9.0 | 6.9 | 7.9 |
| Amphetamines* | 8.5 | 7.2 | 7.8 |
| Methamphetamines* | 3.6 | 2.0 | 2.7 |
| Heroin* | 3.3 | 1.2 | 2.2 |
| Steroids* | 3.2 | 0.9 | 2.0 |
| Ecstasy* | 7.1 | 4.4 | 5.7 |
| Tranquilizers | 5.4 | 5.0 | 5.2 |
| Hallucinogens* | 8.5 | 4.7 | 6.5 |
| Prescription painkillers* | 16.8 | 13.2 | 14.9 |
| Prescription drugs* | 14.3 | 12.4 | 13.3 |
| Over the counter drugs* | 10.3 | 8.8 | 9.5 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011 (Values expressed as percentages)

11th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 22.1 | 19.5 | 20.7 |
| Smokeless tobacco* | 15.6 | 2.7 | 8.8 |
| Cigars* | 16.7 | 6.0 | 11.1 |
| Pipe* | 10.0 | 6.8 | 8.3 |
| Alcohol* | 33.3 | 30.3 | 31.7 |
| Binge drinking# | 23.8 | 18.2 | 20.8 |
| Marijuana* | 21.6 | 15.8 | 18.6 |
| Cocaine* | 2.7 | 1.0 | 1.8 |
| Crack* | 1.8 | 0.6 | 1.2 |
| Inhalants* | 2.1 | 1.5 | 1.8 |
| Amphetamines* | 2.7 | 2.4 | 2.6 |
| Methamphetamines* | 1.9 | 0.7 | 1.3 |
| Heroin* | 1.7 | 0.6 | 1.1 |
| Steroids* | 1.6 | 0.4 | 1.0 |
| Ecstasy* | 2.4 | 1.2 | 1.8 |
| Tranquilizers | 2.2 | 1.7 | 2.0 |
| Hallucinogens* | 3.1 | 1.4 | 2.2 |
| Prescription painkillers* | 6.8 | 5.1 | 5.9 |
| Prescription drugs* | 6.0 | 5.3 | 5.6 |
| Over the counter drugs* | 3.5 | 3.2 | 3.3 |

NOTES: * Differences between males and females are statistically significant ($p < .05$).

Binge drinking during the past two weeks.

Lifetime Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

12th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 45.1 | 39.2 | 42.1 |
| Smokeless tobacco* | 36.6 | 10.7 | 23.6 |
| Cigars* | 48.1 | 23.9 | 35.9 |
| Pipe* | 32.5 | 21.8 | 27.1 |
| Alcohol | 65.6 | 65.1 | 65.4 |
| Marijuana* | 42.2 | 35.1 | 38.6 |
| Cocaine* | 8.1 | 3.8 | 5.9 |
| Crack* | 3.7 | 1.5 | 2.6 |
| Inhalants* | 9.5 | 6.1 | 7.8 |
| Amphetamines* | 9.8 | 6.9 | 8.3 |
| Methamphetamines* | 3.7 | 1.9 | 2.8 |
| Heroin* | 3.4 | 1.4 | 2.4 |
| Steroids* | 3.4 | 0.9 | 2.1 |
| Ecstasy* | 8.4 | 4.9 | 6.7 |
| Tranquilizers* | 6.1 | 4.8 | 5.4 |
| Hallucinogens* | 9.8 | 4.6 | 7.2 |
| Prescription painkillers* | 19.0 | 13.1 | 16.0 |
| Prescription drugs* | 16.5 | 12.7 | 14.6 |
| Over the counter drugs* | 11.7 | 7.8 | 9.7 |

NOTES: * Differences between males and females are statistically significant (p < .05).

Monthly Use of Alcohol, Tobacco, and Other Drugs, 2011
(Values expressed as percentages)

12th Grade

| DRUG | Male | Female | Indiana |
|---------------------------|------|--------|---------|
| Cigarettes* | 27.9 | 21.4 | 24.6 |
| Smokeless tobacco* | 19.2 | 2.4 | 10.7 |
| Cigars* | 25.1 | 8.4 | 16.7 |
| Pipe* | 15.1 | 9.9 | 12.5 |
| Alcohol* | 40.3 | 35.4 | 37.8 |
| Binge drinking** | 29.9 | 23.0 | 26.4 |
| Marijuana* | 23.7 | 16.0 | 19.8 |
| Cocaine* | 2.9 | 1.2 | 2.0 |
| Crack* | 1.7 | 0.6 | 1.2 |
| Inhalants* | 2.4 | 1.0 | 1.7 |
| Amphetamines* | 3.4 | 2.1 | 2.7 |
| Methamphetamines* | 1.8 | 0.8 | 1.3 |
| Heroin* | 1.8 | 0.6 | 1.2 |
| Steroids* | 1.7 | 0.4 | 1.1 |
| Ecstasy* | 3.0 | 1.2 | 2.1 |
| Tranquilizers | 2.4 | 1.6 | 2.0 |
| Hallucinogens* | 3.6 | 1.3 | 2.4 |
| Prescription painkillers* | 8.3 | 4.9 | 6.6 |
| Prescription drugs* | 6.9 | 4.9 | 5.9 |
| Over the counter drugs* | 4.0 | 2.4 | 3.2 |

NOTES: * Differences between males and females are statistically significant (p < .05).
Binge drinking during the past two weeks.



V. PREVALENCE OF USE BY REGION

Prevalence Rates by Region: Lifetime and Monthly Use

| | |
|------------------|-----|
| • Grade 6 | 102 |
| • Grade 7 | 104 |
| • Grade 8 | 106 |
| • Grade 9 | 108 |
| • Grade 10 | 110 |
| • Grade 11 | 112 |
| • Grade 12 | 114 |

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|-------|---------|--------|-----------|-----------|
| Cigarettes | 7.3 | 5.7 ▼ | 8.2 | 4.1 ▼ | 9.8 ▲ | 7.1 | 11.3 ▲ | 4.5 ▼ | 8.9 ▲ |
| Smokeless tobacco | 2.5 | 1.2 ▼ | 2.5 | 1.9 | 3.1 | 2.1 | 4.1 ▲ | 1.6 ▼ | 4.0 ▲ |
| Cigars | 1.9 | 1.6 | 1.8 | 1.2 | 2.2 | 2.0 | 3.3 ▲ | 1.1 ▼ | 2.1 |
| Pipe | 1.3 | 1.4 | 1.3 | 0.7 | 1.5 | 1.3 | 2.1 ▲ | 0.6 ▼ | 1.5 |
| Alcohol | 17.4 | 17.0 | 19.5 ▲ | 12.6 ▼ | 18.0 | 17.5 | 20.0 ▲ | 14.9 ▼ | 18.3 |
| Marijuana | 3.1 | 2.9 | 2.1 ▼ | 2.1 | 2.9 | 4.4 ▲ | 5.9 ▲ | 1.4 ▼ | 2.8 |
| Cocaine | 0.6 | 0.5 | 0.6 | 0.7 | 0.5 | 0.5 | 1.1 ▲ | 0.3 | 0.6 |
| Crack | 0.6 | 0.5 | 0.7 | 0.5 | 0.5 | 0.5 | 1.2 ▲ | 0.4 | 0.5 |
| Inhalants | 1.8 | 1.5 | 2.6 ▲ | 1.7 | 1.8 | 1.4 | 2.4 | 1.8 | 1.9 |
| Amphetamines | 0.4 | 0.4 | 0.3 | 0.6 | 0.5 | 0.4 | 0.8 | 0.3 | 0.4 |
| Methamphetamines | 0.5 | 0.5 | 0.7 | 0.4 | 0.2 | 0.5 | 1.0 ▲ | 0.4 | 0.5 |
| Heroin | 0.4 | 0.5 | 0.4 | 0.3 | 0.4 | 0.4 | 0.8 ▲ | 0.2 | 0.3 |
| Steroids | 0.6 | 0.7 | 0.7 | 0.4 | 0.7 | 0.4 | 0.6 | 0.3 | 0.7 |
| Ecstasy | 0.4 | 0.4 | 0.5 | 0.4 | 0.4 | 0.5 | 0.6 | 0.2 | 0.4 |
| Tranquilizers | 0.5 | 0.7 | 0.4 | 0.3 | 0.5 | 0.3 | 0.8 | 0.3 | 0.6 |
| Hallucinogens | 0.4 | 0.5 | 0.5 | 0.5 | 0.5 | 0.5 | 0.6 | 0.1 | 0.4 |
| Prescription painkillers | 1.1 | 1.3 | 0.8 | 1.4 | 1.0 | 1.1 | 1.8 ▲ | 0.6 ▼ | 0.9 |
| Prescription drugs | 1.0 | 1.2 | 1.0 | 1.2 | 1.1 | 1.1 | 1.3 | 0.8 | 0.8 |
| Over the counter drugs | 1.1 | 1.5 | 1.0 | 0.7 | 1.0 | 1.2 | 1.7 ▲ | 0.7 | 1.0 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|-------|---------|-------|-----------|-----------|
| Cigarettes | 2.6 | 1.8 ▼ | 2.9 | 1.8 | 3.6 ▲ | 2.9 | 4.4 ▲ | 1.5 ▼ | 2.8 |
| Smokeless tobacco | 1.0 | 0.4 ▼ | 0.8 | 0.4 | 1.3 | 1.1 | 1.7 ▲ | 0.5 ▼ | 1.4 ▲ |
| Cigars | 0.7 | 0.4 ▼ | 0.6 | 0.7 | 0.6 | 0.9 | 1.5 ▲ | 0.3 ▼ | 0.7 |
| Pipe | 0.6 | 0.5 | 0.7 | 0.6 | 0.6 | 0.6 | 1.1 ▲ | 0.4 | 0.6 |
| Alcohol | 5.3 | 5.3 | 4.9 | 3.4 ▼ | 5.1 | 5.8 | 6.6 ▲ | 4.8 | 5.7 |
| Binge drinking [#] | 5.8 | 6.7 ▲ | 4.6 ▼ | 4.8 | 6.6 | 6.1 | 6.5 | 4.5 ▼ | 6.2 |
| Marijuana | 1.6 | 1.5 | 1.1 | 1.1 | 1.5 | 2.5 ▲ | 3.0 ▲ | 0.5 ▼ | 1.4 |
| Cocaine | 0.3 | 0.2 | 0.4 | 0.2 | 0.1 | 0.4 | 0.5 | 0.2 | 0.3 |
| Crack | 0.3 | 0.2 | 0.4 | 0.2 | 0.1 | 0.3 | 0.7 ▲ | 0.2 | 0.1 |
| Inhalants | 0.9 | 0.7 | 1.2 | 1.2 | 0.9 | 0.5 ▼ | 1.1 | 1.1 | 0.8 |
| Amphetamines | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.2 | 0.3 | 0.3 | 0.3 |
| Methamphetamines | 0.3 | 0.2 | 0.4 | 0.2 | 0.1 | 0.4 | 0.3 | 0.3 | 0.2 |
| Heroin | 0.2 | 0.2 | 0.3 | 0.3 | 0.1 | 0.2 | 0.3 | 0.1 | 0.2 |
| Steroids | 0.2 | 0.2 | 0.3 | 0.2 | 0.4 | 0.2 | 0.3 | 0.2 | 0.3 |
| Ecstasy | 0.2 | 0.2 | 0.3 | 0.2 | 0.2 | 0.3 | 0.5 | 0.2 | 0.2 |
| Tranquilizers | 0.3 | 0.4 | 0.3 | 0.2 | 0.2 | 0.1 | 0.5 | 0.2 | 0.3 |
| Hallucinogens | 0.3 | 0.2 | 0.4 | 0.3 | 0.2 | 0.3 | 0.3 | 0.1 | 0.2 |
| Prescription painkillers | 0.6 | 0.6 | 0.4 | 0.7 | 0.8 | 0.6 | 1.0 ▲ | 0.4 | 0.6 |
| Prescription drugs | 0.6 | 0.6 | 0.6 | 0.9 | 0.7 | 0.6 | 0.9 | 0.6 | 0.5 |
| Over the counter drugs | 0.7 | 0.6 | 0.7 | 0.7 | 0.8 | 0.9 | 1.0 | 0.5 | 0.5 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 13.2 | 13.1 | 13.4 | 9.9 ▼ | 15.7 ▲ | 10.5 ▼ | 19.7 ▲ | 9.6 ▼ | 17.0 ▲ |
| Smokeless tobacco | 5.1 | 3.5 ▼ | 4.1 ▼ | 3.6 ▼ | 5.8 | 3.7 ▼ | 7.7 ▲ | 4.6 | 9.4 ▲ |
| Cigars | 4.3 | 4.1 | 3.9 | 3.6 | 4.9 | 3.6 ▼ | 6.7 ▲ | 3.0 ▼ | 5.6 ▲ |
| Pipe | 3.2 | 3.3 | 2.7 | 2.7 | 3.3 | 3.1 | 4.8 ▲ | 2.3 ▼ | 4.1 ▲ |
| Alcohol | 26.1 | 28.7 ▲ | 26.7 | 22.1 ▼ | 29.4 ▲ | 22.9 ▼ | 28.9 ▲ | 22.8 ▼ | 29.6 ▲ |
| Marijuana | 7.3 | 8.8 ▲ | 7.1 | 6.3 | 7.8 | 6.9 | 11.3 ▲ | 3.8 ▼ | 7.8 |
| Cocaine | 1.1 | 1.1 | 1.1 | 0.8 | 1.1 | 1.2 | 1.2 | 0.7 ▼ | 1.5 ▲ |
| Crack | 1.1 | 1.1 | 1.1 | 0.6 | 1.1 | 1.2 | 1.0 | 0.9 | 1.5 |
| Inhalants | 3.2 | 3.8 | 2.6 | 2.5 | 2.6 | 3.6 | 3.6 | 3.0 | 3.4 |
| Amphetamines | 1.0 | 1.3 | 0.7 | 0.5 | 1.4 | 1.0 | 1.0 | 0.8 | 1.6 ▲ |
| Methamphetamines | 1.0 | 1.1 | 0.9 | 0.7 | 0.7 | 1.0 | 0.9 | 0.7 | 1.5 ▲ |
| Heroin | 0.8 | 0.8 | 0.5 | 0.5 | 0.8 | 0.9 | 0.8 | 0.6 | 1.3 ▲ |
| Steroids | 1.0 | 0.8 | 0.7 | 0.8 | 1.2 | 1.0 | 1.1 | 0.8 | 1.4 |
| Ecstasy | 1.1 | 1.4 | 0.7 | 0.8 | 1.1 | 1.2 | 1.1 | 0.6 ▼ | 1.5 ▲ |
| Tranquilizers | 0.9 | 1.0 | 0.8 | 0.4 ▼ | 0.9 | 1.1 | 1.0 | 0.7 | 1.3 ▲ |
| Hallucinogens | 0.9 | 1.2 | 0.4 ▼ | 0.4 ▼ | 0.8 | 1.1 | 1.3 | 0.7 | 1.2 |
| Prescription painkillers | 2.7 | 3.4 ▲ | 2.2 | 1.9 | 2.9 | 2.5 | 3.4 ▲ | 2.0 ▼ | 3.0 |
| Prescription drugs | 2.4 | 2.8 | 2.0 | 2.1 | 2.8 | 2.3 | 2.8 | 1.7 ▼ | 2.8 |
| Over the counter drugs | 2.8 | 3.4 | 2.3 | 1.7 ▼ | 2.7 | 3.0 | 3.4 | 2.1 ▼ | 3.3 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|------|---------|-------|-----------|-----------|
| Cigarettes | 6.0 | 5.8 | 5.7 | 4.1 ▼ | 7.0 | 5.0 ▼ | 8.6 ▲ | 4.9 ▼ | 8.1 ▲ |
| Smokeless tobacco | 2.2 | 1.4 ▼ | 1.7 | 1.6 | 2.5 | 1.5 ▼ | 3.2 ▲ | 2.2 | 4.3 ▲ |
| Cigars | 1.6 | 1.4 | 1.5 | 1.4 | 1.6 | 1.4 | 2.7 ▲ | 1.0 ▼ | 2.0 |
| Pipe | 1.5 | 1.7 | 1.3 | 1.5 | 1.3 | 1.2 | 1.6 | 1.3 | 2.1 ▲ |
| Alcohol | 10.2 | 11.4 | 10.1 | 8.5 ▼ | 11.3 | 8.6 ▼ | 11.1 | 9.7 | 12.0 ▲ |
| Binge drinking [#] | 7.9 | 8.8 | 8.4 | 7.6 | 7.9 | 6.8 ▼ | 9.4 ▲ | 6.7 ▼ | 8.8 |
| Marijuana | 4.1 | 4.9 ▲ | 3.7 | 3.9 | 4.6 | 4.0 | 6.2 ▲ | 2.0 ▼ | 4.6 |
| Cocaine | 0.6 | 0.4 | 0.6 | 0.4 | 0.4 | 0.7 | 0.7 | 0.4 | 0.9 |
| Crack | 0.7 | 0.7 | 0.8 | 0.4 | 0.8 | 0.6 | 0.5 | 0.6 | 0.7 |
| Inhalants | 1.4 | 1.7 | 1.1 | 1.2 | 1.2 | 1.7 | 1.3 | 1.5 | 1.4 |
| Amphetamines | 0.5 | 0.7 | 0.4 | 0.2 | 0.8 | 0.5 | 0.5 | 0.5 | 0.7 |
| Methamphetamines | 0.5 | 0.5 | 0.7 | 0.2 | 0.5 | 0.5 | 0.4 | 0.4 | 0.6 |
| Heroin | 0.4 | 0.3 | 0.4 | 0.3 | 0.5 | 0.4 | 0.3 | 0.3 | 0.9 ▲ |
| Steroids | 0.5 | 0.6 | 0.4 | 0.4 | 0.7 | 0.5 | 0.4 | 0.3 | 0.8 ▲ |
| Ecstasy | 0.5 | 0.6 | 0.4 | 0.2 | 0.7 | 0.5 | 0.5 | 0.4 | 0.7 |
| Tranquilizers | 0.5 | 0.7 | 0.4 | 0.1 | 0.6 | 0.5 | 0.5 | 0.4 | 0.6 |
| Hallucinogens | 0.5 | 0.4 | 0.2 ▼ | 0.3 | 0.4 | 0.7 | 0.5 | 0.6 | 0.8 |
| Prescription painkillers | 1.5 | 1.8 | 0.9 ▼ | 1.2 | 1.6 | 1.6 | 1.8 | 1.0 | 1.9 |
| Prescription drugs | 1.4 | 1.9 ▲ | 1.1 | 1.3 | 1.5 | 1.6 | 1.2 | 1.0 | 1.7 |
| Over the counter drugs | 1.6 | 2.2 ▲ | 1.4 | 0.9 ▼ | 1.7 | 1.6 | 1.7 | 1.1 | 1.9 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 20.6 | 22.4 ▲ | 19.8 | 17.1 ▼ | 21.4 | 18.2 ▼ | 25.0 ▲ | 17.3 ▼ | 23.8 ▲ |
| Smokeless tobacco | 8.6 | 5.9 ▼ | 7.5 ▼ | 6.5 ▼ | 11.6 ▲ | 6.7 ▼ | 9.5 | 8.5 | 12.3 ▲ |
| Cigars | 8.1 | 8.0 | 8.2 | 6.6 ▼ | 8.7 | 7.6 | 10.5 ▲ | 6.1 ▼ | 9.2 ▲ |
| Pipe | 6.0 | 6.9 ▲ | 5.3 | 4.0 ▼ | 6.3 | 6.3 | 7.0 | 4.2 ▼ | 6.6 |
| Alcohol | 37.7 | 42.1 ▲ | 38.2 | 34.3 ▼ | 37.7 | 34.5 ▼ | 41.2 ▲ | 33.5 ▼ | 40.8 ▲ |
| Marijuana | 14.6 | 17.7 ▲ | 13.6 | 12.0 ▼ | 14.0 | 16.2 ▲ | 18.1 ▲ | 8.7 ▼ | 14.7 |
| Cocaine | 2.0 | 2.6 ▲ | 1.5 ▼ | 1.1 ▼ | 1.7 | 2.4 | 2.4 | 1.5 | 2.4 |
| Crack | 1.7 | 2.0 | 1.4 | 0.9 ▼ | 1.5 | 1.9 | 1.9 | 1.4 | 1.9 |
| Inhalants | 5.4 | 5.3 | 5.3 | 4.2 ▼ | 5.6 | 5.5 | 5.7 | 5.3 | 6.0 |
| Amphetamines | 2.3 | 2.7 | 2.3 | 1.4 ▼ | 2.4 | 2.3 | 3.0 ▲ | 1.8 | 2.5 |
| Methamphetamines | 1.5 | 1.8 | 1.3 | 0.8 ▼ | 1.4 | 1.8 | 1.6 | 1.2 | 1.6 |
| Heroin | 1.2 | 1.5 | 0.8 ▼ | 0.5 ▼ | 1.1 | 1.5 | 1.5 | 0.8 | 1.3 |
| Steroids | 1.3 | 1.5 | 1.2 | 0.8 ▼ | 1.5 | 1.4 | 1.6 | 1.0 | 1.3 |
| Ecstasy | 2.2 | 2.5 | 2.0 | 1.2 ▼ | 1.9 | 2.8 ▲ | 2.2 | 1.4 ▼ | 2.4 |
| Tranquilizers | 1.9 | 2.1 | 1.8 | 1.0 ▼ | 2.0 | 2.1 | 2.5 | 1.4 ▼ | 2.1 |
| Hallucinogens | 2.3 | 2.7 | 1.8 | 1.4 ▼ | 2.1 | 2.9 ▲ | 2.4 | 1.8 | 2.4 |
| Prescription painkillers | 5.6 | 6.3 | 5.2 | 4.9 | 5.7 | 6.3 ▲ | 6.3 | 3.8 ▼ | 5.4 |
| Prescription drugs | 4.8 | 5.6 ▲ | 3.8 ▼ | 3.5 ▼ | 5.2 | 5.5 ▲ | 5.2 | 3.4 ▼ | 4.8 |
| Over the counter drugs | 5.0 | 5.4 | 4.4 | 4.0 ▼ | 4.9 | 5.5 | 5.8 | 3.7 ▼ | 5.3 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates (p<.05).
▼ indicates that local prevalence rates are significantly lower than state rates (p<.05).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|-------|---------|--------|-----------|-----------|
| Cigarettes | 10.0 | 10.5 | 9.3 | 8.2 ▼ | 10.8 | 9.4 | 11.4 ▲ | 8.5 ▼ | 11.4 ▲ |
| Smokeless tobacco | 3.9 | 2.5 ▼ | 3.4 | 2.0 ▼ | 5.5 ▲ | 3.5 | 3.8 | 3.6 | 5.4 ▲ |
| Cigars | 3.2 | 3.2 | 3.0 | 2.4 | 3.3 | 3.2 | 4.6 ▲ | 2.0 ▼ | 3.8 ▲ |
| Pipe | 3.0 | 3.7 ▲ | 2.6 | 1.7 ▼ | 2.8 | 3.1 | 3.6 | 2.2 ▼ | 3.4 |
| Alcohol | 16.7 | 19.5 ▲ | 14.5 ▼ | 13.4 ▼ | 17.0 | 16.4 | 18.2 | 15.6 | 17.8 |
| Binge drinking [#] | 11.1 | 12.4 ▲ | 10.6 | 9.0 ▼ | 10.4 | 11.7 | 13.1 ▲ | 9.0 ▼ | 11.7 |
| Marijuana | 8.3 | 11.1 ▲ | 7.0 ▼ | 5.9 ▼ | 8.0 | 9.9 ▲ | 10.1 ▲ | 4.7 ▼ | 8.0 |
| Cocaine | 1.0 | 1.3 | 0.8 | 0.5 ▼ | 1.0 | 1.2 | 1.3 | 0.7 | 1.1 |
| Crack | 0.9 | 1.2 ▲ | 0.6 | 0.4 ▼ | 0.8 | 0.9 | 1.3 ▲ | 0.8 | 0.8 |
| Inhalants | 2.3 | 2.1 | 2.1 | 1.5 ▼ | 2.4 | 2.4 | 2.8 | 2.1 | 2.4 |
| Amphetamines | 1.2 | 1.6 | 1.1 | 0.7 | 1.2 | 1.2 | 1.7 ▲ | 0.9 | 1.1 |
| Methamphetamines | 0.9 | 1.1 | 0.8 | 0.3 ▼ | 0.6 | 1.1 | 1.1 | 0.7 | 0.8 |
| Heroin | 0.7 | 0.9 | 0.4 | 0.2 | 0.6 | 0.9 | 1.1 ▲ | 0.5 | 0.6 |
| Steroids | 0.7 | 0.8 | 0.6 | 0.4 | 0.8 | 0.8 | 0.7 | 0.5 | 0.6 |
| Ecstasy | 1.1 | 1.3 | 1.1 | 0.5 ▼ | 1.1 | 1.3 ▲ | 1.3 | 0.7 | 1.1 |
| Tranquilizers | 1.0 | 1.0 | 1.0 | 0.5 ▼ | 1.1 | 1.2 | 1.3 | 0.7 | 1.0 |
| Hallucinogens | 1.2 | 1.4 | 1.0 | 0.7 | 1.0 | 1.5 ▲ | 1.3 | 0.8 ▼ | 1.2 |
| Prescription painkillers | 3.0 | 3.5 | 2.7 | 2.7 | 3.1 | 3.5 | 3.7 | 1.9 ▼ | 2.8 |
| Prescription drugs | 2.7 | 3.7 ▲ | 2.2 | 2.0 | 2.8 | 3.2 ▲ | 2.9 | 1.7 ▼ | 2.5 |
| Over the counter drugs | 2.7 | 3.1 | 2.1 | 2.2 | 2.8 | 3.2 ▲ | 2.9 | 2.0 ▼ | 2.6 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 27.2 | 30.0 ▲ | 30.0 ▲ | 21.5 ▼ | 29.6 ▲ | 21.3 ▼ | 32.3 ▲ | 27.6 | 31.1 ▲ |
| Smokeless tobacco | 13.1 | 9.9 ▼ | 13.3 | 9.3 ▼ | 16.7 ▲ | 9.3 ▼ | 14.8 ▲ | 14.8 ▲ | 19.6 ▲ |
| Cigars | 14.9 | 16.0 | 16.2 | 11.4 ▼ | 16.9 ▲ | 12.0 ▼ | 17.6 ▲ | 15.2 | 17.0 ▲ |
| Pipe | 10.4 | 11.9 ▲ | 10.4 | 7.3 ▼ | 12.7 ▲ | 9.9 | 11.5 | 10.1 | 10.8 |
| Alcohol | 47.7 | 52.1 ▲ | 48.5 | 44.6 ▼ | 49.0 | 42.9 | 50.6 ▲ | 46.3 | 51.6 ▲ |
| Marijuana | 23.0 | 27.1 ▲ | 22.6 | 19.5 ▼ | 24.7 | 22.9 | 26.8 ▲ | 20.1 ▼ | 22.5 |
| Cocaine | 2.9 | 3.5 | 3.3 | 2.1 ▼ | 2.7 | 3.2 | 2.8 | 2.7 | 2.7 |
| Crack | 1.9 | 1.8 | 2.3 | 1.6 | 1.7 | 2.5 ▲ | 1.8 | 1.9 | 1.5 |
| Inhalants | 6.9 | 8.0 ▲ | 6.8 | 5.0 ▼ | 7.5 | 6.1 | 8.0 | 8.3 ▲ | 6.7 |
| Amphetamines | 4.3 | 5.1 | 4.2 | 2.6 ▼ | 4.8 | 4.1 | 4.6 | 5.0 | 4.3 |
| Methamphetamines | 1.9 | 2.1 | 1.9 | 1.1 ▼ | 2.0 | 2.0 | 1.4 | 2.3 | 1.9 |
| Heroin | 1.6 | 1.8 | 1.5 | 1.2 | 2.0 | 2.1 ▲ | 1.4 | 1.4 | 1.2 |
| Steroids | 1.5 | 1.8 | 1.3 | 1.0 ▼ | 1.6 | 1.5 | 1.5 | 1.9 | 1.6 |
| Ecstasy | 3.4 | 4.2 ▲ | 3.9 | 2.5 ▼ | 2.8 | 4.0 ▲ | 3.3 | 2.6 ▼ | 3.3 |
| Tranquilizers | 3.3 | 4.1 ▲ | 3.4 | 2.0 ▼ | 3.6 | 3.2 | 3.6 | 3.6 | 3.3 |
| Hallucinogens | 3.6 | 3.5 | 3.5 | 2.7 ▼ | 3.5 | 4.0 | 3.7 | 3.6 | 3.9 |
| Prescription painkillers | 9.4 | 10.1 | 10.8 ▲ | 7.2 ▼ | 9.3 | 9.4 | 9.9 | 9.1 | 9.8 |
| Prescription drugs | 8.2 | 9.3 | 9.1 | 6.2 ▼ | 8.6 | 7.6 | 8.2 | 9.4 ▲ | 7.9 |
| Over the counter drugs | 7.2 | 7.4 | 8.8 ▲ | 5.4 ▼ | 8.2 | 6.9 | 7.1 | 7.3 | 7.0 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|-------|---------|--------|-----------|-----------|
| Cigarettes | 14.4 | 15.5 | 14.6 | 10.5 ▼ | 15.8 | 10.8 ▼ | 17.3 ▲ | 15.7 | 17.9 ▲ |
| Smokeless tobacco | 6.2 | 4.4 ▼ | 5.7 | 4.3 ▼ | 7.3 ▲ | 4.6 ▼ | 7.5 ▲ | 6.8 | 9.8 ▲ |
| Cigars | 6.0 | 6.4 | 5.9 | 4.2 ▼ | 6.9 | 4.6 ▼ | 8.7 ▲ | 6.6 | 6.5 |
| Pipe | 4.8 | 5.5 | 4.7 | 3.5 ▼ | 5.4 | 4.7 | 5.7 | 4.2 | 4.8 |
| Alcohol | 23.8 | 27.0 ▲ | 23.2 | 22.1 ▼ | 23.0 | 21.0 ▼ | 24.2 | 24.7 | 27.1 ▲ |
| Binge drinking [#] | 15.3 | 16.8 | 16.4 | 12.9 ▼ | 14.9 | 13.9 ▼ | 15.5 | 16.6 | 16.3 |
| Marijuana | 13.0 | 16.1 ▲ | 12.0 | 11.1 ▼ | 14.2 | 12.8 | 15.9 ▲ | 11.4 ▼ | 11.9 |
| Cocaine | 1.4 | 1.6 | 1.3 | 1.1 | 1.3 | 1.7 | 1.2 | 1.4 | 1.3 |
| Crack | 1.0 | 1.2 | 1.0 | 0.7 | 1.0 | 1.2 | 0.8 | 0.9 | 0.8 |
| Inhalants | 2.2 | 2.5 | 1.8 | 1.6 ▼ | 3.0 ▲ | 2.2 | 2.6 | 2.0 | 2.3 |
| Amphetamines | 1.9 | 2.3 | 1.6 | 1.1 ▼ | 2.0 | 2.3 | 2.0 | 1.8 | 1.9 |
| Methamphetamines | 0.9 | 0.9 | 1.1 | 0.5 ▼ | 0.9 | 0.9 | 0.6 | 0.8 | 0.9 |
| Heroin | 0.8 | 1.1 | 0.8 | 0.5 | 0.9 | 1.1 ▲ | 0.6 | 0.6 | 0.8 |
| Steroids | 0.8 | 1.0 | 0.8 | 0.4 ▼ | 0.7 | 0.9 | 0.5 | 0.7 | 0.8 |
| Ecstasy | 1.4 | 1.5 | 1.6 | 1.0 | 1.0 | 2.0 ▲ | 1.5 | 1.1 | 1.3 |
| Tranquilizers | 1.6 | 1.9 | 1.8 | 0.9 ▼ | 1.5 | 1.7 | 2.0 | 1.8 | 1.4 |
| Hallucinogens | 1.5 | 1.6 | 1.5 | 1.0 ▼ | 1.4 | 1.9 | 1.7 | 1.3 | 1.6 |
| Prescription painkillers | 4.7 | 5.2 | 5.4 | 3.6 ▼ | 4.4 | 4.5 | 5.0 | 4.6 | 4.9 |
| Prescription drugs | 4.3 | 5.1 | 4.5 | 2.9 ▼ | 4.6 | 4.3 | 4.3 | 4.6 | 4.0 |
| Over the counter drugs | 3.5 | 3.8 | 4.3 ▲ | 2.5 ▼ | 3.6 | 3.7 | 3.5 | 3.2 | 3.6 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 32.8 | 33.8 | 32.9 | 27.9 ▼ | 34.0 | 28.2 ▼ | 38.9 ▲ | 30.3 ▼ | 39.1 ▲ |
| Smokeless tobacco | 17.2 | 14.7 ▼ | 14.0 ▼ | 12.4 ▼ | 21.8 ▲ | 12.7 ▼ | 19.0 | 18.2 | 24.7 ▲ |
| Cigars | 20.9 | 20.7 | 21.4 | 17.8 ▼ | 23.6 ▲ | 18.3 ▼ | 22.8 | 19.2 ▼ | 24.5 ▲ |
| Pipe | 15.0 | 18.5 ▲ | 14.0 | 11.3 ▼ | 16.9 ▲ | 14.6 | 14.6 | 12.4 ▼ | 16.9 ▲ |
| Alcohol | 55.5 | 59.8 ▲ | 53.7 | 51.4 ▼ | 54.9 | 53.0 ▼ | 58.4 ▲ | 53.5 ▼ | 60.4 ▲ |
| Marijuana | 30.3 | 34.4 ▲ | 28.4 ▼ | 26.9 ▼ | 29.9 | 32.3 ▲ | 35.4 ▲ | 23.2 ▼ | 31.8 ▲ |
| Cocaine | 3.8 | 4.4 | 3.4 | 2.6 ▼ | 3.9 | 4.1 | 3.7 | 3.1 | 4.5 ▲ |
| Crack | 2.0 | 2.1 | 2.0 | 1.5 ▼ | 1.9 | 2.5 ▲ | 1.7 | 2.1 | 2.0 |
| Inhalants | 7.4 | 7.6 | 7.0 | 6.1 ▼ | 7.6 | 6.7 | 8.5 | 7.9 | 8.6 ▲ |
| Amphetamines | 5.9 | 7.2 ▲ | 5.7 | 3.8 ▼ | 7.1 ▲ | 5.4 | 6.2 | 5.2 | 7.1 ▲ |
| Methamphetamines | 2.2 | 1.8 | 2.3 | 1.5 ▼ | 2.3 | 2.2 | 1.7 | 2.4 | 2.7 ▲ |
| Heroin | 1.9 | 2.2 | 1.5 | 1.3 ▼ | 1.7 | 2.2 | 2.0 | 1.6 | 2.1 |
| Steroids | 1.7 | 1.5 | 1.8 | 1.1 ▼ | 1.7 | 1.8 | 1.7 | 1.7 | 1.8 |
| Ecstasy | 4.4 | 5.7 ▲ | 4.1 | 3.3 ▼ | 4.0 | 4.8 | 4.3 | 4.0 | 5.0 |
| Tranquilizers | 4.3 | 5.4 ▲ | 3.9 | 2.6 ▼ | 4.6 | 4.0 | 4.5 | 4.2 | 5.0 ▲ |
| Hallucinogens | 5.1 | 5.3 | 4.4 | 4.3 | 4.8 | 5.4 | 4.5 | 5.1 | 5.9 ▲ |
| Prescription painkillers | 12.6 | 14.4 ▲ | 11.7 | 10.2 ▼ | 13.3 | 12.3 | 12.5 | 10.8 ▼ | 14.7 ▲ |
| Prescription drugs | 11.2 | 13.8 ▲ | 11.5 | 7.6 ▼ | 12.3 | 10.4 | 10.6 | 11.2 | 12.2 ▲ |
| Over the counter drugs | 8.8 | 9.5 | 8.4 | 7.0 ▼ | 9.4 | 8.3 | 8.5 | 8.1 | 10.7 ▲ |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 17.2 | 16.6 | 16.6 | 14.9 ▼ | 18.7 ▲ | 14.1 ▼ | 20.3 ▲ | 17.3 | 21.0 ▲ |
| Smokeless tobacco | 7.9 | 6.4 ▼ | 6.0 ▼ | 4.6 ▼ | 10.7 ▲ | 5.9 ▼ | 9.2 ▲ | 8.5 | 11.9 ▲ |
| Cigars | 8.7 | 9.3 | 8.1 | 7.3 ▼ | 9.3 | 7.7 ▼ | 9.5 | 8.8 | 10.0 ▲ |
| Pipe | 6.6 | 9.0 ▲ | 5.6 | 4.8 ▼ | 6.5 | 6.6 | 6.4 | 5.8 | 7.5 ▲ |
| Alcohol | 28.4 | 31.2 ▲ | 25.2 ▼ | 26.2 ▼ | 27.7 | 26.3 ▼ | 29.0 | 29.5 | 32.0 ▲ |
| Binge drinking [#] | 18.1 | 20.7 ▲ | 15.8 ▼ | 17.2 | 17.9 | 16.1 ▼ | 18.7 | 19.2 | 20.2 ▲ |
| Marijuana | 16.4 | 18.8 ▲ | 14.5 ▼ | 15.0 ▼ | 15.4 | 18.4 ▲ | 19.4 ▲ | 12.5 ▼ | 16.8 |
| Cocaine | 1.6 | 1.9 | 1.4 | 1.2 | 1.0 ▼ | 1.7 | 1.7 | 1.7 | 1.7 |
| Crack | 1.0 | 1.2 | 0.9 | 0.8 | 0.7 | 1.1 | 1.1 | 1.2 | 1.0 |
| Inhalants | 2.1 | 2.6 | 2.1 | 1.6 | 1.9 | 1.8 | 2.4 | 2.7 | 2.3 |
| Amphetamines | 2.4 | 3.3 ▲ | 2.0 | 1.9 | 2.5 | 2.2 | 2.4 | 2.4 | 2.3 |
| Methamphetamines | 0.9 | 1.1 | 0.9 | 0.6 | 0.9 | 1.0 | 0.8 | 1.3 ▲ | 0.9 |
| Heroin | 0.9 | 1.2 | 0.6 | 0.5 ▼ | 0.8 | 1.0 | 1.0 | 0.9 | 0.9 |
| Steroids | 0.8 | 0.8 | 0.7 | 0.5 | 0.8 | 1.0 | 0.8 | 0.8 | 0.8 |
| Ecstasy | 1.7 | 2.4 ▲ | 1.3 ▼ | 1.4 | 1.5 | 2.0 | 1.8 | 1.4 | 1.9 |
| Tranquilizers | 2.0 | 2.3 | 1.6 | 1.3 ▼ | 2.0 | 1.9 | 2.5 | 1.9 | 2.2 |
| Hallucinogens | 1.9 | 1.8 | 1.3 | 1.7 | 1.6 | 2.1 | 1.2 ▼ | 2.2 | 2.3 |
| Prescription painkillers | 5.8 | 6.9 ▲ | 5.2 | 5.2 | 5.5 | 6.0 | 6.1 | 5.0 | 6.5 |
| Prescription drugs | 5.2 | 7.0 ▲ | 4.8 | 3.9 ▼ | 5.6 | 5.1 | 4.8 | 5.3 | 5.4 |
| Over the counter drugs | 3.6 | 4.0 | 3.1 | 3.6 | 3.3 | 3.5 | 3.5 | 3.5 | 3.9 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|------|---------|--------|-----------|-----------|
| Cigarettes | 38.1 | 38.3 | 38.3 | 34.0 ▼ | 37.9 | 34.1 ▼ | 43.4 ▲ | 38.2 | 43.1 ▲ |
| Smokeless tobacco | 20.0 | 18.2 | 19.2 | 16.4 ▼ | 21.3 | 17.0 ▼ | 22.1 | 22.3 ▲ | 25.1 ▲ |
| Cigars | 27.0 | 25.2 | 25.9 | 23.9 ▼ | 28.2 | 25.9 | 30.3 ▲ | 26.3 | 31.4 ▲ |
| Pipe | 20.1 | 25.0 ▲ | 16.5 ▼ | 15.3 ▼ | 21.6 | 22.1 ▲ | 19.7 | 18.6 | 21.9 ▲ |
| Alcohol | 60.6 | 64.9 ▲ | 59.3 | 57.4 ▼ | 59.2 | 58.2 ▼ | 65.9 ▲ | 59.1 | 63.3 ▲ |
| Marijuana | 35.4 | 39.0 ▲ | 32.3 ▼ | 34.2 | 34.6 | 38.5 ▲ | 40.4 ▲ | 29.3 ▼ | 35.5 |
| Cocaine | 5.1 | 5.7 | 5.0 | 4.4 | 5.5 | 4.9 | 4.6 | 4.1 ▼ | 6.6 ▲ |
| Crack | 2.5 | 2.1 | 2.8 | 2.3 | 2.4 | 2.3 | 2.3 | 1.9 | 3.4 ▲ |
| Inhalants | 7.9 | 8.6 | 7.0 | 7.6 | 8.0 | 7.3 | 9.4 | 7.4 | 8.7 |
| Amphetamines | 7.8 | 10.0 ▲ | 7.5 | 6.4 ▼ | 8.4 | 7.4 | 7.7 | 6.4 ▼ | 8.8 |
| Methamphetamines | 2.7 | 2.3 | 2.5 | 2.3 | 2.8 | 2.7 | 2.1 | 2.7 | 3.9 ▲ |
| Heroin | 2.2 | 2.4 | 2.0 | 2.4 | 2.0 | 2.4 | 2.2 | 1.3 ▼ | 2.9 ▲ |
| Steroids | 2.0 | 1.7 | 1.7 | 1.7 | 2.0 | 2.0 | 2.1 | 2.4 | 2.4 |
| Ecstasy | 5.7 | 6.9 ▲ | 5.2 | 4.7 | 4.6 | 6.4 | 5.7 | 3.9 ▼ | 7.2 ▲ |
| Tranquilizers | 5.2 | 6.8 ▲ | 5.1 | 3.9 ▼ | 5.6 | 4.6 | 5.4 | 4.8 | 5.8 |
| Hallucinogens | 6.5 | 6.6 | 5.6 | 6.2 | 6.5 | 6.7 | 5.9 | 5.7 | 8.2 ▲ |
| Prescription painkillers | 14.9 | 17.1 ▲ | 14.7 | 13.8 | 14.0 | 14.6 | 15.8 | 12.3 ▼ | 17.1 ▲ |
| Prescription drugs | 13.3 | 15.7 ▲ | 13.5 | 11.4 ▼ | 13.1 | 13.0 | 14.0 | 12.1 | 14.0 |
| Over the counter drugs | 9.5 | 9.6 | 9.4 | 8.5 | 9.0 | 9.5 | 11.0 | 8.1 ▼ | 11.0 ▲ |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 20.7 | 20.8 | 20.2 | 18.8 ▼ | 19.5 | 17.5 ▼ | 23.5 ▲ | 21.9 | 24.9 ▲ |
| Smokeless tobacco | 8.8 | 7.5 ▼ | 8.2 | 7.0 ▼ | 10.5 ▲ | 7.2 ▼ | 9.0 | 10.6 ▲ | 11.2 ▲ |
| Cigars | 11.1 | 10.2 | 11.3 | 9.7 ▼ | 9.8 | 10.2 | 13.1 ▲ | 11.4 | 13.5 ▲ |
| Pipe | 8.3 | 11.8 ▲ | 6.9 ▼ | 5.9 ▼ | 8.7 | 8.7 | 7.5 | 7.6 | 9.3 |
| Alcohol | 31.7 | 35.0 ▲ | 28.7 ▼ | 30.2 | 29.5 | 29.2 ▼ | 34.1 | 33.0 | 34.7 ▲ |
| Binge drinking [#] | 20.8 | 21.5 | 19.1 | 19.3 | 17.3 ▼ | 20.0 | 23.2 ▲ | 24.6 ▲ | 21.8 |
| Marijuana | 18.6 | 20.6 ▲ | 16.5 ▼ | 18.4 | 17.3 | 21.1 ▲ | 20.2 | 15.5 ▼ | 18.1 |
| Cocaine | 1.8 | 1.8 | 1.9 | 1.8 | 1.6 | 2.0 | 1.2 | 1.4 | 2.4 |
| Crack | 1.2 | 1.0 | 1.4 | 1.1 | 1.2 | 1.2 | 0.5 ▼ | 0.8 | 1.7 ▲ |
| Inhalants | 1.8 | 1.7 | 1.7 | 1.7 | 1.8 | 1.7 | 2.2 | 1.7 | 2.1 |
| Amphetamines | 2.6 | 3.3 | 2.9 | 2.1 | 2.4 | 2.2 | 2.8 | 2.3 | 3.0 |
| Methamphetamines | 1.3 | 1.0 | 1.0 | 1.1 | 1.3 | 1.5 | 0.7 | 1.4 | 1.7 ▲ |
| Heroin | 1.1 | 0.8 | 1.2 | 1.0 | 1.2 | 1.2 | 1.0 | 0.9 | 1.6 |
| Steroids | 1.0 | 0.9 | 0.8 | 0.7 | 1.1 | 1.0 | 0.9 | 1.1 | 1.3 |
| Ecstasy | 1.8 | 1.8 | 1.7 | 2.1 | 1.6 | 1.9 | 1.7 | 1.3 | 2.1 |
| Tranquilizers | 2.0 | 2.4 | 2.0 | 1.5 | 1.6 | 1.8 | 1.6 | 2.3 | 2.4 |
| Hallucinogens | 2.2 | 1.7 | 1.9 | 2.6 | 2.4 | 2.3 | 1.1 ▼ | 2.0 | 3.1 ▲ |
| Prescription painkillers | 5.9 | 6.8 | 5.8 | 5.9 | 5.4 | 5.4 | 6.0 | 4.7 ▼ | 7.3 ▲ |
| Prescription drugs | 5.6 | 7.0 ▲ | 6.3 | 4.8 | 5.2 | 5.3 | 5.3 | 4.6 | 6.3 |
| Over the counter drugs | 3.3 | 3.3 | 3.1 | 3.1 | 2.7 | 3.1 | 3.6 | 3.0 | 4.5 ▲ |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|--------------------------|---------|-----------|---------------|-----------|--------|---------|--------|-----------|-----------|
| Cigarettes | 42.1 | 40.9 | 39.9 | 39.7 ▼ | 43.5 | 38.9 ▼ | 44.8 | 43.2 | 45.5 ▲ |
| Smokeless tobacco | 23.6 | 20.5 ▼ | 18.7 ▼ | 20.4 ▼ | 26.6 ▲ | 21.0 ▼ | 22.9 | 28.7 ▲ | 26.8 ▲ |
| Cigars | 35.9 | 33.9 | 32.4 ▼ | 35.0 | 39.3 ▲ | 33.3 ▼ | 33.7 | 38.1 ▲ | 38.7 ▲ |
| Pipe | 27.1 | 31.5 ▲ | 21.6 ▼ | 23.6 ▼ | 29.9 ▲ | 28.7 | 22.4 ▼ | 27.8 | 27.4 |
| Alcohol | 65.4 | 66.9 | 61.0 ▼ | 63.5 | 63.7 | 63.9 | 67.5 | 65.9 | 68.9 ▲ |
| Marijuana | 38.6 | 40.2 | 34.4 ▼ | 36.8 | 38.3 | 41.0 ▲ | 41.5 | 33.6 ▼ | 41.3 ▲ |
| Cocaine | 5.9 | 6.2 | 4.5 ▼ | 4.6 ▼ | 5.9 | 6.4 | 5.3 | 5.5 | 7.4 ▲ |
| Crack | 2.6 | 2.6 | 1.6 ▼ | 2.7 | 2.6 | 3.2 | 2.4 | 2.2 | 2.9 |
| Inhalants | 7.8 | 7.4 | 7.0 | 6.2 ▼ | 8.1 | 8.1 | 8.1 | 8.5 | 8.4 |
| Amphetamines | 8.3 | 8.5 | 8.3 | 6.2 ▼ | 8.6 | 9.1 | 7.8 | 7.6 | 9.3 |
| Methamphetamines | 2.8 | 2.7 | 2.0 ▼ | 2.1 | 3.4 | 2.9 | 2.5 | 2.6 | 3.5 ▲ |
| Heroin | 2.4 | 3.0 | 1.2 ▼ | 2.2 | 2.3 | 3.0 | 2.4 | 1.9 | 2.7 |
| Steroids | 2.1 | 2.9 ▲ | 1.1 ▼ | 2.1 | 2.4 | 2.0 | 2.4 | 1.5 | 2.4 |
| Ecstasy | 6.7 | 8.0 ▲ | 5.1 ▼ | 5.8 | 6.4 | 8.0 ▲ | 6.2 | 4.9 ▼ | 7.6 ▲ |
| Tranquilizers | 5.4 | 6.4 | 4.3 ▼ | 4.6 | 6.0 | 6.0 | 5.0 | 4.5 | 5.9 |
| Hallucinogens | 7.2 | 6.4 | 5.4 ▼ | 7.5 | 7.4 | 8.2 | 5.3 ▼ | 7.5 | 7.9 |
| Prescription painkillers | 16.0 | 15.2 | 14.5 | 14.4 | 15.9 | 16.7 | 15.4 | 15.2 | 18.6 ▲ |
| Prescription drugs | 14.6 | 14.8 | 14.1 | 12.3 ▼ | 15.1 | 15.1 | 12.0 ▼ | 14.4 | 16.4 ▲ |
| Over the counter drugs | 9.7 | 10.0 | 8.4 | 8.5 | 9.6 | 10.5 | 9.9 | 9.6 | 10.6 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 2011

| DRUG | Indiana | Northwest | North Central | Northeast | West | Central | East | Southwest | Southeast |
|-----------------------------|---------|-----------|---------------|-----------|--------|---------|-------|-----------|-----------|
| Cigarettes | 24.6 | 23.6 | 21.9 ▼ | 23.9 | 25.5 | 22.2 ▼ | 24.0 | 26.5 ▲ | 27.3 ▲ |
| Smokeless tobacco | 10.7 | 9.8 | 7.1 ▼ | 9.5 | 13.3 ▲ | 9.2 ▼ | 8.8 ▼ | 14.2 ▲ | 12.0 ▲ |
| Cigars | 16.7 | 16.4 | 12.7 ▼ | 15.6 | 20.1 ▲ | 15.5 | 16.7 | 18.4 ▲ | 17.4 |
| Pipe | 12.5 | 15.5 ▲ | 8.6 ▼ | 10.4 ▼ | 14.1 ▲ | 14.3 ▲ | 9.6 ▼ | 13.2 | 11.8 |
| Alcohol | 37.8 | 38.9 | 31.2 ▼ | 38.4 | 36.6 | 35.7 ▼ | 36.6 | 42.2 ▲ | 40.6 ▲ |
| Binge drinking [#] | 26.4 | 27.5 | 21.0 ▼ | 25.4 | 27.5 | 23.6 ▼ | 25.1 | 30.3 ▲ | 28.9 ▲ |
| Marijuana | 19.8 | 20.6 | 16.8 ▼ | 20.3 | 20.9 | 21.0 | 21.8 | 15.9 ▼ | 20.7 |
| Cocaine | 2.0 | 2.6 | 1.5 | 2.1 | 2.0 | 2.1 | 2.3 | 1.7 | 2.0 |
| Crack | 1.2 | 1.5 | 0.7 | 1.4 | 1.2 | 1.3 | 1.0 | 0.9 | 1.1 |
| Inhalants | 1.7 | 2.2 | 1.2 | 1.6 | 1.6 | 1.7 | 1.7 | 1.8 | 1.6 |
| Amphetamines | 2.7 | 3.1 | 2.9 | 2.4 | 2.5 | 2.9 | 2.6 | 2.3 | 2.9 |
| Methamphetamines | 1.3 | 1.7 | 0.8 | 1.3 | 1.5 | 1.4 | 1.0 | 1.2 | 1.2 |
| Heroin | 1.2 | 1.9 ▲ | 0.4 ▼ | 1.4 | 1.0 | 1.4 | 1.0 | 0.9 | 1.3 |
| Steroids | 1.1 | 1.7 ▲ | 0.3 ▼ | 0.9 | 1.2 | 1.2 | 1.0 | 0.9 | 1.1 |
| Ecstasy | 2.1 | 3.1 ▲ | 0.8 ▼ | 1.7 | 2.1 | 2.7 ▲ | 2.0 | 1.7 | 2.1 |
| Tranquilizers | 2.0 | 2.7 ▲ | 1.4 | 1.6 | 2.2 | 2.3 | 1.7 | 1.3 ▼ | 2.1 |
| Hallucinogens | 2.4 | 2.5 | 1.5 ▼ | 2.5 | 2.7 | 3.1 ▲ | 1.9 | 2.2 | 2.5 |
| Prescription painkillers | 6.6 | 6.5 | 6.1 | 5.9 | 7.1 | 6.9 | 6.1 | 5.7 | 7.4 |
| Prescription drugs | 5.9 | 6.4 | 5.8 | 4.5 ▼ | 6.0 | 7.1 ▲ | 4.7 | 5.1 | 6.4 |
| Over the counter drugs | 3.2 | 4.1 ▲ | 2.7 | 3.0 | 3.1 | 3.4 | 3.6 | 2.9 | 3.0 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).



VI. PREVALENCE OF USE BY DSA

Prevalence Rates by Defined Service Area (DSA): Lifetime and Monthly Use

| | |
|------------------|-----|
| • Grade 6 | 118 |
| • Grade 7 | 120 |
| • Grade 8 | 122 |
| • Grade 9 | 124 |
| • Grade 10 | 126 |
| • Grade 11 | 128 |
| • Grade 12 | 130 |

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|-------|-------|-------|-------|--------|-------|--------|-------|--------|--------|--------|-------|--------|-------|
| Cigarettes | 7.3 | 6.2 | 6.2 | 6.3 | 6.8 | 1.9 ▼ | 9.7 ▲ | 11.5 ▲ | 5.1 ▼ | 11.2 ▲ | 5.8 ▼ | 9.2 ▲ | 9.0 ▲ | 4.6 ▼ | 6.8 |
| Smokeless tobacco | 2.5 | 1.4 ▼ | 1.6 ▼ | 1.4 ▼ | 2.4 | 1.7 | 3.1 | 3.8 ▲ | 1.7 ▼ | 1.7 ▼ | 3.1 | 3.2 | 4.4 ▲ | 1.5 ▼ | 3.1 |
| Cigars | 1.9 | 1.4 | 1.8 | 1.0 ▼ | 2.4 | 1.1 | 2.0 | 3.1 ▲ | 2.0 | 2.7 ▲ | 1.6 | 2.1 | 2.3 | 1.2 ▼ | 1.5 |
| Pipe | 1.3 | 1.6 | 1.4 | 0.9 | 1.2 | 1.3 | 1.2 | 1.7 | 0.9 | 1.9 | 0.9 | 1.7 | 1.7 | 0.6 ▼ | 1.2 |
| Alcohol | 17.4 | 18.7 | 16.6 | 16.6 | 15.3 | 11.1 ▼ | 17.8 | 20.9 ▲ | 15.6 | 23.0 ▲ | 13.5 ▼ | 20.6 ▲ | 18.2 | 14.9 ▼ | 16.6 |
| Marijuana | 3.1 | 3.4 | 2.2 ▼ | 1.5 ▼ | 2.9 | 1.7 | 2.7 | 5.9 ▲ | 1.9 ▼ | 8.8 ▲ | 2.6 | 2.4 | 2.8 | 1.4 ▼ | 2.6 |
| Cocaine | 0.6 | 0.4 | 0.7 | 0.1 | 0.9 | 1.5 ▲ | 0.1 | 1.1 ▲ | 0.7 | 0.8 | 0.3 | 0.4 | 0.7 | 0.3 | 0.5 |
| Crack | 0.6 | 0.3 | 0.6 | 0.3 | 0.4 | 1.1 | 0.3 | 1.1 ▲ | 0.8 | 0.6 | 0.4 | 0.5 | 0.6 | 0.3 | 0.6 |
| Inhalants | 1.8 | 1.5 | 1.8 | 2.5 | 2.4 | 1.3 | 1.9 | 2.8 ▲ | 1.3 | 1.9 | 1.1 ▼ | 1.4 | 1.9 | 2.2 | 1.6 |
| Amphetamines | 0.4 | 0.2 | 0.6 | 0.1 | 0.4 | 1.1 ▲ | 0.4 | 0.5 | 0.5 | 0.4 | 0.2 | 0.5 | 0.5 | 0.3 | 0.4 |
| Methamphetamines | 0.5 | 0.5 | 0.6 | 0.4 | 0.4 | 0.9 | 0.0 | 0.9 ▲ | 0.6 | 0.6 | 0.4 | 0.5 | 0.6 | 0.3 | 0.6 |
| Heroin | 0.4 | 0.4 | 0.5 | 0.1 | 0.7 | 0.8 | 0.3 | 0.8 ▲ | 0.6 | 0.4 | 0.2 | 0.2 | 0.5 | 0.2 | 0.3 |
| Steroids | 0.6 | 0.5 | 0.7 | 0.6 | 0.4 | 0.8 | 0.7 | 0.7 | 0.5 | 0.4 | 0.5 | 0.5 | 0.7 | 0.1 | 0.9 |
| Ecstasy | 0.4 | 0.3 | 0.5 | 0.2 | 0.6 | 0.9 | 0.3 | 0.6 | 0.4 | 0.5 | 0.6 | 0.2 | 0.5 | 0.2 | 0.3 |
| Tranquilizers | 0.5 | 0.4 | 0.9 ▲ | 0.1 | 0.3 | 0.8 | 0.4 | 0.7 | 0.4 | 0.2 | 0.3 | 0.3 | 0.7 | 0.2 | 0.6 |
| Hallucinogens | 0.4 | 0.4 | 0.5 | 0.3 | 0.4 | 1.1 ▲ | 0.3 | 0.6 | 0.7 | 0.5 | 0.4 | 0.3 | 0.6 | 0.2 | 0.2 |
| Prescription painkillers | 1.1 | 1.1 | 1.2 | 0.5 | 1.9 ▲ | 1.1 | 1.0 | 1.7 ▲ | 1.1 | 1.9 ▲ | 0.5 ▲ | 0.8 | 1.1 | 0.4 ▼ | 0.9 |
| Prescription drugs | 1.0 | 1.1 | 1.2 | 0.8 | 1.7 | 1.3 | 1.0 | 1.3 | 1.0 | 1.6 ▲ | 0.6 | 0.8 | 1.0 | 0.7 | 0.9 |
| Over the counter drugs | 1.1 | 1.3 | 1.4 | 0.5 | 0.6 | 0.8 | 1.0 | 2.0 ▲ | 1.1 | 2.0 ▲ | 0.9 | 0.8 | 1.1 | 0.8 ▼ | 0.8 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|
| Cigarettes | 2.6 | 2.0 | 2.0 | 1.7 | 3.0 | 0.4 | 3.6 ▲ | 5.1 ▲ | 1.9 ▼ | 4.8 ▲ | 1.9 | 3.3 | 2.4 | 1.7 ▼ | 2.4 |
| Smokeless tobacco | 1.0 | 0.4 ▼ | 0.6 | 0.5 | 0.7 | 0.6 | 1.4 | 1.3 | 0.8 | 0.8 | 1.7 ▲ | 1.2 | 1.5 ▲ | 0.4 ▼ | 1.2 |
| Cigars | 0.7 | 0.4 | 0.3 ▼ | 0.4 | 1.3 ▲ | 0.4 | 0.6 | 1.3 ▲ | 0.5 | 1.6 ▲ | 0.4 | 1.0 | 0.5 | 0.4 | 0.6 |
| Pipe | 0.6 | 0.3 | 0.7 | 0.4 | 0.7 | 0.6 | 0.7 | 1.0 | 0.7 | 0.7 | 0.4 | 1.0 | 0.7 | 0.3 | 0.6 |
| Alcohol | 5.3 | 6.3 | 4.4 | 4.3 | 4.3 | 2.6 ▼ | 5.4 | 6.4 | 4.0 ▼ | 9.0 ▲ | 3.6 ▼ | 6.6 | 5.5 | 5.1 | 5.1 |
| Binge drinking# | 5.8 | 8.0 ▲ | 5.6 | 4.0 ▼ | 5.8 | 4.7 | 6.4 | 5.5 | 5.1 | 8.8 ▲ | 4.3 ▼ | 6.1 | 5.8 | 4.3 ▼ | 6.2 |
| Marijuana | 1.6 | 1.8 | 1.2 | 0.5 | 1.7 | 0.8 | 1.5 | 3.2 ▲ | 1.0 ▼ | 5.5 ▲ | 1.4 | 1.0 | 1.3 | 0.5 ▼ | 1.4 |
| Cocaine | 0.3 | 0.2 | 0.2 | 0.1 | 0.6 | 0.6 | 0.1 | 0.5 | 0.5 | 0.5 | 0.1 | 0.3 | 0.3 | 0.2 | 0.2 |
| Crack | 0.3 | 0.1 | 0.2 | 0.1 | 0.3 | 0.6 | 0.1 | 0.7 ▲ | 0.7 ▲ | 0.4 | 0.1 | 0.2 | 0.2 | 0.2 | 0.2 |
| Inhalants | 0.9 | 0.6 | 0.9 | 1.0 | 1.9 ▲ | 0.6 | 1.0 | 1.3 | 0.7 | 0.5 | 0.3 | 0.7 | 0.9 | 1.3 ▲ | 0.7 |
| Amphetamines | 0.2 | 0.0 | 0.3 | 0.1 | 0.4 | 0.6 | 0.2 | 0.4 | 0.3 | 0.2 | 0.1 | 0.1 | 0.3 | 0.2 | 0.3 |
| Methamphetamines | 0.3 | 0.1 | 0.4 | 0.1 | 0.6 | 0.6 | 0.0 | 0.4 | 0.5 | 0.4 | 0.3 | 0.2 | 0.3 | 0.2 | 0.2 |
| Heroin | 0.2 | 0.1 | 0.2 | 0.1 | 0.7 | 0.8 | 0.1 | 0.4 | 0.4 | 0.2 | 0.1 | 0.0 | 0.2 | 0.1 | 0.1 |
| Steroids | 0.2 | 0.1 | 0.1 | 0.5 | 0.4 | 0.4 | 0.4 | 0.1 | 0.2 | 0.2 | 0.1 | 0.2 | 0.2 | 0.1 | 0.4 |
| Ecstasy | 0.2 | 0.1 | 0.2 | 0.2 | 0.4 | 0.4 | 0.2 | 0.4 | 0.4 | 0.2 | 0.1 | 0.2 | 0.2 | 0.3 | 0.2 |
| Tranquilizers | 0.3 | 0.3 | 0.4 | 0.1 | 0.4 | 0.6 | 0.3 | 0.4 | 0.3 | 0.1 | 0.1 | 0.2 | 0.3 | 0.2 | 0.3 |
| Hallucinogens | 0.3 | 0.2 | 0.2 | 0.3 | 0.4 | 0.8 | 0.1 | 0.4 | 0.5 ▲ | 0.2 | 0.1 | 0.2 | 0.3 | 0.2 | 0.1 |
| Prescription painkillers | 0.6 | 0.4 | 0.6 | 0.2 | 1.3 ▲ | 0.6 | 0.8 | 1.0 | 0.8 | 0.9 | 0.3 | 0.5 | 0.6 | 0.2 | 0.6 |
| Prescription drugs | 0.6 | 0.5 | 0.6 | 0.5 | 1.2 | 0.9 | 0.6 | 1.0 | 0.7 | 0.9 | 0.3 | 0.6 | 0.7 | 0.5 | 0.4 |
| Over the counter drugs | 0.7 | 0.5 | 0.6 | 0.4 | 0.7 | 0.6 | 0.8 | 1.3 ▲ | 0.8 | 1.6 ▲ | 0.4 | 0.6 | 0.5 | 0.5 | 0.4 |

NOTES: # Binge drinking during the past two weeks.
 ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|-------|------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 13.2 | 13.4 | 14.1 | 12.5 | 11.9 | 9.2 ▼ | 17.4 ▲ | 19.7 ▲ | 8.2 ▼ | 18.1 ▲ | 12.6 | 15.3 ▲ | 15.2 ▲ | 8.9 ▼ | 15.9 ▲ |
| Smokeless tobacco | 5.1 | 3.5 ▼ | 4.5 | 4.1 | 3.0 ▼ | 2.9 ▼ | 8.4 ▲ | 6.8 ▲ | 2.4 ▼ | 4.5 | 4.4 | 7.2 ▲ | 8.1 ▲ | 4.4 | 8.1 ▲ |
| Cigars | 4.3 | 4.3 | 4.3 | 4.4 | 3.2 | 3.2 | 4.3 | 6.1 ▲ | 2.9 ▼ | 7.3 ▲ | 4.1 | 5.7 ▲ | 5.5 ▲ | 2.4 ▼ | 4.6 |
| Pipe | 3.2 | 2.9 | 3.5 | 3.0 | 2.5 | 2.5 | 3.8 | 4.8 ▲ | 2.7 | 4.5 | 3.0 | 3.5 | 4.0 ▲ | 2.1 ▼ | 3.2 |
| Alcohol | 26.1 | 30.3 ▲ | 26.5 | 26.4 | 26.7 | 20.4 ▼ | 29.0 | 30.5 ▲ | 20.8 ▼ | 37.2 ▲ | 22.9 ▼ | 27.2 | 28.6 ▲ | 21.6 ▼ | 29.7 ▲ |
| Marijuana | 7.3 | 9.8 ▲ | 7.9 | 6.2 | 7.0 | 6.2 | 7.5 | 11.1 ▲ | 5.2 ▼ | 22.4 ▲ | 6.3 | 7.7 | 8.1 | 3.7 ▼ | 6.1 |
| Cocaine | 1.1 | 0.8 | 1.5 | 0.5 | 1.0 | 0.7 | 0.9 | 1.5 | 1.1 | 2.4 ▲ | 1.2 | 0.8 | 1.7 | 0.6 ▼ | 0.8 |
| Crack | 1.1 | 0.8 | 1.5 | 0.7 | 1.0 | 0.4 | 1.2 | 1.3 | 1.1 | 2.7 ▲ | 1.1 | 0.6 | 1.6 | 1.0 | 1.0 |
| Inhalants | 3.2 | 2.9 | 3.9 | 2.9 | 1.7 ▼ | 2.0 ▼ | 4.9 | 4.4 ▲ | 3.2 | 3.0 | 3.2 | 3.1 | 2.9 | 3.0 | 3.6 |
| Amphetamines | 1.0 | 1.0 | 1.5 ▲ | 0.7 | 0.5 | 0.3 | 1.4 | 1.0 | 0.8 | 1.5 | 0.9 | 1.2 | 1.4 | 0.7 | 1.6 ▲ |
| Methamphetamines | 1.0 | 1.0 | 1.2 | 0.8 | 0.5 | 0.5 | 0.6 | 1.2 | 0.6 | 2.7 ▲ | 0.9 | 0.7 | 1.6 ▲ | 0.6 | 1.1 |
| Heroin | 0.8 | 0.7 | 0.9 | 0.5 | 0.5 | 0.3 | 1.2 | 0.8 | 0.8 | 1.2 | 0.9 | 0.5 | 1.3 ▲ | 0.5 | 0.9 |
| Steroids | 1.0 | 0.7 | 0.8 | 0.8 | 0.5 | 0.5 | 1.7 | 1.4 | 1.0 | 1.8 | 1.0 | 0.8 | 1.2 | 0.6 | 1.5 ▲ |
| Ecstasy | 1.1 | 1.1 | 1.4 | 0.9 | 0.3 | 0.7 | 1.2 | 1.3 | 1.1 | 2.7 ▲ | 0.8 | 1.0 | 1.5 ▲ | 0.5 ▼ | 1.0 |
| Tranquilizers | 0.9 | 0.7 | 1.1 | 0.6 | 0.7 | 0.4 | 0.9 | 1.1 | 0.8 | 2.4 ▲ | 0.9 | 1.1 | 1.3 | 0.7 | 1.1 |
| Hallucinogens | 0.9 | 0.8 | 1.2 | 0.3 | 0.5 | 0.3 | 1.4 | 1.3 | 0.9 | 1.2 | 0.9 | 1.2 | 1.2 | 0.8 | 0.8 |
| Prescription painkillers | 2.7 | 3.2 | 3.3 | 1.8 | 2.3 | 1.5 ▼ | 3.8 | 4.1 ▲ | 2.2 | 3.6 | 2.7 | 2.8 | 2.5 | 1.8 ▼ | 3.1 |
| Prescription drugs | 2.4 | 2.9 | 2.5 | 2.1 | 2.0 | 1.6 | 4.3 ▲ | 3.1 ▲ | 1.7 ▼ | 5.1 ▲ | 1.9 | 3.0 | 2.7 | 1.4 ▼ | 2.6 |
| Over the counter drugs | 2.8 | 3.3 | 3.3 | 2.1 | 2.5 | 1.1 ▼ | 3.2 | 4.1 ▲ | 2.3 | 5.1 ▲ | 2.9 | 2.3 | 3.2 | 2.0 ▼ | 3.1 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 7th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|-----------------------------|---------|--------|-------|------|-------|-------|-------|--------|-------|--------|-------|-------|-------|-------|--------|
| Cigarettes | 6.0 | 5.6 | 6.3 | 6.1 | 4.6 | 3.0 ▼ | 9.0 ▲ | 9.4 ▲ | 3.7 ▼ | 7.3 | 5.4 | 7.2 | 7.6 ▲ | 4.2 ▼ | 7.5 ▲ |
| Smokeless tobacco | 2.2 | 1.5 | 1.7 | 2.0 | 1.0 ▼ | 1.3 ▼ | 2.9 | 3.1 ▲ | 0.9 ▼ | 1.5 | 1.4 ▼ | 3.6 ▲ | 3.6 ▲ | 1.9 | 4.0 ▲ |
| Cigars | 1.6 | 1.2 | 1.7 | 1.5 | 1.2 | 1.1 | 1.2 | 2.9 ▲ | 1.0 ▼ | 2.4 | 1.5 | 1.9 | 2.0 | 0.8 ▼ | 1.8 |
| Pipe | 1.5 | 1.5 | 1.9 | 1.0 | 1.4 | 1.4 | 1.7 | 1.9 | 1.1 | 0.9 | 1.0 | 1.1 | 1.7 | 1.1 | 2.2 ▲ |
| Alcohol | 10.2 | 11.9 ▲ | 10.6 | 10.1 | 9.3 | 7.2 ▼ | 13.0 | 12.7 ▲ | 7.3 ▼ | 16.6 ▲ | 8.0 ▼ | 10.6 | 11.3 | 9.3 | 12.5 ▲ |
| Binge drinking [#] | 7.9 | 9.0 | 8.9 | 7.8 | 9.3 | 6.3 ▼ | 8.7 | 10.6 ▲ | 5.5 ▼ | 15.4 ▲ | 6.2 ▼ | 8.6 | 7.4 | 6.7 ▼ | 9.1 |
| Marijuana | 4.1 | 5.1 | 4.6 | 3.7 | 3.7 | 3.5 | 5.2 | 6.4 ▲ | 3.2 | 11.8 ▲ | 3.1 ▼ | 4.0 | 4.7 | 1.8 ▼ | 3.9 |
| Cocaine | 0.6 | 0.1 | 0.9 | 0.4 | 0.3 | 0.3 | 0.6 | 1.0 ▲ | 0.4 | 1.2 | 0.7 | 0.5 | 1.0 ▲ | 0.4 | 0.5 |
| Crack | 0.7 | 0.3 | 1.2 ▲ | 0.5 | 0.5 | 0.3 | 1.2 | 0.7 | 0.4 | 1.5 | 0.6 | 0.5 | 0.9 | 0.5 | 0.7 |
| Inhalants | 1.4 | 1.4 | 1.9 | 1.2 | 1.0 | 0.8 | 1.4 | 1.5 | 1.5 | 0.6 | 1.4 | 1.7 | 1.2 | 1.6 | 1.7 |
| Amphetamines | 0.5 | 0.6 | 0.9 ▲ | 0.5 | 0.2 | 0.2 | 0.6 | 0.5 | 0.3 | 0.6 | 0.4 | 0.8 | 0.6 | 0.4 | 0.9 |
| Methamphetamines | 0.5 | 0.4 | 0.9 ▲ | 0.5 | 0.3 | 0.3 | 0.3 | 0.4 | 0.2 | 1.2 | 0.4 | 0.6 | 0.7 | 0.3 | 0.7 |
| Heroin | 0.4 | 0.1 | 0.6 | 0.5 | 0.5 | 0.3 | 0.6 | 0.3 | 0.3 | 0.3 | 0.4 | 0.5 | 0.9 ▲ | 0.3 | 0.6 |
| Steroids | 0.5 | 0.3 | 0.7 | 0.5 | 0.2 | 0.2 | 1.2 | 0.4 | 0.4 | 0.3 | 0.6 | 0.7 | 0.7 | 0.2 | 0.8 |
| Ecstasy | 0.5 | 0.4 | 0.7 | 0.8 | 0.2 | 0.3 | 0.9 | 0.3 | 0.3 | 0.9 | 0.2 | 0.6 | 0.8 | 0.3 | 0.7 |
| Tranquilizers | 0.5 | 0.4 | 0.9 ▲ | 0.5 | 0.0 | 0.1 | 0.6 | 0.5 | 0.4 | 0.0 | 0.4 | 0.9 ▲ | 0.5 | 0.3 | 0.7 |
| Hallucinogens | 0.5 | 0.4 | 0.4 | 0.3 | 0.2 | 0.3 | 0.9 | 0.4 | 0.6 | 0.0 | 0.6 | 0.7 | 0.8 ▲ | 0.5 | 0.5 |
| Prescription painkillers | 1.5 | 1.8 | 1.6 | 0.8 | 1.0 | 0.7 ▼ | 2.0 | 2.2 ▲ | 1.2 | 1.2 | 1.5 | 2.0 | 1.6 | 0.7 ▼ | 2.0 |
| Prescription drugs | 1.4 | 2.1 | 1.6 | 1.3 | 1.2 | 0.9 | 2.3 | 1.6 | 1.0 | 2.7 ▲ | 1.3 | 1.9 | 1.6 | 0.8 ▼ | 1.6 |
| Over the counter drugs | 1.6 | 2.3 | 2.0 | 1.1 | 1.4 | 0.5 ▼ | 2.0 | 2.3 ▲ | 1.1 | 2.1 | 1.7 | 1.7 | 1.8 | 1.0 ▼ | 1.8 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|-------|--------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 20.6 | 24.1 ▲ | 21.4 | 19.7 | 19.1 | 12.1 ▼ | 22.6 | 25.3 ▲ | 13.9 ▼ | 24.4 ▲ | 20.2 | 22.9 ▲ | 21.6 | 17.3 ▼ | 22.4 ▲ |
| Smokeless tobacco | 8.6 | 5.2 ▼ | 8.9 | 6.1 ▼ | 7.9 | 3.6 ▼ | 12.7 ▲ | 9.6 | 4.9 ▼ | 4.3 ▼ | 11.6 ▲ | 10.7 ▲ | 11.7 ▲ | 8.1 | 10.7 ▲ |
| Cigars | 8.1 | 7.9 | 8.3 | 7.8 | 7.1 | 4.8 ▼ | 9.2 | 11.2 ▲ | 6.7 ▼ | 7.6 | 8.3 | 10.2 ▲ | 8.7 | 6.3 ▼ | 8.2 |
| Pipe | 6.0 | 7.1 ▲ | 6.6 | 4.9 | 4.5 ▼ | 3.6 | 6.2 | 7.4 ▲ | 5.7 | 7.7 ▲ | 5.8 | 6.7 | 5.9 | 3.9 ▼ | 6.4 |
| Alcohol | 37.7 | 44.8 ▲ | 38.7 | 40.5 ▲ | 36.7 | 30.6 ▼ | 38.7 | 40.2 ▲ | 30.9 ▼ | 43.7 ▲ | 34.6 ▼ | 38.7 | 36.5 | 33.6 ▼ | 40.9 ▲ |
| Marijuana | 14.6 | 19.7 ▲ | 14.7 | 14.7 | 12.8 | 11.4 ▼ | 14.5 | 17.7 ▲ | 11.2 ▼ | 28.0 ▲ | 13.8 | 14.8 | 13.4 | 8.7 ▼ | 13.7 |
| Cocaine | 2.0 | 2.5 | 2.5 | 1.5 | 0.6 ▼ | 1.0 ▼ | 1.5 | 2.4 | 2.0 | 3.0 ▲ | 1.8 | 2.7 | 2.2 | 1.4 ▼ | 2.3 |
| Crack | 1.7 | 1.4 | 2.4 ▲ | 1.3 | 1.0 | 0.8 ▼ | 1.3 | 1.8 | 1.9 | 1.9 | 1.7 | 1.8 | 1.8 | 1.4 | 1.6 |
| Inhalants | 5.4 | 5.3 | 5.4 | 5.9 | 4.1 | 3.0 ▼ | 5.6 | 6.3 | 5.1 | 6.0 | 5.7 | 5.7 | 5.4 | 4.7 | 6.4 ▲ |
| Amphetamines | 2.3 | 2.4 | 3.3 ▲ | 1.7 | 1.8 | 1.1 ▼ | 2.1 | 3.7 ▲ | 2.1 | 2.4 | 2.1 | 2.8 | 2.4 | 1.9 | 2.0 |
| Methamphetamines | 1.5 | 1.1 | 2.3 ▲ | 1.1 | 1.0 | 0.5 | 1.2 | 1.8 | 1.6 | 1.9 | 1.7 | 1.9 | 1.6 | 1.1 | 1.4 |
| Heroin | 1.2 | 1.1 | 1.7 ▲ | 0.6 ▼ | 0.6 | 0.5 | 1.0 | 1.4 | 1.4 | 1.4 | 1.4 | 1.5 | 1.3 | 0.7 | 1.0 |
| Steroids | 1.3 | 1.0 | 1.9 ▲ | 0.9 | 0.8 | 0.6 | 1.5 | 1.7 | 1.7 | 1.6 | 1.1 | 1.2 | 1.3 | 0.9 | 1.3 |
| Ecstasy | 2.2 | 2.4 | 2.5 | 1.9 | 1.6 | 0.8 ▼ | 1.5 | 2.3 | 2.3 | 3.7 ▲ | 2.1 | 2.5 | 2.0 | 1.4 ▼ | 2.6 |
| Tranquilizers | 1.9 | 1.8 | 2.5 | 1.7 | 0.9 ▼ | 0.8 ▼ | 2.0 | 2.7 ▲ | 1.9 | 2.5 | 2.0 | 2.3 | 1.9 | 1.3 ▼ | 1.9 |
| Hallucinogens | 2.3 | 2.6 | 2.6 | 1.7 | 1.2 ▼ | 1.2 ▼ | 2.2 | 2.4 | 2.7 | 3.0 | 2.6 | 2.9 | 2.0 | 1.7 | 2.4 |
| Prescription painkillers | 5.6 | 6.6 | 6.0 | 5.5 | 4.5 | 4.1 ▼ | 6.5 | 7.1 ▲ | 5.1 | 7.7 ▲ | 6.1 | 6.0 | 5.0 | 3.8 ▼ | 4.7 ▼ |
| Prescription drugs | 4.8 | 5.8 ▲ | 5.5 | 4.0 | 2.9 ▼ | 3.0 ▼ | 5.9 ▲ | 5.6 | 4.8 | 6.0 ▲ | 4.8 | 5.5 | 4.3 | 3.4 ▼ | 4.5 |
| Over the counter drugs | 5.0 | 5.6 | 5.0 | 4.2 | 4.1 | 3.1 ▼ | 5.4 | 6.8 ▲ | 4.7 | 6.6 ▲ | 4.9 | 5.4 | 4.8 | 3.6 ▼ | 4.9 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 8th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|-----------------------------|---------|--------|-------|-------|--------|--------|-------|--------|--------|--------|-------|-------|-------|-------|--------|
| Cigarettes | 10.0 | 10.8 | 10.8 | 9.0 | 8.9 | 4.9 ▼ | 11.2 | 12.9 ▲ | 6.8 ▼ | 11.8 ▲ | 11.1 | 10.9 | 10.9 | 8.1 ▼ | 10.4 |
| Smokeless tobacco | 3.9 | 2.6 ▼ | 3.8 | 2.3 ▼ | 2.9 | 0.9 ▼ | 6.2 ▲ | 4.3 | 2.6 ▼ | 2.0 ▼ | 5.8 ▲ | 4.5 | 5.3 ▲ | 3.4 | 4.6 ▲ |
| Cigars | 3.2 | 3.4 | 2.9 | 2.8 | 2.1 ▼ | 1.8 ▼ | 3.5 | 5.0 ▲ | 2.5 ▼ | 4.0 | 3.4 | 3.8 | 3.4 | 2.1 ▼ | 3.4 |
| Pipe | 3.0 | 3.8 ▲ | 3.6 | 2.6 | 2.0 | 1.4 ▼ | 2.7 | 3.5 | 2.8 | 4.2 ▲ | 2.4 | 3.4 | 3.2 | 2.1 ▼ | 3.1 |
| Alcohol | 16.7 | 20.8 ▲ | 17.6 | 15.1 | 13.8 ▼ | 11.0 ▼ | 17.2 | 18.5 ▲ | 13.8 ▼ | 20.9 ▲ | 16.4 | 16.7 | 15.6 | 15.5 | 18.8 ▲ |
| Binge drinking [#] | 11.1 | 13.4 ▲ | 11.2 | 11.0 | 9.8 | 8.7 ▼ | 10.2 | 12.8 ▲ | 8.3 ▼ | 17.5 ▲ | 10.8 | 11.7 | 10.5 | 8.8 ▼ | 12.1 |
| Marijuana | 8.3 | 12.4 ▲ | 8.7 | 7.4 | 5.7 ▼ | 5.7 ▼ | 8.0 | 10.2 ▲ | 6.7 ▼ | 18.1 ▲ | 7.7 | 7.8 | 7.9 | 4.9 ▼ | 7.1 ▼ |
| Cocaine | 1.0 | 1.1 | 1.5 ▲ | 0.7 | 0.3 | 0.4 | 0.7 | 1.3 | 1.1 | 1.7 ▲ | 0.9 | 1.2 | 1.1 | 0.7 | 1.1 |
| Crack | 0.9 | 1.0 | 1.3 ▲ | 0.5 | 0.2 | 0.5 | 0.8 | 1.1 | 0.8 | 1.1 | 0.8 | 0.9 | 1.0 | 0.8 | 0.6 |
| Inhalants | 2.3 | 2.0 | 2.4 | 2.3 | 1.4 | 1.3 ▼ | 2.6 | 2.5 | 2.1 | 2.9 | 2.4 | 2.6 | 1.9 | 2.0 | 2.9 ▲ |
| Amphetamines | 1.2 | 1.4 | 1.8 ▲ | 1.0 | 0.8 | 0.5 ▼ | 1.0 | 2.1 ▲ | 1.4 | 1.4 | 0.9 | 1.2 | 1.1 | 0.9 | 0.9 |
| Methamphetamines | 0.9 | 0.7 | 1.4 ▲ | 0.7 | 0.7 | 0.0 | 0.5 | 1.2 | 0.9 | 1.4 | 1.0 | 0.8 | 0.8 | 0.7 | 0.8 |
| Heroin | 0.7 | 0.7 | 1.0 | 0.2 | 0.4 | 0.0 | 0.6 | 1.2 ▲ | 0.8 | 1.2 ▲ | 0.6 | 0.7 | 0.6 | 0.5 | 0.6 |
| Steroids | 0.7 | 0.5 | 1.1 ▲ | 0.4 | 0.4 | 0.2 | 0.9 | 0.9 | 0.9 | 0.9 | 0.6 | 0.5 | 0.6 | 0.6 | 0.7 |
| Ecstasy | 1.1 | 1.3 | 1.4 | 1.1 | 1.0 | 0.3 | 0.7 | 1.2 | 1.2 | 1.8 ▲ | 1.0 | 1.1 | 0.9 | 0.7 | 1.3 |
| Tranquilizers | 1.0 | 0.9 | 1.4 | 0.9 | 0.3 | 0.4 | 1.1 | 1.8 ▲ | 0.9 | 1.8 ▲ | 1.1 | 1.0 | 0.8 | 0.7 | 1.0 |
| Hallucinogens | 1.2 | 1.4 | 1.4 | 0.9 | 0.6 | 0.7 | 1.0 | 1.5 | 1.3 | 1.9 ▲ | 1.5 | 1.1 | 0.9 | 0.8 | 1.1 |
| Prescription painkillers | 3.0 | 3.5 | 3.5 | 2.4 | 2.6 | 2.2 | 3.7 | 4.1 ▲ | 3.0 | 4.5 ▲ | 3.3 | 3.3 | 2.5 | 2.1 ▼ | 2.5 |
| Prescription drugs | 2.7 | 3.4 | 3.6 ▲ | 2.6 | 1.4 ▼ | 1.8 | 3.4 | 3.2 | 2.9 | 3.9 ▲ | 2.5 | 2.7 | 2.2 | 1.9 ▼ | 2.2 |
| Over the counter drugs | 2.7 | 3.1 | 2.9 | 2.2 | 1.6 ▼ | 1.7 | 2.8 | 3.4 | 2.9 | 4.1 ▲ | 2.7 | 2.8 | 2.2 | 2.0 | 2.7 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 27.2 | 29.9 | 30.1 ▲ | 30.7 ▲ | 29.4 | 20.2 ▼ | 33.6 ▲ | 30.6 ▲ | 18.0 ▼ | 20.0 ▼ | 27.7 | 30.3 ▲ | 28.0 | 29.6 ▲ | 29.2 |
| Smokeless tobacco | 13.1 | 6.0 ▼ | 13.0 | 12.9 | 11.7 | 7.9 ▼ | 21.4 ▲ | 14.1 | 8.0 ▼ | 3.5 ▼ | 17.0 ▲ | 15.6 ▲ | 17.1 ▲ | 15.6 ▲ | 18.0 ▲ |
| Cigars | 14.9 | 13.3 | 16.5 ▲ | 17.4 ▲ | 13.6 | 11.4 ▼ | 19.7 ▲ | 16.3 | 11.6 ▼ | 9.2 ▼ | 16.2 | 16.8 ▲ | 15.3 | 17.2 ▲ | 15.7 |
| Pipe | 10.4 | 9.3 | 12.4 ▲ | 9.4 | 10.1 | 7.5 ▼ | 11.3 | 11.3 | 10.5 | 8.8 | 11.5 | 10.6 | 10.2 | 11.9 | 9.7 |
| Alcohol | 47.7 | 57.7 ▲ | 49.0 | 49.5 | 48.1 | 44.5 ▼ | 51.3 | 47.6 | 36.6 ▼ | 46.5 | 47.0 | 52.1 ▲ | 48.8 | 47.6 | 48.3 |
| Marijuana | 23.0 | 30.5 ▲ | 23.8 | 22.6 | 23.0 | 20.7 ▼ | 21.7 | 24.7 | 16.6 ▼ | 29.5 ▲ | 24.6 | 22.4 | 22.9 | 22.4 | 20.2 ▼ |
| Cocaine | 2.9 | 1.7 ▼ | 4.1 ▲ | 3.2 | 3.9 | 2.0 ▼ | 2.9 | 2.9 | 2.7 | 2.6 | 3.3 | 3.1 | 2.9 | 3.1 | 2.1 |
| Crack | 1.9 | 0.5 | 2.2 | 1.6 | 3.0 ▲ | 1.3 ▼ | 2.3 | 2.2 | 2.2 | 2.1 | 2.7 ▲ | 2.2 | 1.4 | 2.0 | 1.5 |
| Inhalants | 6.9 | 7.1 | 7.8 | 7.5 | 7.2 | 4.5 ▼ | 9.0 | 7.7 | 5.6 ▼ | 3.6 ▼ | 8.6 ▲ | 7.5 | 5.9 ▼ | 8.4 ▲ | 8.2 |
| Amphetamines | 4.3 | 2.6 ▼ | 5.9 ▲ | 4.6 | 4.0 | 2.2 ▼ | 6.7 ▲ | 4.2 | 3.2 ▼ | 2.6 ▼ | 4.8 | 5.4 ▲ | 3.9 | 5.1 | 4.9 |
| Methamphetamines | 1.9 | 0.9 | 2.5 ▲ | 1.2 | 2.4 | 0.9 ▼ | 2.9 | 1.7 | 2.1 | 1.4 | 2.0 | 2.0 | 1.9 | 2.7 ▲ | 1.6 |
| Heroin | 1.6 | 0.8 | 2.3 ▲ | 1.2 | 1.3 | 1.2 | 2.6 | 1.4 | 1.9 | 1.5 | 2.2 | 2.0 | 1.4 | 1.4 | 1.1 |
| Steroids | 1.5 | 1.2 | 2.1 ▲ | 0.7 ▼ | 1.3 | 0.9 ▼ | 0.9 | 1.7 | 1.6 | 1.0 | 1.8 | 1.4 | 1.8 | 2.1 | 1.5 |
| Ecstasy | 3.4 | 3.1 | 4.3 ▲ | 4.6 ▲ | 4.0 | 2.0 ▼ | 2.3 | 3.8 | 2.7 | 4.3 | 3.8 | 3.8 | 3.2 | 3.1 | 2.2 ▼ |
| Tranquilizers | 3.3 | 3.0 | 4.6 ▲ | 3.1 | 3.2 | 1.9 ▼ | 4.6 | 3.4 | 2.8 | 2.0 ▼ | 3.6 | 3.8 | 3.1 | 4.0 | 3.0 |
| Hallucinogens | 3.6 | 2.8 | 3.5 | 3.4 | 4.0 | 2.8 | 4.3 | 3.5 | 3.8 | 3.0 | 4.1 | 4.0 | 3.8 | 4.3 | 3.0 |
| Prescription painkillers | 9.4 | 8.1 | 10.6 | 12.6 ▲ | 11.0 | 6.8 ▼ | 11.9 | 9.4 | 7.5 ▼ | 8.1 | 11.2 ▲ | 10.9 ▲ | 8.2 ▼ | 9.8 | 9.3 |
| Prescription drugs | 8.2 | 7.2 | 9.8 ▲ | 11.5 ▲ | 9.1 | 5.3 ▼ | 9.0 | 8.2 | 6.3 ▼ | 6.2 ▼ | 9.1 | 9.0 | 7.4 | 10.4 ▲ | 7.9 |
| Over the counter drugs | 7.2 | 6.4 | 7.8 | 9.0 ▲ | 9.4 ▲ | 5.1 ▼ | 7.2 | 7.6 | 5.6 ▼ | 5.5 ▼ | 9.0 ▲ | 8.1 | 6.8 | 7.6 | 6.9 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 9th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|------|--------|--------|--------|--------|--------|-------|-------|--------|--------|--------|
| Cigarettes | 14.4 | 12.3 | 16.6 ▲ | 14.5 | 14.2 | 9.4 ▼ | 18.0 | 17.2 ▲ | 8.2 ▼ | 8.9 ▼ | 15.4 | 15.9 | 16.0 ▲ | 16.3 ▲ | 17.8 ▲ |
| Smokeless tobacco | 6.2 | 1.9 ▼ | 6.1 | 4.8 | 5.7 | 3.5 ▼ | 7.8 | 7.2 | 3.3 ▼ | 1.9 ▼ | 8.0 ▲ | 7.4 ▲ | 8.7 ▲ | 7.1 | 8.9 ▲ |
| Cigars | 6.0 | 5.5 | 6.5 | 5.2 | 5.7 | 4.1 ▼ | 7.0 | 7.8 ▲ | 4.9 | 3.3 ▼ | 5.7 | 7.4 ▲ | 6.1 | 7.2 | 6.7 |
| Pipe | 4.8 | 4.3 | 5.6 | 4.1 | 5.0 | 3.8 ▼ | 3.2 | 5.9 ▲ | 5.1 | 4.2 | 5.3 | 4.9 | 4.5 | 4.7 | 4.5 |
| Alcohol | 23.8 | 27.2 ▲ | 25.9 ▲ | 25.3 | 21.7 | 22.6 | 19.7 | 23.4 | 16.4 ▼ | 22.8 | 24.3 | 25.0 | 25.4 | 24.6 | 25.2 |
| Binge drinking# | 15.3 | 17.0 | 16.1 | 16.1 | 16.9 | 13.2 ▼ | 10.4 ▼ | 15.7 | 11.8 ▼ | 15.8 | 15.4 | 15.9 | 15.5 | 16.9 | 15.3 |
| Marijuana | 13.0 | 17.0 ▲ | 14.4 | 10.9 ▼ | 12.8 | 12.1 | 12.5 | 15.1 ▲ | 9.4 ▼ | 17.7 ▲ | 13.2 | 12.0 | 12.4 | 12.8 | 10.8 ▼ |
| Cocaine | 1.4 | 0.8 | 1.8 | 0.9 | 1.4 | 1.1 | 2.0 | 1.5 | 1.8 | 1.1 | 1.6 | 1.5 | 1.2 | 1.5 | 1.1 |
| Crack | 1.0 | 0.8 | 1.2 | 0.5 | 1.2 | 0.7 | 0.9 | 0.9 | 1.2 | 0.9 | 1.3 | 1.1 | 0.9 | 0.9 | 0.6 |
| Inhalants | 2.2 | 2.1 | 2.5 | 1.7 | 1.9 | 1.4 ▼ | 4.3 ▲ | 2.2 | 2.2 | 1.2 ▼ | 3.0 ▲ | 2.7 | 2.2 | 2.2 | 2.2 |
| Amphetamines | 1.9 | 0.9 ▼ | 2.8 ▲ | 1.4 | 1.8 | 1.0 ▼ | 3.5 ▲ | 1.8 | 1.8 | 1.4 ▼ | 2.2 | 2.7 ▲ | 1.9 | 2.0 | 1.3 |
| Methamphetamines | 0.9 | 0.2 | 1.1 | 0.9 | 1.1 | 0.6 | 1.4 | 0.8 | 0.9 | 0.5 | 1.2 | 0.9 | 1.1 | 0.8 | 0.6 |
| Heroin | 0.8 | 0.6 | 1.2 | 0.7 | 0.6 | 0.6 | 1.7 | 0.5 | 1.2 | 0.5 | 1.1 | 1.0 | 0.9 | 0.7 | 0.3 |
| Steroids | 0.8 | 0.6 | 1.1 | 0.6 | 1.1 | 0.3 | 0.6 | 0.7 | 1.1 | 0.7 | 0.9 | 0.6 | 0.9 | 0.8 | 0.6 |
| Ecstasy | 1.4 | 0.9 | 1.7 | 1.4 | 1.9 | 0.9 ▼ | 1.4 | 1.7 | 1.2 | 2.1 | 1.7 | 2.0 ▲ | 1.3 | 1.3 | 0.7 |
| Tranquilizers | 1.6 | 1.8 | 2.1 | 1.4 | 2.0 | 0.8 ▼ | 2.6 | 1.6 | 1.6 | 1.1 | 1.6 | 2.2 | 1.5 | 2.0 | 1.0 |
| Hallucinogens | 1.5 | 1.2 | 1.7 | 1.4 | 2.1 | 1.2 | 2.9 ▲ | 1.3 | 1.9 | 1.2 | 1.8 | 1.9 | 1.3 | 1.4 | 1.3 |
| Prescription painkillers | 4.7 | 3.8 | 5.4 | 6.6 ▲ | 5.2 | 3.1 ▼ | 6.1 | 4.9 | 4.0 | 3.3 ▼ | 5.8 ▲ | 5.5 | 3.9 | 5.1 | 4.2 |
| Prescription drugs | 4.3 | 3.8 | 5.4 ▲ | 5.7 ▲ | 4.1 | 2.5 ▼ | 4.3 | 4.1 | 3.5 | 2.7 ▼ | 5.5 ▲ | 5.3 ▲ | 3.8 | 5.1 | 3.7 |
| Over the counter drugs | 3.5 | 2.7 | 4.0 | 4.3 | 4.3 | 2.2 ▼ | 3.8 | 3.6 | 2.8 | 3.2 | 4.6 ▲ | 4.4 ▲ | 3.4 | 3.4 | 3.0 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 32.8 | 32.4 | 36.6 ▲ | 34.2 | 31.7 | 25.6 ▼ | 32.1 | 37.7 ▲ | 27.6 ▼ | 26.3 ▼ | 33.0 | 36.6 ▲ | 36.6 ▲ | 30.0 ▼ | 36.0 ▲ |
| Smokeless tobacco | 17.2 | 12.0 ▼ | 18.4 | 17.3 | 12.6 ▼ | 10.0 ▼ | 23.1 ▲ | 17.1 | 13.6 ▼ | 5.4 ▼ | 20.6 ▲ | 21.2 ▲ | 22.7 ▲ | 17.4 | 22.8 ▲ |
| Cigars | 20.9 | 18.3 ▼ | 24.0 ▲ | 23.0 | 20.0 | 15.9 ▼ | 22.1 | 24.1 ▲ | 19.0 | 14.0 ▼ | 22.4 | 23.5 ▲ | 23.8 ▲ | 19.3 | 21.8 |
| Pipe | 15.0 | 18.1 ▲ | 18.6 ▲ | 14.5 | 12.7 ▼ | 11.0 ▼ | 14.3 | 15.7 | 16.1 | 12.7 ▼ | 16.0 | 14.7 | 15.9 | 12.9 ▼ | 15.8 |
| Alcohol | 55.5 | 63.5 ▲ | 57.6 | 53.9 | 55.8 | 50.1 ▼ | 53.2 | 56.4 | 48.5 ▼ | 58.7 ▲ | 53.3 | 57.0 | 57.8 ▲ | 54.7 | 57.2 |
| Marijuana | 30.3 | 38.2 ▲ | 32.3 | 25.7 ▼ | 30.2 | 27.0 ▼ | 28.1 | 35.1 ▲ | 27.1 ▼ | 39.0 ▲ | 30.3 | 28.6 | 30.5 | 25.3 ▼ | 28.1 ▼ |
| Cocaine | 3.8 | 3.2 | 4.8 ▲ | 3.5 | 3.4 | 2.2 ▼ | 3.5 | 3.4 | 5.4 ▲ | 3.5 | 3.9 | 4.5 | 4.0 | 3.1 | 4.1 |
| Crack | 2.0 | 1.6 | 2.5 | 1.9 | 1.6 | 1.4 ▼ | 1.8 | 1.9 | 3.1 ▲ | 2.0 | 2.4 | 2.0 | 1.9 | 2.2 | 2.1 |
| Inhalants | 7.4 | 5.4 ▼ | 9.3 ▲ | 7.7 | 6.6 | 5.2 ▼ | 7.1 | 9.1 ▲ | 7.8 | 5.5 ▼ | 8.1 | 7.4 | 7.6 | 7.9 | 8.2 |
| Amphetamines | 5.9 | 4.4 ▼ | 9.7 ▲ | 5.9 | 4.7 | 3.0 ▼ | 6.5 | 6.7 | 6.1 | 3.5 ▼ | 6.6 | 6.5 | 6.3 | 5.5 | 6.1 |
| Methamphetamines | 2.2 | 1.1 ▼ | 2.5 | 2.3 | 1.9 | 1.2 ▼ | 2.7 | 2.0 | 2.6 | 1.7 | 2.3 | 2.4 | 2.3 | 2.6 | 2.4 |
| Heroin | 1.9 | 1.7 | 2.2 | 1.0 | 1.8 | 1.1 ▼ | 1.8 | 1.7 | 2.3 | 1.7 | 2.5 | 2.7 ▲ | 1.8 | 1.9 | 1.7 |
| Steroids | 1.7 | 1.2 | 1.7 | 1.1 | 1.8 | 1.0 ▼ | 1.6 | 1.7 | 2.3 ▲ | 1.3 | 2.4 ▲ | 1.3 | 1.7 | 1.7 | 1.8 |
| Ecstasy | 4.4 | 4.8 | 5.9 ▲ | 4.2 | 4.1 | 2.9 ▼ | 3.3 | 4.3 | 5.3 | 4.8 | 4.3 | 4.8 | 4.3 | 4.2 | 4.6 |
| Tranquilizers | 4.3 | 3.7 | 6.6 ▲ | 3.5 | 3.3 | 2.1 ▼ | 4.9 | 4.8 | 4.4 | 3.0 ▼ | 4.5 | 4.4 | 4.5 | 4.6 | 4.4 |
| Hallucinogens | 5.1 | 3.7 ▼ | 5.9 | 5.8 | 4.1 | 4.1 ▼ | 5.1 | 4.4 | 6.1 ▲ | 5.0 | 4.7 | 5.2 | 5.6 | 5.0 | 5.2 |
| Prescription painkillers | 12.6 | 12.1 | 16.1 ▲ | 13.3 | 10.9 | 9.3 ▼ | 13.6 | 13.1 | 11.8 | 10.6 ▼ | 14.1 ▲ | 12.4 | 13.1 | 11.6 | 13.2 |
| Prescription drugs | 11.2 | 10.7 | 16.2 ▲ | 12.9 | 10.5 | 6.8 ▼ | 12.4 | 10.7 | 10.7 | 8.8 ▼ | 12.2 | 10.6 | 10.5 | 11.6 | 12.0 |
| Over the counter drugs | 8.8 | 7.8 | 10.8 ▲ | 9.1 | 7.0 ▼ | 6.7 ▼ | 8.9 | 9.4 | 8.7 | 7.0 ▼ | 9.8 | 8.3 | 10.2 ▲ | 8.3 | 9.5 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 10th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 17.2 | 14.5 ▼ | 19.6 ▲ | 18.5 | 15.1 ▼ | 13.9 ▼ | 18.3 | 20.2 ▲ | 13.3 ▼ | 12.1 ▼ | 17.3 | 19.5 ▲ | 19.8 ▲ | 17.0 | 19.4 ▲ |
| Smokeless tobacco | 7.9 | 4.6 ▼ | 8.9 | 7.3 | 4.8 ▼ | 4.0 ▼ | 12.4 ▲ | 6.6 ▼ | 5.8 ▼ | 2.1 ▼ | 9.0 | 12.2 ▲ | 11.3 ▲ | 8.4 | 10.2 ▲ |
| Cigars | 8.7 | 8.4 | 10.0 ▲ | 8.0 | 7.5 | 7.1 ▼ | 9.4 | 9.0 | 7.5 | 5.9 ▼ | 9.1 | 10.6 ▲ | 9.7 | 9.0 | 9.1 |
| Pipe | 6.6 | 9.4 ▲ | 8.2 ▲ | 5.7 | 5.5 | 5.0 ▼ | 6.3 | 6.0 | 6.1 | 6.6 | 7.0 | 6.4 | 6.7 | 6.1 | 6.8 |
| Alcohol | 28.4 | 32.2 ▲ | 29.9 | 25.5 | 25.6 ▼ | 25.8 ▼ | 26.0 | 27.4 | 24.9 ▼ | 30.0 | 25.0 ▼ | 31.7 ▲ | 29.6 | 29.1 | 31.7 ▲ |
| Binge drinking# | 18.1 | 20.3 | 20.7 ▲ | 16.1 | 15.9 ▼ | 17.2 | 17.6 | 17.3 | 14.5 ▼ | 18.0 | 15.8 ▼ | 21.1 ▲ | 18.0 | 18.3 | 21.2 ▲ |
| Marijuana | 16.4 | 20.4 ▲ | 16.9 | 13.5 ▼ | 15.5 | 15.5 | 15.0 | 18.6 ▲ | 15.1 | 24.3 ▲ | 15.0 | 16.2 | 15.8 | 13.5 ▼ | 15.4 |
| Cocaine | 1.6 | 1.6 | 1.9 | 1.3 | 0.8 ▲ | 1.3 | 1.0 | 1.5 | 1.8 | 1.8 | 1.5 | 1.8 | 1.6 | 1.8 | 1.5 |
| Crack | 1.0 | 1.1 | 1.2 | 0.7 | 0.8 | 0.8 | 0.7 | 0.9 | 0.9 | 1.1 | 1.2 | 1.2 | 0.8 | 1.3 | 0.9 |
| Inhalants | 2.1 | 2.1 | 2.6 | 2.2 | 1.9 | 1.3 ▼ | 2.1 | 2.6 | 2.1 | 1.6 | 2.0 | 2.0 | 2.0 | 2.8 ▲ | 2.2 |
| Amphetamines | 2.4 | 2.4 | 4.1 ▲ | 1.4 | 2.2 | 1.5 ▼ | 2.5 | 2.5 | 2.8 | 1.5 ▼ | 2.4 | 2.4 | 2.1 | 2.6 | 2.1 |
| Methamphetamines | 0.9 | 0.9 | 1.1 | 0.7 | 0.9 | 0.5 ▼ | 1.2 | 0.8 | 0.9 | 0.8 | 1.1 | 0.9 | 0.9 | 1.7 ▲ | 0.7 |
| Heroin | 0.9 | 1.2 | 1.1 | 0.3 | 0.6 | 0.5 ▼ | 0.8 | 0.8 | 0.8 | 0.9 | 1.0 | 1.5 ▲ | 0.9 | 1.0 | 0.7 |
| Steroids | 0.8 | 1.0 | 0.7 | 0.6 | 0.5 | 0.5 | 0.9 | 0.9 | 1.1 | 0.9 | 1.0 | 0.7 | 1.1 | 0.9 | 0.6 |
| Ecstasy | 1.7 | 2.3 | 1.9 | 1.2 | 1.5 | 1.2 | 1.7 | 1.5 | 1.9 | 2.3 | 1.9 | 2.1 | 1.9 | 1.4 | 1.4 |
| Tranquilizers | 2.0 | 2.1 | 2.4 | 0.9 ▼ | 2.1 | 1.0 ▼ | 2.2 | 2.2 | 2.5 | 1.3 ▼ | 2.2 | 2.6 | 1.9 | 2.0 | 1.9 |
| Hallucinogens | 1.9 | 1.6 | 1.7 | 1.9 | 1.8 | 1.7 | 1.9 | 0.8 ▼ | 2.0 | 2.3 | 1.8 | 2.1 | 2.2 | 2.0 | 2.2 |
| Prescription painkillers | 5.8 | 5.8 | 7.6 ▲ | 5.5 | 4.9 | 4.7 ▼ | 6.3 | 6.7 | 5.8 | 5.9 | 5.9 | 6.1 | 5.1 | 4.9 | 6.4 |
| Prescription drugs | 5.2 | 5.4 | 7.7 ▲ | 4.1 | 5.1 | 3.5 ▼ | 5.8 | 5.2 | 5.6 | 4.5 | 5.5 | 5.4 | 4.3 ▼ | 5.5 | 5.4 |
| Over the counter drugs | 3.6 | 3.0 | 4.2 | 3.0 | 3.1 | 3.4 | 3.6 | 3.9 | 4.1 | 2.9 | 4.0 | 3.5 | 3.6 | 3.5 | 3.5 |

NOTES: # Binge drinking during the past two weeks.
 ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 38.1 | 33.9 ▼ | 40.0 | 39.3 | 41.0 | 32.3 ▼ | 38.4 | 42.2 ▲ | 32.7 ▼ | 26.4 ▼ | 42.5 ▼ | 39.0 | 40.4 ▲ | 38.7 | 40.4 |
| Smokeless tobacco | 20.0 | 10.6 ▼ | 22.9 ▲ | 22.2 | 15.2 ▼ | 14.2 ▼ | 22.8 | 20.1 | 16.7 ▼ | 6.2 ▼ | 27.2 ▲ | 23.0 ▲ | 22.4 ▲ | 22.6 ▲ | 24.5 ▲ |
| Cigars | 27.0 | 16.1 ▼ | 29.2 | 26.9 | 26.9 | 23.3 ▼ | 23.5 | 28.4 | 23.7 ▼ | 19.6 ▼ | 32.3 ▲ | 29.9 ▲ | 30.2 ▲ | 27.4 | 27.8 |
| Pipe | 20.1 | 18.9 | 25.0 ▲ | 14.1 ▼ | 17.3 | 15.4 ▼ | 15.3 ▼ | 20.0 | 20.9 | 19.4 | 23.5 ▲ | 19.7 | 22.4 ▲ | 20.0 | 19.4 |
| Alcohol | 60.6 | 63.6 | 63.2 ▲ | 60.0 | 62.9 | 57.4 ▼ | 55.4 | 61.6 | 51.8 ▼ | 64.3 ▲ | 60.8 | 63.3 | 61.9 | 62.2 | 55.8 ▼ |
| Marijuana | 35.4 | 38.4 | 37.0 | 31.1 ▼ | 36.6 | 34.9 | 24.8 ▼ | 40.4 ▲ | 30.0 ▼ | 45.8 ▲ | 38.5 ▲ | 32.4 ▼ | 37.2 | 32.1 ▼ | 28.5 ▼ |
| Cocaine | 5.1 | 3.9 | 6.4 ▲ | 3.4 ▼ | 5.8 | 3.9 ▼ | 4.6 | 5.4 | 5.4 | 3.7 | 5.8 | 4.6 | 6.6 ▲ | 4.0 | 5.6 |
| Crack | 2.5 | 1.8 | 2.8 | 1.7 | 2.6 | 1.9 | 2.0 | 2.9 | 3.2 | 1.7 | 2.1 | 2.5 | 3.5 ▲ | 1.6 ▼ | 2.5 |
| Inhalants | 7.9 | 7.7 | 8.8 | 5.8 ▼ | 8.4 | 7.1 | 7.5 | 9.1 | 7.0 | 5.4 ▼ | 8.8 | 8.9 | 8.5 | 7.3 | 7.8 |
| Amphetamines | 7.8 | 5.7 ▼ | 11.5 ▲ | 7.0 | 6.6 | 6.1 ▼ | 5.9 | 9.0 | 7.1 | 4.6 ▼ | 9.2 | 8.1 | 8.8 | 5.9 ▼ | 7.8 |
| Methamphetamines | 2.7 | 1.1 ▼ | 3.1 | 2.3 | 2.6 | 2.1 | 3.9 | 2.7 | 2.5 | 2.0 | 2.8 | 2.8 | 3.7 ▲ | 2.5 | 3.1 |
| Heroin | 2.2 | 1.4 | 2.8 | 1.1 | 1.9 | 2.4 | 2.6 | 2.4 | 2.3 | 1.8 | 2.5 | 2.9 | 2.6 | 1.4 ▼ | 1.8 |
| Steroids | 2.0 | 1.3 | 1.9 | 1.0 | 1.8 | 1.4 | 1.6 | 2.4 | 2.4 | 1.5 | 2.4 | 1.7 | 2.6 | 2.6 | 1.7 |
| Ecstasy | 5.7 | 6.0 | 7.0 ▲ | 2.7 ▼ | 6.0 | 4.8 | 5.2 | 6.5 | 6.0 | 6.0 | 6.1 | 5.2 | 6.6 | 3.8 ▼ | 5.7 |
| Tranquilizers | 5.2 | 4.6 | 7.7 ▲ | 3.8 | 5.0 | 3.8 ▼ | 5.2 | 6.3 | 4.4 | 2.6 ▼ | 6.1 | 4.4 | 6.1 | 4.2 | 5.5 |
| Hallucinogens | 6.5 | 4.5 ▼ | 7.2 | 3.3 ▼ | 6.8 | 6.5 | 5.2 | 6.2 | 7.4 | 5.3 | 6.3 | 7.1 | 8.5 ▲ | 5.3 | 6.7 |
| Prescription painkillers | 14.9 | 11.5 ▼ | 19.0 ▲ | 14.2 | 16.1 | 13.1 ▼ | 11.7 | 16.3 | 13.7 | 10.6 ▼ | 17.5 ▲ | 16.0 | 16.3 | 12.5 ▼ | 13.5 |
| Prescription drugs | 13.3 | 9.7 ▼ | 18.2 ▲ | 13.7 | 13.7 | 11.2 ▼ | 12.4 | 13.5 | 12.1 | 8.7 ▼ | 15.6 ▲ | 14.9 | 13.3 | 12.7 | 11.8 |
| Over the counter drugs | 9.5 | 7.1 ▼ | 11.3 ▲ | 8.1 | 8.7 | 8.3 | 7.5 | 11.2 ▲ | 9.0 | 7.9 | 9.9 | 10.3 | 11.0 ▲ | 7.9 | 9.0 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 11th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|-------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 20.7 | 16.4 ▼ | 22.3 | 19.5 | 22.9 | 17.6 ▼ | 19.2 | 24.0 ▲ | 17.2 ▼ | 10.2 ▼ | 22.7 | 21.8 | 23.3 ▲ | 21.2 | 23.6 ▲ |
| Smokeless tobacco | 8.8 | 2.8 ▼ | 10.3 ▲ | 9.8 | 5.8 ▼ | 6.2 ▼ | 10.7 | 7.8 | 7.4 | 1.8 ▼ | 12.3 ▲ | 10.9 ▲ | 10.5 ▲ | 9.6 | 12.0 ▲ |
| Cigars | 11.1 | 5.3 ▼ | 11.6 | 12.1 | 10.1 | 9.6 | 7.8 | 13.2 ▲ | 9.7 | 7.0 ▼ | 11.9 | 12.7 | 13.5 ▲ | 11.8 | 10.8 |
| Pipe | 8.3 | 10.1 | 10.6 ▲ | 6.3 | 7.2 | 6.0 ▼ | 5.5 | 8.4 | 8.8 | 7.2 | 8.7 | 7.0 | 9.4 | 8.4 | 8.9 |
| Alcohol | 31.7 | 30.5 | 35.0 ▲ | 29.6 | 30.5 | 30.8 | 23.1 ▼ | 32.5 | 25.0 ▼ | 33.3 | 28.7 ▼ | 33.1 | 35.1 ▲ | 33.0 | 31.1 |
| Binge drinking# | 20.8 | 19.6 | 21.3 | 20.4 | 20.9 | 18.8 | 12.7 ▼ | 22.9 | 17.2 ▼ | 21.1 | 19.8 | 20.9 | 22.5 | 24.5 ▲ | 20.4 |
| Marijuana | 18.6 | 18.3 | 20.9 ▲ | 14.8 ▼ | 18.4 | 19.2 | 9.8 ▼ | 21.5 ▲ | 17.8 | 25.8 ▲ | 19.0 | 15.2 ▼ | 19.2 | 16.7 | 15.3 ▼ |
| Cocaine | 1.8 | 1.1 | 2.2 | 1.0 | 2.2 | 1.6 | 1.3 | 2.0 | 1.5 | 2.1 | 2.0 | 1.6 | 2.2 | 1.2 | 2.4 |
| Crack | 1.2 | 0.8 | 1.4 | 0.6 | 1.0 | 0.8 | 0.7 | 1.3 | 1.8 | 1.2 | 1.2 | 0.8 | 1.5 | 0.7 | 1.7 |
| Inhalants | 1.8 | 1.3 | 2.1 | 1.1 | 1.8 | 1.5 | 1.6 | 2.3 | 1.7 | 1.5 | 2.2 | 1.8 | 1.8 | 1.6 | 2.3 |
| Amphetamines | 2.6 | 1.7 | 3.8 ▲ | 2.8 | 2.6 | 1.8 | 1.6 | 3.6 ▲ | 2.5 | 1.4 ▼ | 2.4 | 2.5 | 2.7 | 1.8 | 3.4 |
| Methamphetamines | 1.3 | 0.3 | 1.4 | 0.6 | 0.7 | 0.8 | 0.7 | 1.3 | 1.6 | 1.5 | 1.3 | 1.5 | 1.6 | 1.2 | 1.6 |
| Heroin | 1.1 | 0.4 | 1.1 | 0.7 | 1.0 | 0.8 | 1.0 | 1.6 | 1.4 | 1.2 | 1.0 | 1.5 | 1.3 | 1.0 | 1.3 |
| Steroids | 1.0 | 0.3 | 1.3 | 0.4 | 0.8 | 0.4 ▼ | 0.7 | 1.0 | 1.2 | 0.9 | 1.3 | 0.8 | 1.1 | 1.2 | 1.3 |
| Ecstasy | 1.8 | 1.5 | 1.9 | 1.0 | 1.9 | 2.1 | 2.0 | 2.3 | 2.5 | 1.7 | 1.4 | 1.7 | 1.9 | 1.2 | 1.6 |
| Tranquilizers | 2.0 | 1.1 | 2.9 ▲ | 1.6 | 1.8 | 1.3 | 1.3 | 2.4 | 1.8 | 1.2 | 1.8 | 2.0 | 2.2 | 1.9 | 2.6 |
| Hallucinogens | 2.2 | 1.5 | 1.9 | 1.3 | 1.7 | 2.7 | 1.6 | 2.2 | 2.8 | 2.2 | 2.1 | 1.1 ▼ | 3.2 ▲ | 1.8 | 3.0 |
| Prescription painkillers | 5.9 | 4.1 ▼ | 8.1 ▲ | 5.7 | 6.4 | 5.5 | 4.9 | 6.4 | 4.8 | 4.0 ▼ | 6.0 | 6.9 | 6.4 | 4.3 ▼ | 6.3 |
| Prescription drugs | 5.6 | 4.1 | 8.1 ▲ | 6.4 | 7.2 | 4.6 | 5.9 | 5.3 | 4.9 | 3.7 ▼ | 6.1 | 6.4 | 5.5 | 4.4 | 5.5 |
| Over the counter drugs | 3.3 | 2.5 | 4.0 | 2.8 | 2.4 | 3.2 | 2.3 | 4.3 | 2.7 | 2.3 | 3.1 | 3.4 | 4.1 | 2.7 | 3.9 |

NOTES: # Binge drinking during the past two weeks.
 ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
 ▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Lifetime Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 42.1 | 37.3 ▼ | 47.3 ▲ | 41.8 | 38.0 ▼ | 38.0 ▼ | 41.0 | 45.4 ▲ | 37.2 ▼ | 35.6 ▼ | 42.3 | 45.0 | 42.7 | 44.6 | 44.4 ▲ |
| Smokeless tobacco | 23.6 | 15.6 ▼ | 26.2 ▲ | 22.0 | 14.2 ▼ | 19.1 ▼ | 29.5 ▲ | 23.6 | 17.7 ▼ | 15.3 ▼ | 26.8 ▲ | 25.9 | 24.8 | 28.5 ▲ | 28.2 ▲ |
| Cigars | 35.9 | 28.4 ▼ | 39.0 ▲ | 37.8 | 29.1 ▼ | 35.0 | 40.0 ▲ | 34.4 | 32.2 ▼ | 26.3 ▼ | 37.1 | 37.4 | 36.9 | 40.0 ▲ | 38.4 ▲ |
| Pipe | 27.1 | 27.8 | 33.7 ▲ | 21.4 ▼ | 17.8 ▼ | 24.9 | 28.9 | 23.4 ▼ | 28.8 | 27.5 | 27.4 | 26.4 | 25.1 ▼ | 29.8 ▲ | 28.8 |
| Alcohol | 65.4 | 67.3 ▲ | 65.3 | 66.0 | 59.2 ▼ | 63.5 | 63.9 | 66.1 | 58.7 ▼ | 69.2 ▲ | 63.6 | 68.0 | 65.5 | 68.1 ▲ | 68.0 ▲ |
| Marijuana | 38.6 | 42.1 | 37.9 | 37.8 | 33.4 ▼ | 37.6 | 37.1 | 41.2 | 34.4 ▼ | 51.5 ▲ | 37.1 | 38.3 | 39.2 | 36.2 | 39.3 |
| Cocaine | 5.9 | 5.6 | 6.4 | 4.8 | 3.6 ▼ | 4.8 | 6.9 | 5.0 | 5.8 | 6.2 | 5.6 | 7.0 | 7.0 ▲ | 5.3 | 6.9 |
| Crack | 2.6 | 2.1 | 2.8 | 0.3 | 2.4 | 2.5 | 3.0 | 2.9 | 2.7 | 3.7 | 2.8 | 2.9 | 2.8 | 2.4 | 2.4 |
| Inhalants | 7.8 | 6.4 | 8.3 | 6.9 | 6.4 | 5.8 ▼ | 8.7 | 8.3 | 7.6 | 8.1 | 8.0 | 8.0 | 8.5 | 8.9 | 7.8 |
| Amphetamines | 8.3 | 6.4 ▼ | 10.1 ▲ | 8.7 | 7.0 | 6.5 ▼ | 9.4 | 8.0 | 8.9 | 9.3 | 7.4 | 9.8 | 8.7 | 7.4 | 8.9 |
| Methamphetamines | 2.8 | 2.2 | 3.3 | 2.0 | 1.7 | 1.9 ▼ | 4.8 ▲ | 2.9 | 2.0 | 3.8 | 2.5 | 3.4 | 3.0 | 2.5 | 3.1 |
| Heroin | 2.4 | 2.0 | 3.3 ▲ | 0.7 | 1.0 ▼ | 2.2 | 2.9 | 2.6 | 2.7 | 4.1 ▲ | 2.5 | 2.7 | 2.5 | 2.0 | 2.1 |
| Steroids | 2.1 | 2.8 | 2.4 | 1.2 | 0.7 | 2.0 | 3.2 ▲ | 2.6 | 2.0 | 1.9 | 2.0 | 2.2 | 2.3 | 1.6 | 2.0 |
| Ecstasy | 6.7 | 7.8 | 7.4 | 4.3 ▼ | 4.6 ▼ | 5.9 | 6.7 | 7.1 | 7.1 | 9.6 ▲ | 6.6 | 7.9 | 7.1 | 4.8 ▼ | 6.1 |
| Tranquilizers | 5.4 | 4.8 | 7.2 ▲ | 4.1 | 3.1 ▼ | 4.6 | 6.9 | 5.4 | 5.2 | 6.4 | 4.9 | 6.4 | 5.6 | 4.7 | 5.5 |
| Hallucinogens | 7.2 | 4.7 ▼ | 7.4 | 5.8 | 5.7 | 7.6 | 7.2 | 5.8 | 7.6 | 9.5 ▲ | 6.1 | 8.4 | 8.1 | 8.1 | 6.8 |
| Prescription painkillers | 16.0 | 13.6 ▼ | 15.9 | 16.6 | 13.8 | 14.5 | 17.9 | 15.4 | 14.2 | 16.3 | 16.9 | 18.4 ▲ | 16.9 | 15.4 | 17.4 |
| Prescription drugs | 14.6 | 12.9 | 16.4 ▲ | 14.0 | 14.6 | 12.4 ▼ | 16.5 | 12.9 | 14.1 | 13.8 | 14.4 | 16.2 | 13.6 | 14.7 | 17.0 ▲ |
| Over the counter drugs | 9.7 | 9.6 | 10.2 | 7.9 | 8.8 | 8.2 | 10.3 | 9.5 | 10.0 | 9.9 | 9.7 | 11.4 | 9.6 | 9.5 | 10.7 |

NOTES: ▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).
▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).

Percentage Reporting Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 12th Grade Students: 2011

| DRUG | Indiana | DSA1 | DSA2 | DSA3 | DSA4 | DSA5 | DSA6 | DSA7 | DSA8 | DSA9 | DSA10 | DSA11 | DSA12 | DSA13 | DSA14 |
|--------------------------|---------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|--------|
| Cigarettes | 24.6 | 19.5 ▼ | 29.1 ▲ | 23.8 | 19.6 ▼ | 22.9 | 24.0 | 25.4 | 20.7 ▼ | 19.5 ▼ | 24.2 | 26.8 | 25.3 | 25.8 | 27.9 ▲ |
| Smokeless tobacco | 10.7 | 7.6 ▼ | 12.3 ▲ | 8.4 | 5.7 ▼ | 8.9 ▼ | 16.3 ▲ | 9.8 | 7.0 ▼ | 6.4 ▼ | 11.8 | 11.3 | 10.4 | 12.8 ▲ | 14.8 ▲ |
| Cigars | 16.7 | 13.0 ▼ | 18.9 ▲ | 16.0 | 10.2 ▼ | 14.9 | 23.0 ▲ | 17.1 | 13.3 ▼ | 11.9 ▼ | 17.8 | 19.5 ▲ | 15.8 | 19.4 ▲ | 18.3 |
| Pipe | 12.5 | 14.4 | 15.3 ▲ | 8.4 ▼ | 7.0 ▼ | 10.7 ▼ | 14.6 | 9.5 ▼ | 13.4 | 14.4 | 14.1 | 11.2 | 10.5 ▼ | 14.7 ▲ | 13.5 |
| Alcohol | 37.8 | 40.6 | 37.1 | 33.1 ▼ | 31.7 ▼ | 38.8 | 35.0 | 36.3 | 30.8 ▼ | 41.2 | 33.8 ▼ | 40.5 | 38.0 | 41.8 ▲ | 43.0 ▲ |
| Binge drinking# | 26.4 | 29.7 ▲ | 25.9 | 22.9 ▼ | 18.1 ▼ | 25.6 | 29.2 | 26.5 | 20.3 ▼ | 28.9 | 22.5 ▼ | 25.9 | 27.0 | 30.4 ▲ | 30.7 ▲ |
| Marijuana | 19.8 | 22.9 ▲ | 19.0 | 16.3 ▼ | 17.8 ▼ | 21.1 | 21.0 | 22.4 ▲ | 16.9 ▼ | 29.6 ▲ | 16.5 ▼ | 18.9 | 19.7 | 17.1 ▼ | 20.5 |
| Cocaine | 2.0 | 2.2 | 2.4 | 1.5 | 1.0 | 2.1 | 3.0 | 2.3 | 1.8 | 3.0 | 1.4 | 2.2 | 2.0 | 1.4 | 2.0 |
| Crack | 1.2 | 1.3 | 1.4 | 0.5 | 0.7 | 1.5 | 2.0 ▲ | 1.2 | 0.9 | 1.9 | 0.7 | 1.3 | 0.9 | 0.9 | 1.1 |
| Inhalants | 1.7 | 2.0 | 2.1 | 0.7 | 1.3 | 1.6 | 2.1 | 2.2 | 1.2 | 2.6 | 1.2 | 1.4 | 1.5 | 1.8 | 1.9 |
| Amphetamines | 2.7 | 2.7 | 3.1 | 3.5 | 2.2 | 2.5 | 3.3 | 2.7 | 2.5 | 3.6 | 2.3 | 3.0 | 2.7 | 2.3 | 2.7 |
| Methamphetamines | 1.3 | 1.2 | 1.7 | 0.5 | 0.7 | 1.3 | 2.5 ▲ | 1.4 | 1.1 | 2.1 | 0.9 | 1.1 | 1.1 | 1.1 | 1.4 |
| Heroin | 1.2 | 1.5 | 1.7 | 0.2 | 0.1 | 1.4 | 1.8 | 1.4 | 0.9 | 2.1 ▲ | 0.9 | 1.1 | 1.2 | 1.1 | 1.0 |
| Steroids | 1.1 | 1.6 | 1.4 | 0.3 | 0.3 | 0.9 | 1.5 | 1.0 | 0.9 | 1.5 | 1.2 | 1.1 | 1.0 | 1.0 | 1.1 |
| Ecstasy | 2.1 | 2.9 | 2.6 | 0.2 | 1.1 | 1.9 | 2.5 | 1.6 | 2.5 | 4.2 ▲ | 1.7 | 2.8 | 1.8 | 1.8 | 1.7 |
| Tranquilizers | 2.0 | 2.6 | 2.4 | 1.8 | 0.6 | 1.6 | 3.1 ▲ | 1.9 | 2.0 | 3.0 | 1.8 | 2.0 | 1.8 | 1.4 | 2.1 |
| Hallucinogens | 2.4 | 1.9 | 2.7 | 1.6 | 1.1 ▼ | 2.6 | 2.9 | 1.8 | 2.6 | 4.7 ▲ | 2.3 | 2.1 | 3.2 ▲ | 2.6 | 1.6 ▼ |
| Prescription painkillers | 6.6 | 5.8 | 6.7 | 7.6 | 5.1 | 5.6 | 8.9 ▲ | 6.3 | 5.4 | 6.7 | 7.3 | 8.4 ▲ | 6.5 | 5.7 | 6.9 |
| Prescription drugs | 5.9 | 6.3 | 6.3 | 5.9 | 5.8 | 4.4 ▼ | 6.9 | 5.1 | 6.4 | 6.8 | 6.1 | 7.1 | 5.3 | 5.1 | 6.6 |
| Over the counter drugs | 3.2 | 4.2 | 4.0 | 2.8 | 2.8 | 2.7 | 3.8 | 3.3 | 3.2 | 4.2 | 2.3 | 3.9 | 2.5 | 3.2 | 2.8 |

NOTES: # Binge drinking during the past two weeks.

▲ indicates that local prevalence rates are significantly higher than state rates ($p < .05$).

▼ indicates that local prevalence rates are significantly lower than state rates ($p < .05$).




VII. TRENDS AND PREVALENCE GRAPHS

Trends Graphs 134


Prevalence Graphs 148

Trends lifetime Alcohol Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Monthly Alcohol Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Lifetime Cigarettes Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Monthly Cigarettes Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Lifetime Smokeless Tobacco Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Monthly Smokeless Tobacco Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Lifetime Marijuana Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Monthly Marijuana Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Lifetime Inhalants Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Monthly Inhalants Use Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Binge Drinking Among Indiana Students: 1993 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Trends of Binge Drinking by Indiana and U.S.A. 12th Graders: 1975 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010
Johnston, et al., National Institute on Drug Abuse, 2009

Trends of Monthly Cigarette Use by Indiana and U.S.A. 12th Graders: 1975 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010
Johnston, et al., National Institute on Drug Abuse, 2009

Trends of Monthly Marijuana Use by Indiana and U.S.A. 12th Graders: 1975 - 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010
Johnston, et al., National Institute on Drug Abuse, 2009

Monthly Use of Selected Drugs and Binge Drinking by Indiana and U.S.A. 12th Graders




Source: Indiana Prevention Resource Center at Indiana University, 2010
Johnston, et al., National Institute on Drug Abuse, 2009

Alcohol Use Among Indiana Students 6th - 12th Graders, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Cigarette Use Among Indiana Students 6th - 12th Graders, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Smokeless Tobacco Use Among Indiana Students 6th - 12th Graders, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Marijuana Use Among Indiana Students 6th - 12th Graders, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Inhalants Use Among Indiana Students 6th - 12th Graders, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Binge Drinking Among Indiana Students by Grade and by Gender, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Monthly Alcohol Use Among Indiana Students by Grade and by Gender, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Monthly Cigarette Use Among Indiana Students by Grade and by Gender, 2011




Source: Indiana Prevention Resource Center at Indiana University, 2010

Monthly Smokeless Tobacco Use Among Indiana Students by Grade and by Gender, 2011



Source: Indiana Prevention Resource Center at Indiana University, 2010

Monthly Marijuana Use Among Indiana Students by Grade and by Gender, 2011



Source: Indiana Prevention Resource Center at Indiana University, 2010



VIII FREQUENCY TABLES

| | |
|----------------------|-----|
| General | 160 |
| Male | 194 |
| Female | 228 |

**Personal and Family Information
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------|----------------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Hispanic | No Answer | 7.2 | 6.0 | 6.0 | 6.2 | 6.6 | 5.8 | 5.3 |
| | Hispanic | 8.1 | 8.8 | 8.6 | 8.4 | 7.4 | 7.5 | 6.7 |
| | Non-Hispanic | 84.7 | 85.2 | 85.4 | 85.3 | 86.0 | 86.6 | 88.0 |
| Race | No Answer | 3.3 | 3.7 | 3.8 | 4.1 | 3.4 | 2.9 | 3.0 |
| | White | 78.8 | 80.2 | 78.7 | 79.4 | 79.3 | 81.2 | 83.1 |
| | Black/African American | 6.6 | 5.1 | 6.2 | 5.6 | 7.1 | 6.1 | 5.8 |
| | Asian | 1.5 | 1.5 | 1.6 | 1.7 | 1.7 | 1.9 | 1.6 |
| | Native Hawaiian/Pacific Islander | .1 | .2 | .3 | .4 | .4 | .4 | .4 |
| | American Indian/Alaskan Native | .5 | .6 | .6 | .6 | .7 | .6 | .5 |
| | More than one race | 5.9 | 5.8 | 6.4 | 5.9 | 5.4 | 4.9 | 4.0 |
| | Race not known or other | 3.3 | 3.0 | 2.3 | 2.4 | 2.0 | 1.9 | 1.5 |
| Age | No Answer | 1.0 | 1.2 | 1.1 | 1.2 | 1.5 | 1.3 | 1.1 |
| | 10 years or younger | .3 | .0 | .0 | .0 | .0 | .0 | .0 |
| | 11 years | 22.7 | .2 | .0 | .0 | .0 | .0 | .0 |
| | 12 years | 65.7 | 21.6 | .1 | .0 | .0 | .0 | .0 |
| | 13 years | 9.8 | 68.0 | 21.9 | .1 | .0 | .0 | .0 |
| | 14 years | .5 | 8.6 | 66.9 | 21.0 | .1 | .0 | .0 |
| | 15 years | .0 | .4 | 9.4 | 67.7 | 18.7 | .3 | .0 |
| | 16 years | .0 | .0 | .5 | 9.5 | 70.6 | 20.5 | .3 |
| | 17 years | .0 | .0 | .0 | .5 | 8.7 | 70.7 | 20.8 |
| | 18 years or older | .0 | .0 | .0 | .0 | .4 | 7.3 | 77.8 |
| Grades | No Answer | 4.8 | 4.2 | 4.1 | 3.8 | 2.9 | 2.3 | 1.8 |
| | Mostly A's | 37.5 | 36.8 | 35.4 | 33.4 | 31.7 | 31.5 | 36.0 |
| | Mostly B's | 36.1 | 35.4 | 35.4 | 36.1 | 38.0 | 40.5 | 39.0 |
| | Mostly C's | 16.6 | 17.9 | 19.9 | 21.1 | 21.9 | 21.5 | 20.0 |
| | Mostly D's | 3.5 | 3.9 | 3.8 | 3.9 | 4.1 | 3.3 | 2.5 |
| | Mostly F's | 1.5 | 1.9 | 1.4 | 1.6 | 1.3 | 1.0 | .8 |
| Live with | No Answer | 1.8 | 1.6 | 1.6 | 1.6 | 1.1 | 1.1 | .9 |
| | Both parents | 59.5 | 59.6 | 56.2 | 56.2 | 55.5 | 56.0 | 56.7 |
| | Mother only | 14.1 | 13.5 | 15.5 | 15.7 | 16.4 | 16.3 | 16.3 |
| | Father only | 2.5 | 3.0 | 3.3 | 3.7 | 4.2 | 4.5 | 4.4 |
| | Mother & stepfather | 12.8 | 13.2 | 14.4 | 13.9 | 13.5 | 12.8 | 11.1 |
| | Father & stepmother | 2.9 | 3.3 | 3.6 | 3.7 | 3.9 | 3.5 | 3.0 |
| | Other | 6.4 | 5.8 | 5.4 | 5.1 | 5.2 | 5.8 | 7.7 |
| Employed - You | No Answer | 18.5 | 15.4 | 15.0 | 12.6 | 10.1 | 7.0 | 5.3 |
| | Yes, full time | .9 | 1.0 | 1.2 | 1.5 | 1.8 | 2.2 | 3.4 |
| | Yes, part time | 6.5 | 8.3 | 10.8 | 13.2 | 20.5 | 36.3 | 47.3 |
| | No | 74.1 | 75.2 | 73.1 | 72.7 | 67.6 | 54.5 | 44.0 |
| Employed - Mom | No Answer | 6.8 | 5.7 | 5.1 | 5.3 | 4.6 | 4.2 | 3.9 |
| | Yes, full time | 52.4 | 55.3 | 58.2 | 59.7 | 62.5 | 64.3 | 64.5 |
| | Yes, part time | 21.3 | 19.7 | 17.4 | 15.8 | 14.6 | 13.7 | 12.9 |
| | No | 19.5 | 19.3 | 19.3 | 19.2 | 18.3 | 17.8 | 18.7 |
| Employed - Dad | No Answer | 8.5 | 7.4 | 7.7 | 8.2 | 7.9 | 7.6 | 7.2 |
| | Yes, full time | 70.5 | 73.8 | 75.2 | 76.0 | 77.7 | 78.4 | 78.3 |
| | Yes, part time | 12.4 | 9.9 | 8.1 | 6.6 | 5.1 | 4.4 | 4.0 |
| | No | 8.6 | 8.9 | 9.0 | 9.2 | 9.3 | 9.6 | 10.6 |

Lifetime Use of Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | | Grade | | | | | | |
|-------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | | .3 | .3 | .4 | .3 | .4 | .5 | .5 |
| | Never | | 92.4 | 86.5 | 79.0 | 72.5 | 66.8 | 61.4 | 57.4 |
| | 1-5 times | | 5.5 | 8.6 | 11.6 | 12.6 | 13.9 | 14.6 | 14.8 |
| | 6-19 times | | .8 | 2.0 | 3.1 | 4.0 | 4.7 | 5.3 | 5.6 |
| | 20-40 times | | .3 | .8 | 1.8 | 2.3 | 3.0 | 2.8 | 3.6 |
| | 40+ times | | .6 | 1.8 | 4.1 | 8.4 | 11.3 | 15.3 | 18.1 |
| Smokeless Tobacco | No Answer | | .2 | .2 | .3 | .5 | .5 | .5 | .6 |
| | Never | | 97.3 | 94.6 | 91.1 | 86.4 | 82.3 | 79.4 | 75.8 |
| | 1-5 times | | 1.9 | 3.7 | 5.4 | 7.3 | 8.8 | 9.7 | 10.8 |
| | 6-19 times | | .3 | .6 | 1.1 | 2.0 | 2.6 | 2.7 | 3.3 |
| | 20-40 times | | .1 | .3 | .6 | 1.0 | 1.4 | 1.5 | 2.2 |
| | 40+ times | | .2 | .5 | 1.4 | 2.8 | 4.4 | 6.0 | 7.3 |
| Cigars | No Answer | | .3 | .3 | .5 | .5 | .7 | .7 | .8 |
| | Never | | 97.8 | 95.4 | 91.4 | 84.6 | 78.4 | 72.2 | 63.3 |
| | 1-5 times | | 1.4 | 3.2 | 5.4 | 8.8 | 11.3 | 13.1 | 15.6 |
| | 6-19 times | | .2 | .5 | 1.4 | 3.0 | 4.5 | 6.1 | 8.4 |
| | 20-40 times | | .1 | .2 | .6 | 1.5 | 2.2 | 3.1 | 4.9 |
| | 40+ times | | .2 | .3 | .8 | 1.6 | 2.9 | 4.7 | 7.0 |
| Pipe | No Answer | | .4 | .5 | .6 | .8 | .9 | .9 | 1.1 |
| | Never | | 98.3 | 96.3 | 93.4 | 88.8 | 84.1 | 78.9 | 71.8 |
| | 1-5 times | | .8 | 2.1 | 3.5 | 5.5 | 7.7 | 9.7 | 12.2 |
| | 6-19 times | | .2 | .5 | 1.1 | 1.9 | 3.0 | 4.3 | 6.4 |
| | 20-40 times | | .1 | .3 | .6 | 1.2 | 1.7 | 2.2 | 3.4 |
| | 40+ times | | .2 | .4 | .9 | 1.8 | 2.6 | 4.0 | 5.2 |
| Alcohol | No Answer | | .6 | .6 | .7 | .8 | .9 | .9 | 1.2 |
| | Never | | 82.0 | 73.3 | 61.6 | 51.5 | 43.6 | 38.5 | 33.5 |
| | 1-5 times | | 14.6 | 18.6 | 23.1 | 23.9 | 24.1 | 22.1 | 19.7 |
| | 6-19 times | | 1.8 | 4.6 | 8.2 | 12.3 | 14.1 | 15.5 | 15.0 |
| | 20-40 times | | .5 | 1.5 | 3.1 | 5.1 | 7.1 | 8.7 | 10.4 |
| | 40+ times | | .5 | 1.4 | 3.3 | 6.5 | 10.3 | 14.3 | 20.2 |
| Marijuana | No Answer | | .3 | .4 | .5 | .7 | .8 | .9 | 1.1 |
| | Never | | 96.6 | 92.3 | 84.9 | 76.3 | 68.8 | 63.6 | 60.3 |
| | 1-5 times | | 1.9 | 4.0 | 7.0 | 9.3 | 10.9 | 11.8 | 12.5 |
| | 6-19 times | | .5 | 1.2 | 2.7 | 4.0 | 5.4 | 6.0 | 6.6 |
| | 20-40 times | | .2 | .7 | 1.5 | 2.6 | 3.4 | 4.1 | 4.4 |
| | 40+ times | | .5 | 1.3 | 3.4 | 7.1 | 10.7 | 13.5 | 15.1 |
| Cocaine | No Answer | | .3 | .3 | .4 | .5 | .7 | .7 | .9 |
| | Never | | 99.1 | 98.6 | 97.6 | 96.6 | 95.6 | 94.2 | 93.2 |
| | 1-5 times | | .3 | .6 | 1.2 | 1.7 | 2.4 | 3.2 | 3.5 |
| | 6-19 times | | .1 | .2 | .3 | .5 | .6 | .8 | 1.0 |
| | 20-40 times | | .1 | .1 | .2 | .3 | .3 | .3 | .5 |
| | 40+ times | | .2 | .2 | .3 | .5 | .5 | .8 | .8 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | | Grade | | | | | | |
|------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Crack | No Answer | | .3 | .3 | .4 | .5 | .6 | .5 | .6 |
| | Never | | 99.1 | 98.6 | 97.9 | 97.6 | 97.4 | 97.0 | 96.8 |
| | 1-5 times | | .3 | .6 | 1.0 | 1.0 | 1.1 | 1.2 | 1.4 |
| | 6-19 times | | .1 | .1 | .2 | .3 | .3 | .4 | .4 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .2 | .3 | .2 |
| | 40+ times | | .2 | .2 | .3 | .3 | .4 | .5 | .6 |
| Inhalants | No Answer | | 1.2 | .5 | .5 | .6 | .6 | .6 | .7 |
| | Never | | 97.0 | 96.3 | 94.1 | 92.5 | 92.0 | 91.5 | 91.5 |
| | 1-5 times | | 1.1 | 2.2 | 3.6 | 4.4 | 4.6 | 4.5 | 4.4 |
| | 6-19 times | | .4 | .6 | 1.0 | 1.3 | 1.6 | 1.8 | 1.8 |
| | 20-40 times | | .1 | .2 | .3 | .5 | .6 | .8 | .7 |
| | 40+ times | | .2 | .3 | .5 | .6 | .7 | .8 | 1.0 |
| Amphetamines | No Answer | | 1.6 | .7 | .6 | .6 | .7 | .7 | .7 |
| | Never | | 98.0 | 98.2 | 97.1 | 95.1 | 93.3 | 91.6 | 90.9 |
| | 1-5 times | | .2 | .6 | 1.3 | 2.0 | 2.6 | 3.3 | 3.1 |
| | 6-19 times | | .1 | .2 | .5 | 1.0 | 1.4 | 1.8 | 2.0 |
| | 20-40 times | | .0 | .1 | .2 | .6 | .8 | 1.1 | 1.2 |
| | 40+ times | | .1 | .2 | .3 | .7 | 1.1 | 1.6 | 1.9 |
| Methamphetamines | No Answer | | 1.0 | .5 | .5 | .6 | .6 | .6 | .6 |
| | Never | | 98.5 | 98.5 | 98.0 | 97.6 | 97.2 | 96.7 | 96.5 |
| | 1-5 times | | .2 | .6 | .8 | 1.0 | 1.1 | 1.3 | 1.3 |
| | 6-19 times | | .1 | .1 | .2 | .4 | .4 | .4 | .5 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .2 | .4 | .3 |
| | 40+ times | | .1 | .2 | .3 | .3 | .4 | .7 | .7 |
| Heroin | No Answer | | .9 | .5 | .5 | .6 | .6 | .6 | .8 |
| | Never | | 98.7 | 98.7 | 98.4 | 97.8 | 97.5 | 97.1 | 96.8 |
| | 1-5 times | | .2 | .4 | .6 | .8 | .9 | 1.0 | 1.0 |
| | 6-19 times | | .0 | .2 | .2 | .3 | .3 | .4 | .4 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .2 | .2 | .3 |
| | 40+ times | | .1 | .1 | .2 | .3 | .4 | .6 | .7 |
| Steroids | No Answer | | .7 | .5 | .5 | .6 | .6 | .6 | .8 |
| | Never | | 98.8 | 98.5 | 98.2 | 97.9 | 97.7 | 97.5 | 97.1 |
| | 1-5 times | | .3 | .6 | .7 | .8 | .8 | .9 | .9 |
| | 6-19 times | | .1 | .2 | .2 | .3 | .2 | .4 | .4 |
| | 20-40 times | | .0 | .0 | .1 | .2 | .2 | .2 | .3 |
| | 40+ times | | .1 | .2 | .3 | .4 | .4 | .6 | .6 |
| Ecstasy | No Answer | | 1.7 | .8 | .5 | .7 | .7 | .7 | .8 |
| | Never | | 97.9 | 98.2 | 97.3 | 95.9 | 94.9 | 93.6 | 92.6 |
| | 1-5 times | | .2 | .7 | 1.4 | 2.1 | 2.8 | 3.7 | 4.2 |
| | 6-19 times | | .1 | .2 | .3 | .6 | .7 | 1.0 | 1.2 |
| | 20-40 times | | .0 | .1 | .2 | .3 | .5 | .4 | .5 |
| | 40+ times | | .1 | .1 | .3 | .4 | .5 | .6 | .7 |

Lifetime Use of Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|--------------------------|-------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Tranquilizers | No Answer | 1.5 | .8 | .6 | .8 | .7 | .8 | .9 |
| | Never | 98.0 | 98.3 | 97.5 | 95.9 | 95.0 | 94.0 | 93.6 |
| | 1-5 times | .3 | .7 | 1.1 | 1.7 | 2.1 | 2.5 | 2.4 |
| | 6-19 times | .1 | .1 | .3 | .8 | .9 | 1.2 | 1.3 |
| | 20-40 times | .0 | .1 | .1 | .4 | .5 | .5 | .6 |
| | 40+ times | .1 | .1 | .3 | .5 | .7 | .9 | 1.1 |
| Hallucinogens | No Answer | 1.8 | .8 | .7 | .9 | .9 | .9 | 1.1 |
| | Never | 97.7 | 98.3 | 97.0 | 95.6 | 94.1 | 92.6 | 91.7 |
| | 1-5 times | .2 | .5 | 1.4 | 2.1 | 3.1 | 4.1 | 4.2 |
| | 6-19 times | .1 | .1 | .4 | .6 | .9 | 1.2 | 1.5 |
| | 20-40 times | .0 | .1 | .2 | .4 | .4 | .4 | .5 |
| | 40+ times | .1 | .2 | .3 | .4 | .6 | .8 | 1.0 |
| Prescription painkillers | No Answer | .9 | .6 | .7 | .7 | .7 | .8 | .9 |
| | Never | 98.0 | 96.8 | 93.8 | 89.9 | 86.7 | 84.3 | 83.1 |
| | 1-5 times | .6 | 1.5 | 3.1 | 4.7 | 5.7 | 6.2 | 6.1 |
| | 6-19 times | .1 | .6 | 1.2 | 2.2 | 3.1 | 3.6 | 3.8 |
| | 20-40 times | .1 | .3 | .6 | 1.0 | 1.6 | 1.9 | 2.3 |
| | 40+ times | .2 | .3 | .7 | 1.5 | 2.2 | 3.2 | 3.8 |
| Prescription drugs | No Answer | 1.0 | .6 | .7 | .8 | .9 | .9 | 1.0 |
| | Never | 98.0 | 97.0 | 94.5 | 91.0 | 88.0 | 85.8 | 84.4 |
| | 1-5 times | .6 | 1.5 | 2.6 | 4.0 | 5.0 | 5.2 | 5.7 |
| | 6-19 times | .1 | .4 | .9 | 1.7 | 2.4 | 3.1 | 3.3 |
| | 20-40 times | .1 | .2 | .6 | .9 | 1.4 | 1.9 | 1.8 |
| | 40+ times | .3 | .3 | .7 | 1.6 | 2.3 | 3.1 | 3.8 |
| Over the counter drugs | No Answer | .8 | .8 | .7 | .8 | .8 | .8 | .9 |
| | Never | 98.1 | 96.5 | 94.3 | 92.0 | 90.4 | 89.7 | 89.4 |
| | 1-5 times | .7 | 1.9 | 3.1 | 3.9 | 4.7 | 4.7 | 4.9 |
| | 6-19 times | .2 | .4 | .8 | 1.5 | 1.9 | 2.2 | 2.2 |
| | 20-40 times | .1 | .2 | .5 | .6 | .8 | 1.0 | 1.0 |
| | 40+ times | .2 | .3 | .6 | 1.1 | 1.4 | 1.6 | 1.8 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | | Grade | | | | | | |
|-------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | | 1.7 | 1.5 | 1.8 | 2.1 | 1.9 | 1.8 | 1.9 |
| | Never | | 95.7 | 92.5 | 88.2 | 83.6 | 80.9 | 77.4 | 73.5 |
| | 1-5 times | | 1.9 | 3.8 | 5.7 | 6.5 | 6.9 | 7.0 | 7.8 |
| | 6-19 times | | .3 | 1.0 | 1.7 | 2.5 | 2.9 | 3.1 | 3.6 |
| | 20-40 times | | .1 | .5 | 1.0 | 1.6 | 1.9 | 2.5 | 2.5 |
| | 40+ times | | .2 | .7 | 1.7 | 3.8 | 5.5 | 8.1 | 10.6 |
| Smokeless Tobacco | No Answer | | 1.9 | 1.7 | 2.2 | 2.3 | 2.2 | 2.2 | 2.1 |
| | Never | | 97.1 | 96.1 | 93.9 | 91.5 | 89.8 | 89.0 | 87.2 |
| | 1-5 times | | .7 | 1.5 | 2.3 | 3.3 | 3.6 | 3.6 | 4.0 |
| | 6-19 times | | .2 | .3 | .7 | 1.0 | 1.3 | 1.4 | 1.8 |
| | 20-40 times | | .1 | .2 | .3 | .7 | .8 | 1.1 | 1.3 |
| | 40+ times | | .1 | .2 | .6 | 1.2 | 2.1 | 2.8 | 3.6 |
| Cigars | No Answer | | 2.2 | 1.7 | 2.4 | 2.6 | 2.3 | 2.3 | 2.3 |
| | Never | | 97.2 | 96.7 | 94.4 | 91.4 | 89.0 | 86.6 | 81.0 |
| | 1-5 times | | .5 | 1.2 | 2.4 | 4.2 | 6.0 | 7.5 | 11.4 |
| | 6-19 times | | .1 | .2 | .5 | 1.0 | 1.5 | 2.1 | 3.0 |
| | 20-40 times | | .0 | .1 | .2 | .4 | .5 | .6 | 1.0 |
| | 40+ times | | .1 | .1 | .2 | .3 | .7 | .9 | 1.3 |
| Pipe | No Answer | | 2.2 | 1.8 | 2.6 | 2.8 | 2.5 | 2.6 | 2.6 |
| | Never | | 97.2 | 96.7 | 94.4 | 92.5 | 90.9 | 89.1 | 84.9 |
| | 1-5 times | | .4 | 1.0 | 2.0 | 3.0 | 3.9 | 5.2 | 8.5 |
| | 6-19 times | | .1 | .2 | .6 | .9 | 1.2 | 1.7 | 2.1 |
| | 20-40 times | | .0 | .1 | .2 | .4 | .7 | .6 | .7 |
| | 40+ times | | .0 | .1 | .2 | .5 | .8 | .8 | 1.2 |
| Alcohol | No Answer | | 2.4 | 1.9 | 2.5 | 2.6 | 2.4 | 2.4 | 2.2 |
| | Never | | 92.2 | 87.9 | 80.8 | 73.6 | 69.2 | 65.9 | 60.0 |
| | 1-5 times | | 4.6 | 8.3 | 12.8 | 17.6 | 20.4 | 21.8 | 25.0 |
| | 6-19 times | | .5 | 1.3 | 2.6 | 4.1 | 5.1 | 6.7 | 8.5 |
| | 20-40 times | | .1 | .3 | .7 | 1.1 | 1.5 | 1.6 | 2.2 |
| | 40+ times | | .1 | .3 | .6 | 1.1 | 1.4 | 1.6 | 2.1 |
| Marijuana | No Answer | | 2.3 | 1.8 | 2.5 | 2.6 | 2.5 | 2.4 | 2.5 |
| | Never | | 96.1 | 94.1 | 89.2 | 84.4 | 81.1 | 79.0 | 77.7 |
| | 1-5 times | | 1.0 | 2.5 | 4.8 | 6.5 | 7.8 | 8.7 | 8.6 |
| | 6-19 times | | .3 | .8 | 1.5 | 2.6 | 3.3 | 3.6 | 3.7 |
| | 20-40 times | | .1 | .4 | .9 | 1.5 | 2.0 | 2.2 | 2.4 |
| | 40+ times | | .2 | .4 | 1.1 | 2.3 | 3.4 | 4.0 | 5.1 |
| Cocaine | No Answer | | 2.5 | 1.9 | 2.6 | 2.8 | 2.5 | 2.5 | 2.6 |
| | Never | | 97.3 | 97.5 | 96.4 | 95.8 | 95.9 | 95.6 | 95.4 |
| | 1-5 times | | .2 | .4 | .6 | .8 | .9 | 1.1 | 1.2 |
| | 6-19 times | | .1 | .1 | .2 | .3 | .3 | .3 | .4 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .1 | .2 | .2 |
| | 40+ times | | .0 | .1 | .1 | .1 | .2 | .2 | .2 |

Monthly Use of Alcohol, Tobacco, and Other Drugs by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------|-------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Crack | No Answer | 2.5 | 1.9 | 2.6 | 2.9 | 2.6 | 2.5 | 2.5 |
| | Never | 97.3 | 97.4 | 96.5 | 96.2 | 96.5 | 96.3 | 96.3 |
| | 1-5 times | .1 | .4 | .5 | .5 | .5 | .6 | .6 |
| | 6-19 times | .0 | .1 | .2 | .3 | .2 | .2 | .2 |
| | 20-40 times | .0 | .1 | .1 | .1 | .1 | .2 | .1 |
| | 40+ times | .1 | .1 | .1 | .1 | .1 | .2 | .1 |
| Inhalants | No Answer | 3.1 | 2.0 | 2.7 | 2.9 | 2.6 | 2.6 | 2.6 |
| | Never | 96.0 | 96.6 | 95.0 | 94.9 | 95.3 | 95.6 | 95.7 |
| | 1-5 times | .6 | 1.1 | 1.6 | 1.6 | 1.5 | 1.3 | 1.0 |
| | 6-19 times | .2 | .2 | .4 | .4 | .4 | .4 | .4 |
| | 20-40 times | .1 | .1 | .1 | .1 | .1 | .1 | .2 |
| | 40+ times | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| Amphetamines | No Answer | 3.5 | 2.3 | 2.7 | 2.8 | 2.7 | 2.6 | 2.6 |
| | Never | 96.3 | 97.2 | 96.1 | 95.3 | 94.9 | 94.8 | 94.7 |
| | 1-5 times | .1 | .4 | .8 | 1.1 | 1.5 | 1.6 | 1.6 |
| | 6-19 times | .1 | .1 | .2 | .5 | .5 | .6 | .6 |
| | 20-40 times | .0 | .0 | .1 | .2 | .2 | .3 | .2 |
| | 40+ times | .0 | .0 | .1 | .1 | .1 | .2 | .2 |
| Methamphetamines | No Answer | 3.1 | 2.1 | 2.8 | 2.9 | 2.7 | 2.6 | 2.6 |
| | Never | 96.6 | 97.4 | 96.3 | 96.3 | 96.4 | 96.1 | 96.1 |
| | 1-5 times | .2 | .3 | .6 | .4 | .5 | .7 | .6 |
| | 6-19 times | .1 | .1 | .2 | .3 | .3 | .3 | .3 |
| | 20-40 times | .0 | .0 | .1 | .1 | .1 | .1 | .2 |
| | 40+ times | .1 | .0 | .1 | .1 | .1 | .2 | .2 |
| Heroin | No Answer | 3.0 | 2.1 | 2.8 | 2.9 | 2.7 | 2.7 | 2.6 |
| | Never | 96.8 | 97.5 | 96.6 | 96.3 | 96.4 | 96.2 | 96.2 |
| | 1-5 times | .1 | .3 | .4 | .4 | .5 | .6 | .5 |
| | 6-19 times | .0 | .1 | .2 | .2 | .2 | .2 | .3 |
| | 20-40 times | .0 | .0 | .1 | .1 | .1 | .2 | .2 |
| | 40+ times | .1 | .1 | .1 | .1 | .1 | .2 | .1 |
| Steroids | No Answer | 2.7 | 2.0 | 2.8 | 2.9 | 2.7 | 2.7 | 2.6 |
| | Never | 97.0 | 97.5 | 96.5 | 96.3 | 96.5 | 96.4 | 96.3 |
| | 1-5 times | .1 | .3 | .4 | .4 | .4 | .5 | .5 |
| | 6-19 times | .0 | .1 | .2 | .2 | .2 | .2 | .3 |
| | 20-40 times | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| | 40+ times | .0 | .1 | .1 | .1 | .1 | .2 | .1 |
| Ecstasy | No Answer | 3.7 | 2.3 | 2.8 | 3.0 | 2.8 | 2.7 | 2.7 |
| | Never | 96.1 | 97.2 | 96.1 | 95.6 | 95.5 | 95.5 | 95.2 |
| | 1-5 times | .2 | .3 | .8 | 1.0 | 1.3 | 1.1 | 1.4 |
| | 6-19 times | .1 | .1 | .2 | .3 | .2 | .4 | .4 |
| | 20-40 times | .0 | .1 | .1 | .1 | .1 | .2 | .2 |
| | 40+ times | .0 | .0 | .1 | .1 | .1 | .1 | .1 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | | Grade | | | | | | |
|--------------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Tranquilizers | No Answer | | 3.5 | 2.2 | 2.8 | 3.0 | 3.0 | 2.8 | 2.7 |
| | Never | | 96.2 | 97.3 | 96.2 | 95.4 | 95.0 | 95.3 | 95.3 |
| | 1-5 times | | .2 | .3 | .7 | 1.0 | 1.2 | 1.2 | 1.3 |
| | 6-19 times | | .0 | .1 | .2 | .4 | .4 | .5 | .4 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| | 40+ times | | .0 | .0 | .0 | .1 | .3 | .1 | .1 |
| Hallucinogens | No Answer | | 3.8 | 2.4 | 2.9 | 3.0 | 2.8 | 2.8 | 2.8 |
| | Never | | 96.0 | 97.1 | 95.9 | 95.4 | 95.3 | 95.0 | 94.8 |
| | 1-5 times | | .2 | .3 | .7 | 1.0 | 1.3 | 1.6 | 1.6 |
| | 6-19 times | | .0 | .1 | .3 | .3 | .3 | .4 | .5 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .1 | .1 | .2 |
| | 40+ times | | .0 | .1 | .1 | .1 | .1 | .1 | .2 |
| Prescription painkillers | No Answer | | 2.9 | 2.2 | 2.9 | 3.0 | 2.7 | 2.8 | 2.7 |
| | Never | | 96.5 | 96.4 | 94.0 | 92.3 | 91.5 | 91.3 | 90.8 |
| | 1-5 times | | .4 | 1.0 | 2.1 | 3.1 | 3.8 | 3.6 | 3.9 |
| | 6-19 times | | .1 | .3 | .6 | .9 | 1.2 | 1.3 | 1.6 |
| | 20-40 times | | .1 | .1 | .2 | .4 | .4 | .6 | .6 |
| | 40+ times | | .1 | .1 | .1 | .3 | .3 | .4 | .5 |
| Prescription drugs | No Answer | | 3.0 | 2.2 | 3.0 | 3.1 | 2.8 | 2.8 | 2.7 |
| | Never | | 96.4 | 96.4 | 94.3 | 92.6 | 92.0 | 91.6 | 91.3 |
| | 1-5 times | | .4 | .9 | 1.8 | 2.7 | 3.3 | 3.4 | 3.6 |
| | 6-19 times | | .1 | .3 | .5 | .9 | 1.0 | 1.2 | 1.3 |
| | 20-40 times | | .1 | .1 | .2 | .4 | .5 | .5 | .5 |
| | 40+ times | | .1 | .1 | .1 | .3 | .4 | .4 | .5 |
| Over the counter drugs | No Answer | | 3.2 | 2.6 | 3.2 | 3.4 | 3.1 | 3.0 | 3.0 |
| | Never | | 96.1 | 95.8 | 94.1 | 93.1 | 93.3 | 93.6 | 93.8 |
| | 1-5 times | | .5 | 1.2 | 1.9 | 2.4 | 2.4 | 2.2 | 2.1 |
| | 6-19 times | | .1 | .2 | .5 | .6 | .7 | .6 | .6 |
| | 20-40 times | | .0 | .1 | .2 | .3 | .2 | .3 | .2 |
| | 40+ times | | .1 | .1 | .2 | .3 | .2 | .3 | .2 |

Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | 2.2 | 2.1 | 2.5 | 2.9 | 2.7 | 2.6 | 2.6 |
| | Never | 91.1 | 85.8 | 78.5 | 72.0 | 66.4 | 61.3 | 57.4 |
| | 10 years or younger | 3.4 | 4.3 | 4.6 | 4.5 | 4.6 | 4.2 | 3.4 |
| | 11 years | 2.0 | 2.8 | 3.0 | 2.6 | 2.6 | 2.3 | 1.7 |
| | 12 years | 1.1 | 3.0 | 3.7 | 3.8 | 3.6 | 3.4 | 2.7 |
| | 13 years | .2 | 1.8 | 5.3 | 5.3 | 4.9 | 4.5 | 3.7 |
| | 14 years | .0 | .2 | 2.2 | 6.0 | 5.2 | 5.0 | 4.4 |
| | 15 years | .0 | .0 | .2 | 2.6 | 7.1 | 6.8 | 6.4 |
| | 16 years | .0 | .0 | .0 | .2 | 2.7 | 7.3 | 8.1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .2 | 2.5 | 9.6 |
| Smokeless Tobacco | No Answer | 2.5 | 2.3 | 2.9 | 3.2 | 3.2 | 3.1 | 3.2 |
| | Never | 95.5 | 93.4 | 89.7 | 85.2 | 81.2 | 78.6 | 75.0 |
| | 10 years or younger | .9 | 1.2 | 1.3 | 1.4 | 1.3 | 1.3 | 1.3 |
| | 11 years | .6 | .9 | 1.0 | .8 | .8 | .6 | .5 |
| | 12 years | .4 | 1.2 | 1.5 | 1.5 | 1.4 | 1.2 | 1.0 |
| | 13 years | .1 | .9 | 2.3 | 2.3 | 2.3 | 2.1 | 1.7 |
| | 14 years | .0 | .1 | 1.2 | 3.5 | 3.1 | 2.6 | 2.3 |
| | 15 years | .0 | .0 | .1 | 1.8 | 4.7 | 4.2 | 3.9 |
| | 16 years | .0 | .0 | .0 | .2 | 1.9 | 4.5 | 5.6 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.7 | 5.4 |
| Cigars | No Answer | 2.6 | 2.5 | 3.1 | 3.4 | 3.4 | 3.3 | 3.2 |
| | Never | 95.9 | 94.1 | 90.2 | 83.9 | 78.3 | 72.7 | 64.2 |
| | 10 years or younger | .7 | .9 | 1.0 | 1.1 | 1.2 | 1.2 | 1.1 |
| | 11 years | .4 | .7 | .9 | .9 | .8 | .7 | .6 |
| | 12 years | .3 | 1.0 | 1.3 | 1.6 | 1.4 | 1.1 | 1.1 |
| | 13 years | .1 | .7 | 2.2 | 2.8 | 2.6 | 2.4 | 1.9 |
| | 14 years | .0 | .1 | 1.1 | 4.0 | 3.6 | 3.3 | 2.9 |
| | 15 years | .0 | .0 | .2 | 2.1 | 6.1 | 5.8 | 5.2 |
| | 16 years | .0 | .0 | .0 | .1 | 2.5 | 6.9 | 8.6 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 2.4 | 11.3 |
| Pipe | No Answer | 2.8 | 2.7 | 3.4 | 4.0 | 4.0 | 3.9 | 3.9 |
| | Never | 96.5 | 95.2 | 92.3 | 88.0 | 84.0 | 79.4 | 73.1 |
| | 10 years or younger | .3 | .5 | .5 | .6 | .7 | .7 | .7 |
| | 11 years | .2 | .4 | .6 | .5 | .5 | .3 | .3 |
| | 12 years | .2 | .7 | .8 | .9 | .8 | .9 | .5 |
| | 13 years | .0 | .4 | 1.4 | 1.9 | 1.5 | 1.4 | 1.1 |
| | 14 years | .0 | .0 | .8 | 2.5 | 2.6 | 2.1 | 1.7 |
| | 15 years | .0 | .0 | .1 | 1.4 | 4.0 | 4.1 | 3.3 |
| | 16 years | .0 | .0 | .0 | .1 | 1.8 | 5.3 | 5.6 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.9 | 9.8 |

Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|-----------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Alcohol | No Answer | 3.2 | 2.9 | 3.5 | 3.6 | 3.5 | 3.5 | 3.2 |
| | Never | 81.6 | 74.1 | 62.7 | 52.8 | 45.2 | 40.0 | 35.0 |
| | 10 years or younger | 8.2 | 7.7 | 7.3 | 5.9 | 5.0 | 4.7 | 3.5 |
| | 11 years | 4.2 | 5.3 | 4.6 | 3.5 | 2.5 | 1.8 | 1.4 |
| | 12 years | 2.3 | 6.2 | 7.0 | 5.8 | 4.6 | 4.1 | 3.0 |
| | 13 years | .4 | 3.3 | 9.9 | 10.2 | 8.1 | 6.6 | 5.7 |
| | 14 years | .0 | .4 | 4.4 | 11.9 | 11.0 | 9.1 | 7.9 |
| | 15 years | .0 | .0 | .5 | 5.7 | 14.6 | 13.7 | 12.9 |
| | 16 years | .0 | .0 | .0 | .5 | 5.1 | 12.6 | 15.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .4 | 3.9 | 12.4 |
| | | | | | | | | |
| Marijuana | No Answer | 2.7 | 2.6 | 3.0 | 3.3 | 3.3 | 3.1 | 3.2 |
| | Never | 94.9 | 91.4 | 84.3 | 76.3 | 69.3 | 64.4 | 61.5 |
| | 10 years or younger | .6 | 1.1 | 1.4 | 1.3 | 1.5 | 1.3 | 1.0 |
| | 11 years | .8 | 1.3 | 1.3 | 1.3 | 1.1 | 1.0 | .8 |
| | 12 years | .7 | 1.9 | 2.4 | 2.5 | 2.2 | 2.1 | 1.8 |
| | 13 years | .2 | 1.5 | 4.8 | 4.8 | 4.0 | 3.4 | 2.8 |
| | 14 years | .0 | .2 | 2.5 | 6.8 | 5.7 | 5.1 | 4.1 |
| | 15 years | .0 | .0 | .3 | 3.4 | 9.0 | 8.0 | 6.7 |
| | 16 years | .0 | .0 | .0 | .3 | 3.6 | 8.9 | 9.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .3 | 2.7 | 9.1 |
| | | | | | | | | |
| Cocaine | No Answer | 2.7 | 2.4 | 3.0 | 3.5 | 3.5 | 3.4 | 3.4 |
| | Never | 97.0 | 96.9 | 95.6 | 94.6 | 93.7 | 92.7 | 92.1 |
| | 10 years or younger | .1 | .2 | .2 | .2 | .2 | .3 | .3 |
| | 11 years | .1 | .1 | .2 | .2 | .1 | .2 | .1 |
| | 12 years | .0 | .2 | .3 | .2 | .2 | .2 | .1 |
| | 13 years | .0 | .2 | .3 | .3 | .3 | .3 | .3 |
| | 14 years | .0 | .0 | .3 | .5 | .5 | .4 | .4 |
| | 15 years | .0 | .0 | .1 | .4 | .9 | .8 | .7 |
| | 16 years | .0 | .0 | .0 | .1 | .5 | 1.1 | 1.1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .6 | 1.6 |
| | | | | | | | | |
| Crack | No Answer | 2.8 | 2.5 | 3.1 | 3.6 | 3.5 | 3.4 | 3.4 |
| | Never | 96.9 | 96.8 | 95.8 | 95.2 | 95.1 | 94.9 | 94.8 |
| | 10 years or younger | .2 | .2 | .2 | .2 | .2 | .2 | .2 |
| | 11 years | .1 | .1 | .2 | .1 | .1 | .2 | .1 |
| | 12 years | .1 | .1 | .2 | .2 | .1 | .1 | .1 |
| | 13 years | .0 | .2 | .2 | .2 | .2 | .2 | .2 |
| | 14 years | .0 | .0 | .2 | .3 | .2 | .2 | .2 |
| | 15 years | .0 | .0 | .0 | .2 | .4 | .3 | .3 |
| | 16 years | .0 | .0 | .0 | .0 | .2 | .3 | .3 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .1 | .4 |
| | | | | | | | | |

Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Inhalants | No Answer | 3.6 | 2.8 | 3.1 | 3.6 | 3.6 | 3.5 | 3.5 |
| | Never | 95.2 | 95.1 | 93.1 | 91.9 | 91.4 | 91.1 | 91.2 |
| | 10 years or younger | .6 | .6 | .5 | .4 | .4 | .4 | .4 |
| | 11 years | .3 | .5 | .6 | .4 | .3 | .3 | .2 |
| | 12 years | .2 | .6 | .8 | .6 | .5 | .5 | .4 |
| | 13 years | .1 | .5 | 1.2 | 1.1 | .8 | .8 | .7 |
| | 14 years | .0 | .0 | .6 | 1.3 | 1.1 | 1.0 | .7 |
| | 15 years | .0 | .0 | .1 | .6 | 1.4 | 1.2 | .9 |
| | 16 years | .0 | .0 | .0 | .1 | .6 | 1.0 | 1.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .3 | 1.1 |
| Amphetamines | No Answer | 3.9 | 2.9 | 3.1 | 3.7 | 3.6 | 3.5 | 3.5 |
| | Never | 95.9 | 96.5 | 95.3 | 93.6 | 92.3 | 91.1 | 90.8 |
| | 10 years or younger | .1 | .1 | .2 | .2 | .2 | .3 | .2 |
| | 11 years | .1 | .1 | .2 | .1 | .1 | .2 | .1 |
| | 12 years | .0 | .2 | .3 | .3 | .3 | .3 | .3 |
| | 13 years | .0 | .2 | .5 | .6 | .6 | .6 | .4 |
| | 14 years | .0 | .0 | .3 | .9 | 1.0 | .9 | .7 |
| | 15 years | .0 | .0 | .1 | .5 | 1.4 | 1.3 | 1.1 |
| | 16 years | .0 | .0 | .0 | .0 | .6 | 1.4 | 1.6 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .4 | 1.3 |
| Methamphetamines | No Answer | 3.5 | 2.7 | 3.0 | 3.5 | 3.6 | 3.5 | 3.5 |
| | Never | 96.3 | 96.7 | 95.9 | 95.3 | 94.9 | 94.6 | 94.5 |
| | 10 years or younger | .1 | .2 | .2 | .2 | .1 | .3 | .2 |
| | 11 years | .0 | .2 | .1 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .1 | .2 | .1 | .1 | .1 | .1 |
| | 13 years | .0 | .1 | .3 | .2 | .2 | .1 | .1 |
| | 14 years | .0 | .0 | .2 | .3 | .3 | .2 | .1 |
| | 15 years | .0 | .0 | .1 | .2 | .4 | .4 | .3 |
| | 16 years | .0 | .0 | .0 | .0 | .3 | .5 | .4 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .3 | .7 |
| Heroin | No Answer | 3.4 | 2.7 | 3.2 | 3.6 | 3.6 | 3.5 | 3.5 |
| | Never | 96.4 | 96.9 | 96.0 | 95.4 | 95.1 | 94.9 | 94.8 |
| | 10 years or younger | .1 | .1 | .2 | .2 | .2 | .3 | .2 |
| | 11 years | .0 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .1 | .1 | .2 | .1 | .1 | .1 |
| | 13 years | .0 | .1 | .2 | .2 | .2 | .1 | .1 |
| | 14 years | .0 | .0 | .2 | .1 | .2 | .1 | .2 |
| | 15 years | .0 | .0 | .0 | .2 | .3 | .3 | .3 |
| | 16 years | .0 | .0 | .0 | .0 | .2 | .4 | .3 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .2 | .5 |

Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|---------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Steroids | No Answer | 3.1 | 2.7 | 3.2 | 3.6 | 3.7 | 3.6 | 3.5 |
| | Never | 96.6 | 96.7 | 95.9 | 95.5 | 95.2 | 95.2 | 95.1 |
| | 10 years or younger | .1 | .2 | .2 | .1 | .2 | .2 | .2 |
| | 11 years | .1 | .1 | .2 | .1 | .0 | .1 | .0 |
| | 12 years | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 13 years | .0 | .1 | .2 | .1 | .1 | .1 | .1 |
| | 14 years | .0 | .0 | .1 | .2 | .1 | .1 | .2 |
| | 15 years | .0 | .0 | .0 | .2 | .3 | .2 | .2 |
| | 16 years | .0 | .0 | .0 | .0 | .2 | .3 | .3 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .2 | .4 |
| | | | | | | | | |
| Ecstasy | No Answer | 4.1 | 2.9 | 3.3 | 3.7 | 3.6 | 3.5 | 3.4 |
| | Never | 95.6 | 96.4 | 95.2 | 94.0 | 93.2 | 92.3 | 91.6 |
| | 10 years or younger | .1 | .2 | .2 | .2 | .2 | .3 | .2 |
| | 11 years | .0 | .1 | .2 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .2 | .2 | .2 | .1 | .1 | .1 |
| | 13 years | .0 | .2 | .4 | .4 | .3 | .2 | .3 |
| | 14 years | .0 | .0 | .4 | .8 | .6 | .5 | .3 |
| | 15 years | .0 | .0 | .1 | .5 | 1.2 | .9 | .7 |
| | 16 years | .0 | .0 | .0 | .1 | .6 | 1.4 | 1.6 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .6 | 1.7 |
| | | | | | | | | |
| Tranquilizers | No Answer | 3.8 | 3.0 | 3.4 | 3.9 | 3.7 | 3.6 | 3.6 |
| | Never | 95.9 | 96.5 | 95.3 | 93.9 | 93.2 | 92.7 | 92.5 |
| | 10 years or younger | .1 | .1 | .2 | .2 | .2 | .3 | .2 |
| | 11 years | .1 | .1 | .2 | .1 | .2 | .1 | .1 |
| | 12 years | .1 | .1 | .2 | .2 | .2 | .2 | .2 |
| | 13 years | .0 | .2 | .4 | .6 | .4 | .4 | .4 |
| | 14 years | .0 | .0 | .3 | .8 | .7 | .6 | .5 |
| | 15 years | .0 | .0 | .0 | .3 | .9 | 1.1 | .7 |
| | 16 years | .0 | .0 | .0 | .1 | .4 | .7 | 1.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .3 | .8 |
| | | | | | | | | |
| Hallucinogens | No Answer | 4.2 | 3.0 | 3.5 | 3.9 | 3.9 | 3.7 | 3.8 |
| | Never | 95.6 | 96.4 | 95.0 | 93.7 | 92.6 | 91.7 | 90.9 |
| | 10 years or younger | .1 | .1 | .2 | .2 | .2 | .3 | .2 |
| | 11 years | .0 | .1 | .2 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .2 | .3 | .3 | .2 | .2 | .1 |
| | 13 years | .0 | .2 | .5 | .5 | .4 | .3 | .3 |
| | 14 years | .0 | .0 | .4 | .8 | .6 | .5 | .4 |
| | 15 years | .0 | .0 | .1 | .5 | 1.2 | .9 | .7 |
| | 16 years | .0 | .0 | .0 | .1 | .7 | 1.6 | 1.3 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .8 | 2.1 |
| | | | | | | | | |

Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|--------------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Prescription painkillers | No Answer | 3.4 | 2.7 | 3.3 | 3.8 | 3.6 | 3.6 | 3.6 |
| | Never | 95.9 | 95.3 | 92.4 | 88.7 | 86.2 | 84.3 | 83.4 |
| | 10 years or younger | .3 | .3 | .4 | .4 | .3 | .4 | .3 |
| | 11 years | .2 | .4 | .5 | .4 | .3 | .2 | .2 |
| | 12 years | .2 | .6 | .7 | .8 | .7 | .5 | .4 |
| | 13 years | .0 | .5 | 1.6 | 1.6 | 1.3 | 1.2 | .8 |
| | 14 years | .0 | .1 | .9 | 2.7 | 2.2 | 1.8 | 1.6 |
| | 15 years | .0 | .0 | .2 | 1.6 | 3.5 | 3.2 | 2.5 |
| | 16 years | .0 | .0 | .0 | .1 | 1.7 | 3.4 | 3.9 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.3 | 3.4 |
| | | | | | | | | |
| Prescription drugs | No Answer | 3.5 | 2.9 | 3.3 | 3.8 | 3.6 | 3.6 | 3.5 |
| | Never | 95.8 | 95.4 | 92.8 | 89.5 | 87.2 | 85.4 | 84.3 |
| | 10 years or younger | .4 | .4 | .4 | .4 | .4 | .4 | .3 |
| | 11 years | .1 | .3 | .4 | .3 | .2 | .2 | .2 |
| | 12 years | .1 | .5 | .6 | .6 | .5 | .4 | .3 |
| | 13 years | .0 | .5 | 1.4 | 1.4 | 1.2 | 1.1 | .7 |
| | 14 years | .0 | .1 | .8 | 2.4 | 1.9 | 1.7 | 1.4 |
| | 15 years | .0 | .0 | .1 | 1.3 | 3.3 | 2.8 | 2.3 |
| | 16 years | .0 | .0 | .0 | .1 | 1.6 | 3.2 | 3.5 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.3 | 3.4 |
| | | | | | | | | |
| Over the counter drugs | No Answer | 3.7 | 3.2 | 3.7 | 4.1 | 4.0 | 3.9 | 3.8 |
| | Never | 95.6 | 94.7 | 92.4 | 90.5 | 89.1 | 88.7 | 88.6 |
| | 10 years or younger | .3 | .4 | .4 | .5 | .3 | .4 | .3 |
| | 11 years | .2 | .3 | .4 | .3 | .3 | .1 | .2 |
| | 12 years | .2 | .7 | .8 | .7 | .5 | .4 | .3 |
| | 13 years | .1 | .6 | 1.4 | 1.2 | 1.0 | .8 | .6 |
| | 14 years | .0 | .1 | .8 | 1.7 | 1.5 | 1.3 | 1.0 |
| | 15 years | .0 | .0 | .1 | .9 | 2.2 | 1.9 | 1.4 |
| | 16 years | .0 | .0 | .0 | .1 | 1.0 | 1.9 | 2.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .6 | 1.7 |
| | | | | | | | | |

**Perceived Availability of Drugs
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------------------|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Availability of cigarettes | No Answer | 9.4 | 8.0 | 7.3 | 6.5 | 5.0 | 4.5 | 4.0 |
| | Very difficult | 62.4 | 53.2 | 42.5 | 32.3 | 25.8 | 19.2 | 9.3 |
| | Fairly difficult | 10.2 | 12.7 | 13.5 | 12.6 | 11.4 | 8.7 | 3.4 |
| | Fairly easy | 9.0 | 13.3 | 18.0 | 21.6 | 23.0 | 22.2 | 10.5 |
| | Easy | 9.1 | 12.8 | 18.8 | 26.9 | 34.7 | 45.3 | 72.8 |
| Availability of alcohol | No Answer | 9.7 | 8.1 | 7.3 | 6.3 | 4.8 | 4.3 | 4.0 |
| | Very difficult | 53.4 | 42.8 | 32.5 | 23.5 | 18.3 | 15.3 | 11.8 |
| | Fairly difficult | 14.2 | 16.2 | 15.7 | 14.3 | 13.0 | 10.8 | 10.8 |
| | Fairly easy | 11.6 | 16.7 | 21.0 | 25.2 | 27.7 | 28.9 | 32.5 |
| | Easy | 11.1 | 16.3 | 23.5 | 30.6 | 36.3 | 40.6 | 41.0 |
| Availability of marijuana | No Answer | 10.4 | 8.8 | 8.0 | 7.1 | 5.6 | 5.0 | 4.6 |
| | Very difficult | 81.3 | 74.4 | 61.3 | 45.5 | 35.9 | 29.7 | 24.4 |
| | Fairly difficult | 3.6 | 6.2 | 9.5 | 11.4 | 11.6 | 11.5 | 12.1 |
| | Fairly easy | 2.0 | 4.8 | 9.6 | 15.0 | 18.4 | 20.2 | 23.2 |
| | Easy | 2.7 | 5.8 | 11.6 | 20.9 | 28.4 | 33.6 | 35.7 |
| Availability of drugs | No Answer | 10.6 | 9.0 | 8.4 | 7.9 | 6.4 | 5.7 | 5.1 |
| | Very difficult | 84.6 | 81.4 | 75.4 | 64.9 | 59.1 | 53.6 | 49.5 |
| | Fairly difficult | 2.4 | 4.9 | 8.2 | 13.6 | 17.0 | 19.0 | 21.7 |
| | Fairly easy | 1.0 | 2.1 | 4.0 | 7.0 | 9.5 | 11.6 | 12.6 |
| | Easy | 1.5 | 2.6 | 3.9 | 6.7 | 8.0 | 10.0 | 11.2 |

**Binge Drinking in the Past Two Weeks
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Binge drinking | No Answer | 3.1 | 2.6 | 2.1 | 2.4 | 2.1 | 2.4 | 2.2 |
| | None | 91.0 | 89.5 | 86.7 | 82.3 | 79.7 | 76.8 | 71.3 |
| | Once | 2.5 | 3.2 | 4.4 | 6.1 | 7.1 | 8.1 | 9.8 |
| | Twice | 1.2 | 1.8 | 2.7 | 3.7 | 4.3 | 4.8 | 6.5 |
| | 3-5 times | 1.2 | 1.6 | 2.4 | 3.2 | 4.1 | 4.8 | 6.4 |
| | 6-9 times | .4 | .6 | .8 | 1.1 | 1.3 | 1.6 | 2.0 |
| | 10+ times | .5 | .7 | .8 | 1.3 | 1.3 | 1.5 | 1.8 |

**Main Sources of Alcoholic Beverages in the Past Month
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------------|---|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Main sources of alcoholic beverages | No Answer | 6.5 | 5.7 | 5.1 | 5.7 | 6.0 | 6.2 | 7.0 |
| | No drink | 88.0 | 84.3 | 78.7 | 71.9 | 67.4 | 64.4 | 57.6 |
| | Restaurants/bars/clubs | .1 | .0 | .1 | .1 | .2 | .2 | .3 |
| | Public events | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| | Had someone else buy it | .3 | .9 | 1.9 | 4.3 | 6.9 | 9.3 | 12.6 |
| | Liquor stores/convenience stores/supermarkets | .1 | .1 | .1 | .2 | .3 | .5 | .9 |
| | Received from person 21 or older | .8 | 1.4 | 2.4 | 3.7 | 5.4 | 6.9 | 9.9 |
| | Took it from a store | .1 | .2 | .3 | .3 | .3 | .5 | .3 |
| | Family members | 2.0 | 3.4 | 5.0 | 5.4 | 5.0 | 4.4 | 4.1 |
| | Other ways | 2.0 | 4.0 | 6.5 | 8.3 | 8.5 | 7.6 | 7.2 |

Most Important Reasons for Drinking by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | | Grade | | | | | | |
|---|-----------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Did not drink alcoholic beverages | No answer | | 18.6 | 26.3 | 36.6 | 46.1 | 52.7 | 56.6 | 62.4 |
| | Yes | | 81.4 | 73.7 | 63.4 | 53.9 | 47.3 | 43.4 | 37.6 |
| To experiment | No answer | | 89.2 | 83.9 | 77.9 | 76.1 | 75.4 | 77.4 | 78.2 |
| | Yes | | 10.8 | 16.1 | 22.1 | 23.9 | 24.6 | 22.6 | 21.8 |
| To relax or relieve tension | No answer | | 98.7 | 96.6 | 92.8 | 88.7 | 84.8 | 81.4 | 78.6 |
| | Yes | | 1.3 | 3.4 | 7.2 | 11.3 | 15.2 | 18.6 | 21.4 |
| To feel good or get high | No answer | | 98.9 | 97.3 | 94.8 | 91.1 | 87.9 | 85.4 | 82.9 |
| | Yes | | 1.1 | 2.7 | 5.2 | 8.9 | 12.1 | 14.6 | 17.1 |
| To seek deeper insights and understanding | No answer | | 99.5 | 99.2 | 98.5 | 97.7 | 97.7 | 97.3 | 96.7 |
| | Yes | | .5 | .8 | 1.5 | 2.3 | 2.3 | 2.7 | 3.3 |
| To have a good time with friends | No answer | | 98.0 | 94.4 | 88.5 | 79.4 | 72.2 | 66.7 | 60.3 |
| | Yes | | 2.0 | 5.6 | 11.5 | 20.6 | 27.8 | 33.3 | 39.7 |
| To fit in with a group I like | No answer | | 99.1 | 98.4 | 97.9 | 97.6 | 97.4 | 97.2 | 97.1 |
| | Yes | | .9 | 1.6 | 2.1 | 2.4 | 2.6 | 2.8 | 2.9 |
| To get away from my problems | No answer | | 98.2 | 95.9 | 93.0 | 90.6 | 89.2 | 88.6 | 88.8 |
| | Yes | | 1.8 | 4.1 | 7.0 | 9.4 | 10.8 | 11.4 | 11.2 |
| Because of boredom | No answer | | 98.1 | 95.7 | 92.7 | 90.8 | 88.7 | 88.1 | 87.4 |
| | Yes | | 1.9 | 4.3 | 7.3 | 9.2 | 11.3 | 11.9 | 12.6 |
| Because of anger | No answer | | 98.4 | 97.1 | 95.0 | 93.8 | 93.2 | 92.8 | 93.1 |
| | Yes | | 1.6 | 2.9 | 5.0 | 6.2 | 6.8 | 7.2 | 6.9 |
| To get through the day | No answer | | 99.4 | 98.8 | 98.1 | 97.5 | 97.5 | 97.5 | 97.6 |
| | Yes | | .6 | 1.2 | 1.9 | 2.5 | 2.5 | 2.5 | 2.4 |
| To increase the effects of other drugs | No answer | | 99.7 | 99.5 | 99.0 | 98.3 | 97.8 | 97.5 | 97.2 |
| | Yes | | .3 | .5 | 1.0 | 1.7 | 2.2 | 2.5 | 2.8 |
| To decrease the effects of other drugs | No answer | | 99.8 | 99.8 | 99.7 | 99.6 | 99.6 | 99.6 | 99.5 |
| | Yes | | .2 | .2 | .3 | .4 | .4 | .4 | .5 |
| To get to sleep | No answer | | 99.4 | 99.1 | 98.3 | 97.8 | 97.5 | 97.1 | 96.4 |
| | Yes | | .6 | .9 | 1.7 | 2.2 | 2.5 | 2.9 | 3.6 |
| Because it tastes good | No answer | | 97.3 | 94.1 | 90.1 | 86.4 | 84.5 | 82.7 | 81.1 |
| | Yes | | 2.7 | 5.9 | 9.9 | 13.6 | 15.5 | 17.3 | 18.9 |
| Because I am hooked | No answer | | 99.5 | 99.2 | 99.0 | 98.8 | 98.8 | 98.8 | 98.6 |
| | Yes | | .5 | .8 | 1.0 | 1.2 | 1.2 | 1.2 | 1.4 |

Consequences of Alcohol, Tobacco, and Other Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|-----------------------|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Had a hangover | No Answer | 3.9 | 3.5 | 2.9 | 3.6 | 3.3 | 3.2 | 3.0 |
| | Never | 92.7 | 88.3 | 82.3 | 74.1 | 68.8 | 64.8 | 59.4 |
| | Once | 1.8 | 3.7 | 6.1 | 8.5 | 10.2 | 11.6 | 11.7 |
| | Twice | .6 | 1.7 | 3.2 | 4.8 | 5.8 | 6.5 | 7.3 |
| | 3-5 times | .4 | 1.4 | 2.6 | 4.4 | 5.9 | 6.7 | 8.9 |
| | 6-10 times | .2 | .4 | 1.1 | 1.7 | 2.4 | 2.7 | 4.1 |
| | 11 or more times | .4 | 1.0 | 1.9 | 2.9 | 3.6 | 4.6 | 5.6 |
| Had nausea or vomited | No Answer | 4.2 | 3.8 | 3.2 | 3.8 | 3.6 | 3.5 | 3.3 |
| | Never | 92.5 | 89.4 | 85.3 | 78.8 | 73.3 | 68.1 | 62.7 |
| | Once | 1.9 | 3.7 | 5.7 | 8.2 | 10.4 | 12.1 | 12.9 |
| | Twice | .6 | 1.4 | 2.3 | 3.4 | 4.7 | 5.6 | 7.4 |
| | 3-5 times | .4 | .8 | 1.9 | 3.2 | 4.4 | 6.0 | 7.7 |
| | 6-10 times | .1 | .4 | .7 | 1.2 | 1.7 | 2.3 | 2.9 |
| | 11 or more times | .3 | .6 | .9 | 1.3 | 1.8 | 2.3 | 2.9 |
| Had a memory loss | No Answer | 4.4 | 4.1 | 3.4 | 4.1 | 3.9 | 3.7 | 3.5 |
| | Never | 93.5 | 91.6 | 88.5 | 82.9 | 78.1 | 75.4 | 71.8 |
| | Once | 1.2 | 2.1 | 3.7 | 5.4 | 7.2 | 7.7 | 8.5 |
| | Twice | .4 | .8 | 1.6 | 2.4 | 3.4 | 3.9 | 4.7 |
| | 3-5 times | .2 | .6 | 1.3 | 2.4 | 3.4 | 4.4 | 5.6 |
| | 6-10 times | .1 | .2 | .6 | 1.0 | 1.6 | 2.1 | 2.4 |
| | 11 or more times | .2 | .5 | 1.0 | 1.7 | 2.4 | 2.7 | 3.5 |
| Poor on school test | No Answer | 4.5 | 4.0 | 3.5 | 4.1 | 3.9 | 3.7 | 3.5 |
| | Never | 92.7 | 91.7 | 90.7 | 89.0 | 88.7 | 89.0 | 89.4 |
| | Once | 1.3 | 1.8 | 2.2 | 2.4 | 2.3 | 2.3 | 2.1 |
| | Twice | .5 | .8 | 1.1 | 1.2 | 1.3 | 1.2 | 1.3 |
| | 3-5 times | .4 | .7 | 1.1 | 1.4 | 1.5 | 1.6 | 1.6 |
| | 6-10 times | .1 | .3 | .5 | .6 | .7 | .8 | .7 |
| | 11 or more times | .4 | .7 | 1.0 | 1.4 | 1.5 | 1.4 | 1.4 |
| Missed school | No Answer | 4.6 | 4.1 | 3.5 | 4.1 | 3.9 | 3.7 | 3.5 |
| | Never | 92.9 | 92.5 | 91.6 | 90.1 | 89.8 | 89.8 | 88.8 |
| | Once | .9 | 1.2 | 1.7 | 1.8 | 2.0 | 2.2 | 2.3 |
| | Twice | .5 | .7 | 1.0 | 1.1 | 1.2 | 1.1 | 1.4 |
| | 3-5 times | .6 | .7 | .9 | 1.1 | 1.2 | 1.3 | 1.6 |
| | 6-10 times | .2 | .3 | .5 | .7 | .7 | .7 | .8 |
| | 11 or more times | .4 | .6 | .8 | 1.1 | 1.1 | 1.1 | 1.5 |

**Consequences of Alcohol, Tobacco, and Other Drug Use
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Got into a fight | No Answer | 4.6 | 4.1 | 3.6 | 4.2 | 4.0 | 3.8 | 3.6 |
| | Never | 91.1 | 89.5 | 87.4 | 85.1 | 83.9 | 83.7 | 82.4 |
| | Once | 2.0 | 2.6 | 3.6 | 4.2 | 4.7 | 4.6 | 5.4 |
| | Twice | .7 | 1.2 | 1.7 | 2.1 | 2.6 | 2.6 | 3.2 |
| | 3-5 times | .6 | 1.1 | 1.6 | 1.9 | 2.2 | 2.7 | 2.7 |
| | 6-10 times | .3 | .5 | .8 | 1.0 | 1.0 | 1.0 | 1.1 |
| | 11 or more times | .7 | 1.1 | 1.4 | 1.6 | 1.7 | 1.6 | 1.6 |
| | | | | | | | | |
| Damaged property | No Answer | 4.6 | 4.2 | 3.5 | 4.0 | 3.9 | 3.7 | 3.5 |
| | Never | 94.0 | 93.4 | 92.8 | 91.6 | 91.6 | 91.8 | 91.3 |
| | Once | .6 | 1.0 | 1.4 | 1.5 | 1.6 | 1.5 | 1.8 |
| | Twice | .3 | .4 | .6 | .8 | .7 | .9 | 1.0 |
| | 3-5 times | .2 | .4 | .7 | .8 | .9 | .8 | 1.0 |
| | 6-10 times | .1 | .2 | .3 | .4 | .4 | .4 | .4 |
| | 11 or more times | .2 | .4 | .6 | .9 | .8 | .8 | 1.0 |
| | | | | | | | | |
| Driven by someone under the influence | No Answer | 3.6 | 3.0 | 2.6 | 3.0 | 2.8 | 2.9 | 2.7 |
| | Yes | 19.9 | 24.5 | 32.5 | 38.6 | 44.7 | 48.0 | 50.7 |
| | No | 76.5 | 72.4 | 64.9 | 58.4 | 52.4 | 49.1 | 46.6 |
| Used alcohol or drugs to relax or fit in | No Answer | 4.1 | 3.7 | 3.0 | 3.4 | 3.3 | 3.3 | 3.0 |
| | Yes | 2.9 | 5.9 | 10.1 | 14.9 | 18.5 | 20.8 | 23.3 |
| | No | 93.0 | 90.4 | 86.9 | 81.8 | 78.2 | 75.9 | 73.7 |
| Used alcohol or drugs alone | No Answer | 4.6 | 4.1 | 3.3 | 3.8 | 3.4 | 3.4 | 3.1 |
| | Yes | 3.3 | 7.6 | 12.4 | 17.2 | 20.4 | 21.4 | 22.1 |
| | No | 92.1 | 88.3 | 84.3 | 79.0 | 76.3 | 75.2 | 74.7 |
| Forgot things you did while high | No Answer | 7.4 | 6.0 | 4.8 | 5.0 | 4.3 | 4.1 | 3.7 |
| | Yes | 2.1 | 4.6 | 8.6 | 13.8 | 18.0 | 21.2 | 24.6 |
| | No | 90.5 | 89.4 | 86.6 | 81.2 | 77.6 | 74.6 | 71.7 |
| Had been told to cut down | No Answer | 9.0 | 7.3 | 5.4 | 5.5 | 4.5 | 4.2 | 3.7 |
| | Yes | 2.1 | 3.3 | 5.1 | 7.1 | 8.3 | 8.7 | 8.9 |
| | No | 88.9 | 89.4 | 89.5 | 87.4 | 87.2 | 87.0 | 87.4 |
| Got into trouble | No Answer | 9.3 | 7.2 | 5.3 | 5.5 | 4.5 | 4.2 | 3.8 |
| | Yes | 2.6 | 4.1 | 6.7 | 9.4 | 11.2 | 12.5 | 12.8 |
| | No | 88.2 | 88.6 | 88.1 | 85.2 | 84.4 | 83.3 | 83.4 |

Gambling during the Last Twelve Months by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Played cards | No Answer | 2.9 | 2.6 | 2.5 | 3.1 | 3.1 | 3.0 | 3.0 |
| | Not at all | 82.2 | 78.8 | 76.3 | 74.9 | 75.1 | 75.3 | 73.4 |
| | Less than once a month | 7.8 | 9.6 | 11.0 | 11.4 | 11.4 | 11.3 | 12.4 |
| | About once a month | 3.8 | 5.0 | 5.9 | 6.0 | 6.0 | 6.1 | 6.6 |
| | About once a week | 2.3 | 2.7 | 3.2 | 3.3 | 3.1 | 3.0 | 3.3 |
| | Daily | 1.0 | 1.3 | 1.1 | 1.4 | 1.3 | 1.3 | 1.4 |
| Bet on games | No Answer | 3.1 | 2.8 | 2.8 | 3.2 | 3.4 | 3.2 | 3.2 |
| | Not at all | 81.7 | 77.7 | 75.4 | 73.6 | 74.4 | 75.1 | 73.9 |
| | Less than once a month | 7.2 | 8.7 | 9.5 | 9.8 | 9.4 | 9.0 | 10.0 |
| | About once a month | 3.8 | 4.9 | 5.9 | 6.4 | 6.0 | 6.0 | 6.2 |
| | About once a week | 2.4 | 3.2 | 3.8 | 4.0 | 4.0 | 4.0 | 4.1 |
| | Daily | 1.8 | 2.6 | 2.6 | 2.9 | 2.7 | 2.6 | 2.6 |
| Bet on sports | No Answer | 3.4 | 2.8 | 3.1 | 3.4 | 3.5 | 3.3 | 3.3 |
| | Not at all | 82.7 | 79.5 | 77.8 | 77.3 | 78.8 | 80.7 | 80.2 |
| | Less than once a month | 8.1 | 9.9 | 10.6 | 10.6 | 9.8 | 8.9 | 9.4 |
| | About once a month | 2.6 | 3.9 | 4.3 | 4.3 | 3.8 | 3.5 | 3.8 |
| | About once a week | 1.9 | 2.2 | 2.5 | 2.3 | 2.2 | 2.0 | 1.8 |
| | Daily | 1.3 | 1.7 | 1.8 | 2.1 | 1.8 | 1.6 | 1.5 |
| Bought lottery tickets | No Answer | 3.4 | 3.1 | 3.0 | 3.5 | 3.6 | 3.5 | 3.3 |
| | Not at all | 82.5 | 80.7 | 80.2 | 79.9 | 80.8 | 80.8 | 67.4 |
| | Less than once a month | 8.9 | 9.8 | 10.4 | 10.3 | 10.0 | 9.8 | 18.5 |
| | About once a month | 3.1 | 3.8 | 4.0 | 3.8 | 3.5 | 3.6 | 6.7 |
| | About once a week | 1.5 | 1.7 | 1.8 | 1.6 | 1.4 | 1.4 | 3.0 |
| | Daily | .6 | .9 | .7 | .9 | .8 | .9 | 1.2 |
| Gambled in a casino | No Answer | 3.7 | 3.0 | 3.2 | 3.5 | 3.6 | 3.3 | 3.2 |
| | Not at all | 95.6 | 95.9 | 95.6 | 95.0 | 95.0 | 94.9 | 94.2 |
| | Less than once a month | .3 | .5 | .6 | .5 | .6 | .7 | 1.4 |
| | About once a month | .1 | .1 | .2 | .2 | .2 | .3 | .4 |
| | About once a week | .1 | .1 | .1 | .1 | .1 | .2 | .3 |
| | Daily | .2 | .3 | .3 | .6 | .5 | .6 | .6 |
| Played online | No Answer | 3.6 | 3.0 | 3.1 | 3.5 | 3.7 | 3.4 | 3.2 |
| | Not at all | 87.3 | 86.6 | 86.0 | 86.6 | 87.6 | 88.7 | 88.7 |
| | Less than once a month | 2.6 | 3.1 | 3.3 | 2.8 | 2.7 | 2.2 | 2.4 |
| | About once a month | 1.2 | 1.4 | 1.6 | 1.5 | 1.5 | 1.5 | 1.4 |
| | About once a week | 1.9 | 2.0 | 2.2 | 1.9 | 1.7 | 1.5 | 1.6 |
| | Daily | 3.5 | 3.8 | 3.8 | 3.6 | 2.9 | 2.7 | 2.6 |
| Felt bad about the amount bet | No Answer | 10.7 | 9.9 | 9.4 | 9.0 | 8.8 | 8.1 | 7.4 |
| | Yes | 3.1 | 2.9 | 2.6 | 2.5 | 2.3 | 2.3 | 2.6 |
| | No | 15.1 | 17.6 | 19.8 | 20.6 | 20.1 | 19.7 | 23.7 |
| | I do not bet for money | 71.2 | 69.5 | 68.3 | 67.9 | 68.7 | 69.9 | 66.3 |
| Would like to stop betting but could not | No Answer | 11.3 | 10.4 | 9.7 | 9.4 | 9.1 | 8.3 | 7.5 |
| | Yes | 2.4 | 2.0 | 2.0 | 1.7 | 1.4 | 1.4 | 1.4 |
| | No | 15.4 | 18.0 | 19.9 | 20.7 | 20.4 | 20.0 | 24.3 |
| | I do not bet for money | 70.9 | 69.6 | 68.4 | 68.2 | 69.1 | 70.3 | 66.8 |

**Perceived Risk of Selected Substance Use Behavior
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|---------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Smoke 1+ pack cigarettes per day | No Answer | 3.8 | 3.3 | 2.9 | 3.8 | 3.6 | 3.5 | 3.6 |
| | No risk | 7.0 | 5.6 | 4.9 | 4.7 | 4.5 | 4.6 | 4.3 |
| | Slight risk | 10.3 | 8.0 | 8.7 | 8.4 | 7.4 | 7.6 | 7.3 |
| | Moderate risk | 23.3 | 22.9 | 24.6 | 23.8 | 22.7 | 23.0 | 22.4 |
| | Great risk | 55.7 | 60.2 | 58.9 | 59.3 | 61.7 | 61.3 | 62.5 |
| Smoke marijuana occasionally | No Answer | 4.1 | 3.6 | 3.2 | 4.0 | 3.8 | 3.7 | 3.7 |
| | No risk | 7.5 | 7.9 | 10.7 | 15.7 | 18.9 | 22.3 | 23.1 |
| | Slight risk | 10.1 | 12.2 | 17.0 | 21.2 | 24.2 | 25.3 | 27.1 |
| | Moderate risk | 29.5 | 29.8 | 29.9 | 27.8 | 26.4 | 24.7 | 24.0 |
| | Great risk | 48.9 | 46.5 | 39.3 | 31.3 | 26.8 | 24.1 | 22.2 |
| Smoke marijuana regularly | No Answer | 4.8 | 4.2 | 3.9 | 4.7 | 4.3 | 4.2 | 4.2 |
| | No risk | 7.1 | 6.4 | 7.5 | 9.9 | 11.0 | 12.3 | 13.0 |
| | Slight risk | 4.6 | 5.1 | 7.9 | 11.4 | 13.9 | 15.5 | 16.6 |
| | Moderate risk | 12.3 | 12.5 | 15.8 | 19.3 | 21.1 | 22.7 | 23.6 |
| | Great risk | 71.2 | 71.7 | 65.0 | 54.7 | 49.7 | 45.3 | 42.6 |
| Occasionally consume 1-2 alcoholic drinks | No Answer | 4.8 | 4.3 | 3.8 | 4.5 | 4.3 | 4.1 | 4.0 |
| | No risk | 20.0 | 21.0 | 24.2 | 29.3 | 29.7 | 33.5 | 35.6 |
| | Slight risk | 37.8 | 38.3 | 39.0 | 36.7 | 37.9 | 36.2 | 37.0 |
| | Moderate risk | 19.6 | 18.7 | 17.4 | 15.8 | 15.4 | 14.1 | 12.4 |
| | Great risk | 17.8 | 17.7 | 15.6 | 13.7 | 12.7 | 12.2 | 11.0 |
| Binge drink weekly | No Answer | 4.5 | 3.7 | 3.4 | 4.1 | 4.0 | 3.9 | 3.8 |
| | No risk | 8.4 | 7.1 | 6.8 | 7.0 | 6.6 | 7.2 | 6.9 |
| | Slight risk | 13.4 | 13.0 | 13.9 | 15.9 | 15.5 | 16.1 | 17.8 |
| | Moderate risk | 29.3 | 29.9 | 31.5 | 32.0 | 31.5 | 32.4 | 33.5 |
| | Great risk | 44.4 | 46.3 | 44.4 | 41.0 | 42.4 | 40.4 | 38.0 |

**Perceived Peer Approval of Using Selected Substances
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Smoke 1+ pack cigarettes per day | No Answer | 5.4 | 4.7 | 3.8 | 4.6 | 4.2 | 4.1 | 4.1 |
| | Strongly approve | 2.2 | 2.2 | 2.1 | 2.3 | 2.4 | 2.8 | 2.8 |
| | Approve | 1.0 | 1.8 | 3.0 | 4.5 | 5.2 | 6.4 | 7.1 |
| | Do not know | 11.6 | 12.8 | 15.6 | 17.2 | 17.7 | 17.4 | 18.5 |
| | Disapprove | 9.8 | 11.4 | 14.1 | 17.0 | 17.5 | 18.5 | 20.5 |
| | Strongly disapprove | 70.1 | 67.1 | 61.4 | 54.4 | 52.9 | 50.8 | 47.1 |
| Smoke marijuana occasionally | No Answer | 5.5 | 4.7 | 3.9 | 4.7 | 4.5 | 4.3 | 4.2 |
| | Strongly approve | 2.4 | 2.7 | 3.4 | 4.8 | 6.0 | 6.9 | 6.8 |
| | Approve | 1.4 | 3.3 | 7.3 | 11.3 | 15.3 | 17.2 | 18.4 |
| | Do not know | 9.9 | 11.7 | 14.9 | 17.4 | 18.0 | 18.3 | 19.1 |
| | Disapprove | 7.3 | 8.7 | 11.0 | 13.7 | 13.8 | 13.8 | 15.0 |
| | Strongly disapprove | 73.6 | 68.9 | 59.5 | 48.0 | 42.5 | 39.4 | 36.4 |
| Smoke marijuana regularly | No Answer | 5.8 | 4.9 | 4.2 | 5.1 | 4.8 | 4.5 | 4.5 |
| | Strongly approve | 2.5 | 2.8 | 3.6 | 4.7 | 5.8 | 6.6 | 6.5 |
| | Approve | .9 | 2.1 | 4.5 | 7.1 | 9.5 | 10.2 | 10.7 |
| | Do not know | 9.5 | 10.8 | 13.9 | 16.0 | 16.7 | 16.9 | 17.2 |
| | Disapprove | 5.3 | 6.4 | 8.6 | 11.5 | 12.5 | 13.4 | 14.8 |
| | Strongly disapprove | 76.0 | 72.9 | 65.2 | 55.6 | 50.8 | 48.4 | 46.4 |
| Occasionally consume 1-2 alcoholic drinks | No Answer | 5.7 | 5.0 | 4.2 | 4.8 | 4.7 | 4.4 | 4.4 |
| | Strongly approve | 2.5 | 3.0 | 3.7 | 4.9 | 6.1 | 7.5 | 8.8 |
| | Approve | 3.7 | 6.7 | 12.4 | 19.1 | 23.5 | 27.1 | 31.4 |
| | Do not know | 15.5 | 17.6 | 21.6 | 24.2 | 24.6 | 24.0 | 23.5 |
| | Disapprove | 12.3 | 12.9 | 13.7 | 14.1 | 13.6 | 12.6 | 11.2 |
| | Strongly disapprove | 60.3 | 54.9 | 44.4 | 32.9 | 27.5 | 24.4 | 20.6 |
| Binge drink weekly | No Answer | 5.7 | 4.9 | 4.1 | 4.9 | 4.6 | 4.5 | 4.4 |
| | Strongly approve | 2.4 | 2.6 | 2.9 | 3.7 | 4.3 | 5.0 | 5.5 |
| | Approve | 1.1 | 2.4 | 4.5 | 7.1 | 9.1 | 10.7 | 12.9 |
| | Do not know | 11.0 | 12.8 | 16.6 | 19.9 | 20.8 | 20.6 | 21.9 |
| | Disapprove | 7.9 | 9.1 | 12.1 | 15.4 | 16.2 | 17.4 | 18.0 |
| | Strongly disapprove | 71.9 | 68.1 | 59.7 | 49.0 | 44.9 | 41.9 | 37.3 |

**Parental Attitudes toward Drug Use and Antisocial Behavior
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Parents feel about alcohol | No Answer | 4.5 | 4.0 | 3.5 | 4.4 | 4.4 | 4.1 | 4.4 |
| | Very wrong | 84.7 | 81.7 | 76.6 | 69.1 | 64.5 | 59.6 | 51.2 |
| | Wrong | 6.5 | 8.1 | 10.9 | 14.0 | 15.8 | 16.7 | 18.7 |
| | A little bit wrong | 3.1 | 4.3 | 6.5 | 9.4 | 11.4 | 14.3 | 18.4 |
| | Not at all wrong | 1.3 | 1.8 | 2.5 | 3.2 | 3.9 | 5.4 | 7.4 |
| Parents feel about cigarettes | No Answer | 4.4 | 4.1 | 3.5 | 4.5 | 4.4 | 4.2 | 4.3 |
| | Very wrong | 88.6 | 87.0 | 84.1 | 79.3 | 76.5 | 71.4 | 62.3 |
| | Wrong | 4.9 | 6.0 | 7.9 | 10.1 | 11.6 | 13.8 | 17.5 |
| | A little bit wrong | 1.0 | 1.5 | 2.6 | 3.7 | 4.6 | 6.5 | 9.5 |
| | Not at all wrong | 1.0 | 1.4 | 1.8 | 2.3 | 3.0 | 4.1 | 6.3 |
| Parents feel about marijuana | No Answer | 4.6 | 4.1 | 3.6 | 4.6 | 4.5 | 4.3 | 4.5 |
| | Very wrong | 91.8 | 90.0 | 86.8 | 81.4 | 78.3 | 75.4 | 72.0 |
| | Wrong | 2.0 | 3.2 | 5.3 | 7.6 | 9.2 | 10.6 | 12.6 |
| | A little bit wrong | .5 | 1.2 | 2.2 | 3.7 | 4.8 | 5.8 | 6.8 |
| | Not at all wrong | 1.0 | 1.4 | 2.1 | 2.8 | 3.2 | 4.0 | 4.2 |
| Parents feel about stealing | No Answer | 4.7 | 4.2 | 3.7 | 4.7 | 4.6 | 4.4 | 4.5 |
| | Very wrong | 83.9 | 82.8 | 81.1 | 78.5 | 79.1 | 79.1 | 78.7 |
| | Wrong | 8.6 | 9.3 | 10.6 | 12.0 | 12.1 | 11.8 | 12.5 |
| | A little bit wrong | 1.7 | 2.3 | 2.8 | 3.0 | 2.7 | 2.8 | 2.5 |
| | Not at all wrong | 1.0 | 1.4 | 1.8 | 1.7 | 1.6 | 1.9 | 1.9 |
| Parents feel about graffiti | No Answer | 4.9 | 4.3 | 3.8 | 4.7 | 4.5 | 4.4 | 4.4 |
| | Very wrong | 85.2 | 84.0 | 81.0 | 78.1 | 77.7 | 76.9 | 75.6 |
| | Wrong | 6.7 | 7.8 | 9.9 | 11.5 | 12.2 | 12.6 | 13.8 |
| | A little bit wrong | 1.8 | 2.3 | 3.3 | 3.7 | 3.5 | 3.8 | 3.9 |
| | Not at all wrong | 1.4 | 1.7 | 2.1 | 2.1 | 2.1 | 2.3 | 2.3 |
| Parents feel about fighting | No Answer | 4.9 | 4.4 | 3.9 | 4.9 | 4.6 | 4.5 | 4.5 |
| | Very wrong | 67.6 | 63.6 | 58.0 | 56.2 | 55.5 | 56.6 | 55.4 |
| | Wrong | 17.2 | 18.6 | 20.7 | 21.4 | 22.4 | 22.5 | 24.4 |
| | A little bit wrong | 7.6 | 9.9 | 12.9 | 13.4 | 13.4 | 12.5 | 12.0 |
| | Not at all wrong | 2.7 | 3.5 | 4.5 | 4.1 | 4.0 | 3.9 | 3.6 |

**Family Management and Conflict
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Clear rules | No Answer | 3.4 | 3.7 | 3.3 | 5.0 | 5.0 | 5.1 | 5.2 |
| | YES! | 65.1 | 60.0 | 55.8 | 52.8 | 53.2 | 54.1 | 53.1 |
| | yes | 27.2 | 30.6 | 33.9 | 34.9 | 34.9 | 33.8 | 34.1 |
| | no | 2.9 | 4.2 | 5.2 | 5.4 | 5.2 | 5.0 | 5.5 |
| | NO! | 1.3 | 1.6 | 1.8 | 1.9 | 1.8 | 2.0 | 2.1 |
| Ask about homework | No Answer | 3.3 | 3.7 | 3.5 | 5.1 | 5.3 | 5.5 | 5.6 |
| | YES! | 69.7 | 61.4 | 53.1 | 46.7 | 40.7 | 36.4 | 29.9 |
| | yes | 21.4 | 25.2 | 30.1 | 31.6 | 33.8 | 33.6 | 34.0 |
| | no | 4.0 | 7.1 | 9.7 | 12.3 | 14.9 | 17.4 | 21.4 |
| | NO! | 1.6 | 2.6 | 3.6 | 4.2 | 5.3 | 7.1 | 9.1 |
| Parents know whereabouts | No Answer | 3.6 | 4.0 | 3.6 | 5.4 | 5.5 | 5.7 | 5.7 |
| | YES! | 76.8 | 70.7 | 64.2 | 57.1 | 54.1 | 50.8 | 44.7 |
| | yes | 14.4 | 18.5 | 23.7 | 28.2 | 30.8 | 32.4 | 34.9 |
| | no | 3.4 | 4.8 | 6.2 | 7.0 | 7.3 | 8.1 | 10.8 |
| | NO! | 1.8 | 2.1 | 2.3 | 2.3 | 2.4 | 3.0 | 3.8 |
| Clear rules about alcohol use | No Answer | 4.6 | 4.6 | 4.2 | 5.9 | 5.9 | 6.0 | 6.1 |
| | YES! | 80.8 | 76.4 | 68.7 | 61.0 | 57.7 | 55.0 | 49.9 |
| | yes | 9.5 | 12.2 | 17.6 | 22.1 | 24.4 | 25.8 | 28.8 |
| | no | 3.2 | 4.7 | 7.2 | 8.5 | 9.6 | 10.3 | 12.0 |
| | NO! | 2.0 | 2.0 | 2.4 | 2.5 | 2.5 | 2.9 | 3.2 |
| Parents ask before most family decisions made | No Answer | 7.1 | 6.2 | 5.3 | 6.9 | 6.8 | 6.9 | 6.8 |
| | YES! | 36.9 | 31.9 | 27.5 | 24.4 | 22.5 | 22.0 | 20.7 |
| | yes | 32.9 | 34.5 | 35.7 | 36.0 | 37.0 | 37.1 | 37.9 |
| | no | 13.4 | 16.4 | 19.8 | 21.4 | 22.3 | 22.7 | 23.7 |
| | NO! | 9.7 | 11.0 | 11.7 | 11.3 | 11.4 | 11.4 | 10.8 |
| Ask parents for help if problem | No Answer | 4.8 | 5.1 | 4.4 | 6.6 | 6.5 | 6.5 | 6.5 |
| | YES! | 65.7 | 55.9 | 47.9 | 42.4 | 40.5 | 40.7 | 41.1 |
| | yes | 19.4 | 23.7 | 27.7 | 29.7 | 31.6 | 32.5 | 33.8 |
| | no | 5.4 | 8.4 | 11.4 | 12.5 | 13.1 | 12.4 | 11.9 |
| | NO! | 4.8 | 6.9 | 8.6 | 8.8 | 8.3 | 7.9 | 6.7 |
| Parents give me chances for fun with them | No Answer | 5.2 | 5.5 | 4.8 | 7.0 | 6.8 | 6.7 | 6.8 |
| | YES! | 54.1 | 48.1 | 41.1 | 35.1 | 32.5 | 31.0 | 29.5 |
| | yes | 27.1 | 28.7 | 32.3 | 34.1 | 35.0 | 35.9 | 36.1 |
| | no | 9.5 | 12.3 | 15.3 | 17.0 | 18.7 | 19.0 | 20.3 |
| | NO! | 4.1 | 5.4 | 6.5 | 6.8 | 7.0 | 7.4 | 7.3 |

Family Management and Conflict by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Parents know when home late | No Answer | 6.0 | 6.3 | 5.3 | 7.3 | 7.2 | 7.2 | 7.4 |
| | YES! | 69.2 | 63.2 | 58.8 | 52.0 | 49.8 | 47.3 | 41.0 |
| | yes | 17.6 | 21.3 | 25.2 | 28.6 | 30.3 | 30.6 | 32.6 |
| | no | 5.0 | 6.7 | 8.0 | 9.2 | 9.8 | 11.2 | 13.9 |
| | NO! | 2.1 | 2.6 | 2.7 | 2.9 | 2.8 | 3.7 | 5.1 |
| Parents catch you drinking | No Answer | 6.0 | 5.9 | 5.3 | 7.5 | 7.3 | 7.2 | 7.1 |
| | YES! | 71.4 | 59.5 | 46.2 | 33.8 | 27.8 | 25.1 | 20.6 |
| | yes | 12.3 | 16.9 | 20.9 | 22.1 | 22.1 | 20.3 | 19.4 |
| | no | 6.3 | 11.8 | 19.0 | 26.2 | 31.1 | 33.7 | 37.5 |
| | NO! | 4.0 | 5.9 | 8.5 | 10.4 | 11.7 | 13.8 | 15.4 |
| Parents catch you with gun | No Answer | 6.2 | 6.0 | 5.3 | 7.6 | 7.3 | 7.4 | 7.2 |
| | YES! | 79.1 | 72.7 | 65.6 | 57.2 | 52.9 | 50.2 | 43.8 |
| | yes | 8.9 | 12.5 | 16.8 | 20.1 | 21.5 | 22.3 | 24.2 |
| | no | 3.1 | 5.5 | 8.2 | 10.4 | 12.9 | 13.9 | 17.5 |
| | NO! | 2.6 | 3.3 | 4.1 | 4.8 | 5.4 | 6.2 | 7.2 |
| Parents catch you skip school | No Answer | 5.3 | 5.6 | 4.8 | 7.1 | 6.7 | 6.8 | 6.9 |
| | YES! | 79.6 | 73.8 | 67.9 | 59.8 | 55.7 | 52.3 | 45.5 |
| | yes | 10.1 | 13.5 | 17.5 | 22.0 | 24.0 | 25.8 | 27.9 |
| | no | 3.0 | 4.3 | 6.6 | 7.6 | 9.6 | 10.5 | 13.8 |
| | NO! | 2.0 | 2.8 | 3.1 | 3.5 | 4.0 | 4.5 | 5.9 |
| Family argue repetitively | No Answer | 5.8 | 5.2 | 4.6 | 6.2 | 6.2 | 6.2 | 6.3 |
| | YES! | 17.5 | 20.7 | 22.9 | 23.7 | 23.3 | 22.3 | 20.5 |
| | yes | 19.6 | 21.4 | 23.6 | 25.4 | 26.5 | 25.6 | 25.3 |
| | no | 31.6 | 32.8 | 33.3 | 32.3 | 31.8 | 33.2 | 34.4 |
| | NO! | 25.4 | 19.9 | 15.7 | 12.4 | 12.2 | 12.7 | 13.5 |
| Family have serious arguments | No Answer | 5.9 | 5.6 | 4.9 | 6.5 | 6.5 | 6.5 | 6.7 |
| | YES! | 14.1 | 15.8 | 17.4 | 18.1 | 17.6 | 17.0 | 15.0 |
| | yes | 14.7 | 15.5 | 17.9 | 18.8 | 19.4 | 18.9 | 18.6 |
| | no | 29.5 | 32.1 | 34.2 | 34.8 | 35.4 | 35.7 | 37.4 |
| | NO! | 35.8 | 31.1 | 25.7 | 21.9 | 21.1 | 21.9 | 22.3 |
| Family insult each other | No Answer | 7.1 | 6.2 | 5.7 | 7.1 | 7.4 | 7.2 | 7.5 |
| | YES! | 14.7 | 16.5 | 18.3 | 18.6 | 17.8 | 16.7 | 14.6 |
| | yes | 16.6 | 17.6 | 19.6 | 20.2 | 20.3 | 19.3 | 19.0 |
| | no | 25.7 | 28.4 | 29.3 | 30.2 | 31.1 | 31.8 | 32.8 |
| | NO! | 35.9 | 31.3 | 27.1 | 23.8 | 23.4 | 24.9 | 26.1 |

**Interaction with Antisocial Peers
by Indiana 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Number of best friends suspended | No Answer | 5.0 | 5.3 | 4.8 | 7.3 | 7.0 | 6.8 | 6.9 |
| | None of my friends | 71.2 | 65.9 | 62.9 | 62.1 | 65.4 | 69.3 | 72.5 |
| | 1 of my friends | 14.7 | 16.4 | 17.5 | 16.7 | 15.4 | 13.9 | 12.0 |
| | 2 of my friends | 4.8 | 6.3 | 7.6 | 7.1 | 6.5 | 5.6 | 4.4 |
| | 3 of my friends | 1.6 | 2.2 | 2.7 | 2.3 | 2.2 | 1.7 | 1.4 |
| | 4 of my friends | 2.8 | 3.9 | 4.5 | 4.4 | 3.6 | 2.8 | 2.8 |
| Number of best friends carry guns | No Answer | 4.8 | 5.4 | 4.8 | 7.4 | 7.0 | 6.8 | 7.0 |
| | None of my friends | 91.6 | 89.6 | 88.9 | 85.5 | 84.9 | 84.9 | 83.4 |
| | 1 of my friends | 2.3 | 3.0 | 3.7 | 4.0 | 4.6 | 4.7 | 5.7 |
| | 2 of my friends | .6 | .8 | 1.1 | 1.3 | 1.5 | 1.6 | 1.8 |
| | 3 of my friends | .2 | .3 | .5 | .5 | .5 | .7 | .6 |
| | 4 of my friends | .5 | .9 | 1.1 | 1.3 | 1.5 | 1.4 | 1.5 |
| Number of best friends sold drugs | No Answer | 4.9 | 5.6 | 4.9 | 7.6 | 7.3 | 7.0 | 7.2 |
| | None of my friends | 91.6 | 86.5 | 81.7 | 73.2 | 70.7 | 69.7 | 68.9 |
| | 1 of my friends | 2.2 | 4.3 | 7.1 | 9.2 | 10.3 | 10.8 | 11.3 |
| | 2 of my friends | .6 | 1.5 | 2.8 | 4.5 | 5.4 | 5.8 | 6.1 |
| | 3 of my friends | .2 | .7 | 1.3 | 1.9 | 2.3 | 2.4 | 2.5 |
| | 4 of my friends | .4 | 1.5 | 2.1 | 3.7 | 4.0 | 4.3 | 4.0 |
| Number of best friends stolen vehicle | No Answer | 4.9 | 5.6 | 4.9 | 7.5 | 7.3 | 7.0 | 7.1 |
| | None of my friends | 92.6 | 90.5 | 89.3 | 85.2 | 85.4 | 87.1 | 87.7 |
| | 1 of my friends | 1.6 | 2.3 | 3.3 | 4.0 | 4.1 | 3.1 | 2.6 |
| | 2 of my friends | .4 | .6 | 1.0 | 1.3 | 1.4 | 1.2 | 1.1 |
| | 3 of my friends | .2 | .3 | .5 | .7 | .5 | .5 | .5 |
| | 4 of my friends | .4 | .7 | 1.0 | 1.3 | 1.2 | 1.1 | 1.0 |
| Number of best friends arrested | No Answer | 5.1 | 5.6 | 4.9 | 7.6 | 7.3 | 7.0 | 7.2 |
| | None of my friends | 87.3 | 83.1 | 79.3 | 74.9 | 74.3 | 74.6 | 74.1 |
| | 1 of my friends | 5.2 | 6.8 | 9.2 | 9.7 | 10.2 | 10.5 | 10.5 |
| | 2 of my friends | 1.2 | 2.0 | 3.0 | 3.6 | 4.0 | 3.8 | 4.1 |
| | 3 of my friends | .5 | .9 | 1.4 | 1.4 | 1.6 | 1.6 | 1.5 |
| | 4 of my friends | .7 | 1.5 | 2.1 | 2.7 | 2.6 | 2.4 | 2.6 |
| Number of best friends dropped out of school | No Answer | 5.2 | 5.7 | 5.1 | 7.7 | 7.3 | 7.0 | 7.1 |
| | None of my friends | 90.5 | 88.6 | 87.8 | 82.1 | 79.9 | 78.7 | 78.9 |
| | 1 of my friends | 3.2 | 3.6 | 4.3 | 6.2 | 8.0 | 8.7 | 8.3 |
| | 2 of my friends | .6 | .9 | 1.2 | 1.7 | 2.2 | 2.9 | 2.6 |
| | 3 of my friends | .2 | .4 | .5 | .8 | 1.0 | 1.1 | 1.3 |
| | 4 of my friends | .4 | .8 | 1.1 | 1.4 | 1.5 | 1.6 | 1.7 |
| Number of best friends been members of gangs | No Answer | 6.1 | 6.4 | 5.7 | 8.3 | 8.1 | 7.8 | 7.8 |
| | None of my friends | 86.0 | 83.1 | 81.5 | 80.0 | 81.1 | 82.9 | 84.0 |
| | 1 of my friends | 3.9 | 4.7 | 5.8 | 4.9 | 4.4 | 4.0 | 3.3 |
| | 2 of my friends | 1.4 | 1.9 | 2.4 | 2.1 | 2.2 | 1.8 | 1.8 |
| | 3 of my friends | .8 | .9 | 1.3 | 1.3 | 1.0 | .9 | .8 |
| | 4 of my friends | 1.8 | 2.9 | 3.3 | 3.3 | 3.1 | 2.6 | 2.3 |

Interaction with Prosocial Peers by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Number of best friends participated in school activities | No Answer | 7.5 | 7.2 | 6.7 | 8.7 | 8.6 | 7.9 | 8.3 |
| | None of my friends | 21.3 | 21.6 | 21.5 | 21.0 | 20.5 | 20.9 | 20.6 |
| | 1 of my friends | 10.5 | 9.2 | 9.1 | 8.4 | 8.6 | 8.5 | 8.5 |
| | 2 of my friends | 14.8 | 13.9 | 13.6 | 13.2 | 13.7 | 13.4 | 12.8 |
| | 3 of my friends | 11.8 | 12.1 | 12.9 | 12.1 | 12.2 | 11.5 | 11.0 |
| | 4 of my friends | 34.2 | 36.1 | 36.2 | 36.7 | 36.4 | 37.8 | 38.8 |
| | | | | | | | | |
| Number of best friends commitment to stay drug-free | No Answer | 8.8 | 8.1 | 7.4 | 9.5 | 9.0 | 8.5 | 8.6 |
| | None of my friends | 18.9 | 22.4 | 27.5 | 30.2 | 31.3 | 34.9 | 36.3 |
| | 1 of my friends | 4.1 | 5.4 | 7.9 | 9.5 | 11.4 | 11.8 | 11.8 |
| | 2 of my friends | 4.4 | 5.6 | 7.8 | 9.2 | 10.0 | 9.9 | 10.3 |
| | 3 of my friends | 5.7 | 6.6 | 8.2 | 8.7 | 9.0 | 8.4 | 8.3 |
| | 4 of my friends | 58.0 | 51.9 | 41.1 | 32.8 | 29.3 | 26.5 | 24.8 |
| | | | | | | | | |
| Number of best friends liked school | No Answer | 7.9 | 7.6 | 7.0 | 9.3 | 8.8 | 8.4 | 8.5 |
| | None of my friends | 25.5 | 29.1 | 32.4 | 34.4 | 35.3 | 36.1 | 35.5 |
| | 1 of my friends | 9.4 | 10.2 | 10.4 | 10.3 | 11.2 | 10.8 | 10.7 |
| | 2 of my friends | 13.5 | 13.3 | 13.9 | 13.7 | 14.3 | 14.2 | 14.4 |
| | 3 of my friends | 12.3 | 12.1 | 11.9 | 11.1 | 10.8 | 10.2 | 10.3 |
| | 4 of my friends | 31.4 | 27.7 | 24.3 | 21.2 | 19.6 | 20.2 | 20.6 |
| | | | | | | | | |
| Number of best friends regularly attended religious activities | No Answer | 9.9 | 8.8 | 7.5 | 9.7 | 8.9 | 8.4 | 8.4 |
| | None of my friends | 21.9 | 22.9 | 25.3 | 26.0 | 27.3 | 29.2 | 30.0 |
| | 1 of my friends | 14.4 | 14.8 | 16.5 | 17.1 | 17.5 | 17.8 | 18.0 |
| | 2 of my friends | 17.5 | 17.6 | 18.6 | 17.3 | 17.8 | 17.3 | 18.0 |
| | 3 of my friends | 13.8 | 14.1 | 13.0 | 12.2 | 11.9 | 10.9 | 10.2 |
| | 4 of my friends | 22.5 | 21.9 | 19.1 | 17.7 | 16.6 | 16.4 | 15.3 |
| | | | | | | | | |
| Number of best friends tried to do well in school | No Answer | 8.2 | 7.6 | 7.3 | 9.1 | 8.9 | 8.2 | 8.6 |
| | None of my friends | 7.6 | 9.4 | 10.0 | 10.4 | 10.3 | 11.0 | 10.2 |
| | 1 of my friends | 4.1 | 4.6 | 5.8 | 6.1 | 5.6 | 5.7 | 5.8 |
| | 2 of my friends | 5.6 | 6.8 | 8.4 | 9.0 | 10.0 | 10.1 | 10.4 |
| | 3 of my friends | 9.3 | 11.7 | 13.5 | 13.5 | 14.4 | 14.3 | 14.0 |
| | 4 of my friends | 65.3 | 59.9 | 55.0 | 51.9 | 50.7 | 50.6 | 51.0 |
| | | | | | | | | |

Rewards for Antisocial Involvement by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|--------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cool if smoke cigarettes | No Answer | 6.5 | 6.8 | 5.8 | 8.8 | 8.2 | 7.8 | 7.8 |
| | No or very little chance | 77.7 | 71.4 | 66.9 | 61.2 | 61.2 | 62.0 | 61.5 |
| | Little chance | 6.8 | 9.5 | 12.2 | 13.2 | 14.4 | 14.2 | 15.0 |
| | Some chance | 4.2 | 6.5 | 8.8 | 10.2 | 10.3 | 9.8 | 10.3 |
| | Pretty good chance | 2.0 | 2.8 | 3.5 | 3.9 | 3.4 | 3.3 | 2.8 |
| | Very good chance | 2.7 | 2.9 | 2.8 | 2.7 | 2.6 | 2.8 | 2.6 |
| | | | | | | | | |
| Cool if drink alcohol regularly | No Answer | 6.6 | 7.0 | 5.9 | 9.0 | 8.4 | 8.0 | 8.1 |
| | No or very little chance | 77.5 | 69.7 | 62.4 | 52.8 | 48.9 | 49.1 | 46.7 |
| | Little chance | 7.0 | 9.7 | 12.5 | 13.5 | 14.3 | 14.0 | 14.2 |
| | Some chance | 3.8 | 6.8 | 10.1 | 12.6 | 14.5 | 14.1 | 15.3 |
| | Pretty good chance | 2.2 | 3.5 | 5.3 | 7.7 | 9.1 | 9.6 | 10.1 |
| | Very good chance | 2.9 | 3.3 | 3.8 | 4.4 | 4.8 | 5.3 | 5.6 |
| | | | | | | | | |
| Cool if smoke marijuana | No Answer | 6.7 | 7.1 | 5.9 | 8.9 | 8.4 | 8.0 | 8.0 |
| | No or very little chance | 81.0 | 73.8 | 66.1 | 56.6 | 53.3 | 54.1 | 53.0 |
| | Little chance | 4.2 | 6.5 | 9.1 | 10.4 | 12.0 | 11.8 | 13.4 |
| | Some chance | 2.6 | 4.7 | 7.9 | 10.9 | 12.0 | 12.2 | 13.1 |
| | Pretty good chance | 1.8 | 3.4 | 5.3 | 7.2 | 7.8 | 7.7 | 6.9 |
| | Very good chance | 3.7 | 4.5 | 5.5 | 6.0 | 6.4 | 6.2 | 5.5 |
| | | | | | | | | |
| Cool if carry a gun | No Answer | 6.8 | 7.2 | 6.1 | 9.0 | 8.5 | 8.1 | 8.1 |
| | No or very little chance | 78.0 | 75.3 | 74.2 | 71.1 | 72.5 | 73.9 | 72.4 |
| | Little chance | 5.6 | 6.8 | 8.2 | 8.7 | 8.4 | 8.1 | 9.1 |
| | Some chance | 3.4 | 4.1 | 4.9 | 5.4 | 5.0 | 4.7 | 5.2 |
| | Pretty good chance | 1.9 | 2.5 | 2.7 | 2.2 | 2.1 | 1.8 | 2.0 |
| | Very good chance | 4.2 | 4.1 | 4.0 | 3.6 | 3.4 | 3.4 | 3.2 |
| | | | | | | | | |

School Commitment
by Indiana 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|----------------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Days of school skipped during last month | No Answer | 5.1 | 5.5 | 5.0 | 7.3 | 7.0 | 6.4 | 6.6 |
| | None | 87.7 | 85.1 | 82.6 | 79.6 | 78.4 | 76.8 | 70.9 |
| | 1 | 3.4 | 4.1 | 5.6 | 5.8 | 6.6 | 7.3 | 9.4 |
| | 2 | 1.4 | 2.1 | 2.7 | 2.6 | 3.1 | 3.7 | 5.3 |
| | 3 | 1.0 | 1.2 | 1.6 | 1.9 | 1.8 | 2.3 | 3.3 |
| | 4-5 | .7 | .9 | 1.4 | 1.5 | 1.6 | 1.8 | 2.5 |
| | 6-10 | .3 | .4 | .5 | .5 | .6 | .8 | 1.0 |
| | 11 or more | .4 | .6 | .6 | .8 | .8 | .9 | 1.0 |
| Feel schoolwork is meaningful | No Answer | 5.4 | 6.1 | 5.0 | 7.7 | 7.1 | 6.7 | 6.8 |
| | Almost always | 42.6 | 32.3 | 27.0 | 21.8 | 19.0 | 18.6 | 15.9 |
| | Often | 20.7 | 22.5 | 23.1 | 21.6 | 20.7 | 20.2 | 19.1 |
| | Sometimes | 19.5 | 22.5 | 25.6 | 27.3 | 29.6 | 29.3 | 30.6 |
| | Seldom | 5.5 | 9.3 | 11.8 | 13.9 | 15.6 | 17.4 | 19.0 |
| | Never | 6.3 | 7.4 | 7.5 | 7.7 | 8.0 | 7.8 | 8.6 |
| How interesting are courses | No Answer | 9.0 | 7.6 | 5.9 | 8.3 | 7.6 | 7.2 | 7.1 |
| | Very interesting and stimulating | 19.3 | 13.1 | 10.2 | 8.5 | 7.7 | 8.4 | 8.2 |
| | Quite interesting | 26.2 | 23.2 | 21.0 | 20.3 | 19.2 | 19.5 | 18.7 |
| | Fairly interesting | 27.2 | 31.1 | 34.3 | 35.2 | 35.5 | 35.7 | 35.1 |
| | Slightly dull | 10.5 | 13.6 | 16.2 | 16.0 | 17.2 | 16.8 | 17.2 |
| | Very dull | 7.7 | 11.3 | 12.4 | 11.8 | 12.8 | 12.5 | 13.9 |
| Importance of school to later life | No Answer | 6.6 | 7.0 | 5.8 | 8.5 | 7.8 | 7.4 | 7.3 |
| | Very important | 62.0 | 49.6 | 42.8 | 32.4 | 27.4 | 25.3 | 21.9 |
| | Quite important | 15.5 | 20.0 | 20.7 | 21.6 | 21.3 | 20.2 | 20.1 |
| | Fairly important | 8.6 | 12.5 | 16.0 | 19.3 | 20.7 | 22.0 | 23.3 |
| | Slightly important | 5.1 | 7.7 | 10.9 | 13.4 | 16.9 | 19.0 | 20.4 |
| | Not at all important | 2.1 | 3.2 | 3.8 | 4.9 | 6.0 | 6.1 | 7.1 |
| Grades better than other students | No Answer | 6.7 | 7.3 | 5.8 | 8.8 | 8.1 | 7.6 | 7.3 |
| | YES! | 26.2 | 25.1 | 24.5 | 23.2 | 22.8 | 23.8 | 25.7 |
| | yes | 44.8 | 43.7 | 44.6 | 42.9 | 44.4 | 44.9 | 44.6 |
| | no | 17.4 | 18.4 | 19.9 | 19.8 | 20.0 | 19.4 | 18.7 |
| | NO! | 4.9 | 5.6 | 5.2 | 5.3 | 4.8 | 4.3 | 3.6 |
| How often did you enjoy school | No Answer | 6.2 | 6.8 | 5.5 | 8.5 | 8.0 | 7.5 | 7.4 |
| | Never | 8.0 | 9.2 | 9.5 | 10.1 | 10.5 | 10.5 | 10.2 |
| | Seldom | 8.6 | 11.1 | 12.5 | 13.9 | 15.7 | 17.2 | 16.8 |
| | Sometimes | 23.6 | 25.1 | 28.0 | 29.2 | 30.7 | 31.4 | 31.6 |
| | Often | 25.8 | 26.9 | 26.9 | 25.4 | 24.3 | 23.7 | 24.2 |
| | A lot | 27.8 | 20.9 | 17.6 | 12.9 | 10.8 | 9.6 | 9.8 |

School Commitment
by Indiana 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| How often did you hate school | No Answer | 6.4 | 6.9 | 5.7 | 8.6 | 8.1 | 7.6 | 7.3 |
| | Never | 22.5 | 15.2 | 11.7 | 8.7 | 7.3 | 6.7 | 6.6 |
| | Seldom | 25.7 | 27.1 | 26.3 | 23.4 | 22.2 | 20.8 | 21.2 |
| | Sometimes | 23.4 | 24.9 | 27.9 | 29.1 | 29.9 | 31.1 | 31.6 |
| | Often | 10.2 | 12.2 | 14.1 | 16.0 | 17.5 | 18.4 | 18.9 |
| | A lot | 11.8 | 13.7 | 14.3 | 14.2 | 15.1 | 15.3 | 14.3 |
| How often did you do your best work | No Answer | 6.4 | 7.0 | 5.7 | 8.8 | 8.2 | 7.6 | 7.3 |
| | Never | 1.6 | 2.3 | 2.3 | 2.4 | 2.7 | 2.4 | 2.5 |
| | Seldom | 2.3 | 3.6 | 4.6 | 4.8 | 5.0 | 5.2 | 5.7 |
| | Sometimes | 7.5 | 10.6 | 14.3 | 16.3 | 18.0 | 18.9 | 19.8 |
| | Often | 17.2 | 21.9 | 26.2 | 27.6 | 29.0 | 29.5 | 31.6 |
| | A lot | 65.1 | 54.6 | 46.9 | 40.1 | 37.2 | 36.4 | 33.0 |

Opportunities for Prosocial Involvement by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Students have chances to help decide activities and rules | No Answer | 9.1 | 9.3 | 7.5 | 11.4 | 10.6 | 9.9 | 9.5 |
| | YES! | 28.5 | 23.1 | 20.3 | 20.6 | 18.7 | 18.8 | 16.7 |
| | yes | 33.2 | 32.3 | 33.0 | 32.7 | 31.8 | 31.7 | 32.3 |
| | no | 19.3 | 23.6 | 26.9 | 25.0 | 27.2 | 27.3 | 28.2 |
| | NO! | 9.9 | 11.8 | 12.4 | 10.2 | 11.8 | 12.2 | 13.3 |
| Teachers ask to work on special class projects | No Answer | 9.5 | 9.5 | 7.7 | 11.7 | 10.8 | 10.2 | 9.7 |
| | YES! | 23.5 | 18.6 | 15.6 | 12.1 | 11.1 | 10.9 | 10.9 |
| | yes | 31.3 | 30.3 | 31.1 | 28.1 | 28.9 | 29.2 | 31.1 |
| | no | 24.7 | 29.5 | 33.1 | 36.5 | 36.9 | 37.4 | 36.7 |
| | NO! | 11.0 | 12.1 | 12.5 | 11.6 | 12.3 | 12.3 | 11.7 |
| Chances for kids to get involved in activities outside of class | No Answer | 9.5 | 9.9 | 8.0 | 12.1 | 11.1 | 10.4 | 10.1 |
| | YES! | 66.8 | 64.9 | 60.4 | 54.9 | 51.8 | 50.4 | 48.2 |
| | yes | 18.3 | 19.5 | 25.1 | 25.7 | 29.2 | 31.0 | 32.7 |
| | no | 3.5 | 3.7 | 4.3 | 5.0 | 5.6 | 5.6 | 6.3 |
| | NO! | 1.9 | 2.0 | 2.2 | 2.2 | 2.4 | 2.6 | 2.6 |
| Chances for kids to talk with teacher one-on-one | No Answer | 10.1 | 10.3 | 8.5 | 12.5 | 11.5 | 10.9 | 10.4 |
| | YES! | 49.9 | 43.2 | 39.4 | 32.9 | 30.3 | 30.6 | 30.6 |
| | yes | 27.8 | 30.6 | 33.7 | 36.1 | 38.1 | 39.3 | 41.0 |
| | no | 8.8 | 11.4 | 13.6 | 13.9 | 15.4 | 14.7 | 13.7 |
| | NO! | 3.4 | 4.5 | 4.8 | 4.6 | 4.6 | 4.5 | 4.4 |
| Chances to be a part of class discussion and activities | No Answer | 11.2 | 11.1 | 9.2 | 13.2 | 11.9 | 11.2 | 10.8 |
| | YES! | 45.8 | 39.8 | 35.1 | 28.2 | 25.4 | 25.7 | 25.6 |
| | yes | 30.3 | 34.1 | 39.6 | 42.2 | 45.3 | 46.8 | 48.5 |
| | no | 9.0 | 10.5 | 11.8 | 12.4 | 13.1 | 12.2 | 11.5 |
| | NO! | 3.6 | 4.4 | 4.4 | 4.0 | 4.3 | 4.0 | 3.7 |

Rewards for Prosocial Involvement by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Teachers notice when doing a good job and let me know | No Answer | 9.9 | 10.1 | 8.3 | 12.2 | 11.4 | 10.6 | 10.1 |
| | YES! | 42.0 | 33.0 | 28.1 | 21.2 | 18.4 | 18.2 | 17.1 |
| | yes | 31.4 | 34.4 | 37.9 | 37.8 | 39.2 | 40.2 | 41.7 |
| | no | 11.5 | 15.4 | 18.2 | 20.9 | 22.7 | 23.0 | 22.8 |
| | NO! | 5.3 | 7.2 | 7.5 | 7.9 | 8.3 | 8.0 | 8.2 |
| Feel safe at school | No Answer | 9.7 | 10.0 | 8.2 | 12.3 | 11.3 | 10.7 | 10.1 |
| | YES! | 55.7 | 46.0 | 39.8 | 30.6 | 30.2 | 32.2 | 33.5 |
| | yes | 24.7 | 30.9 | 36.6 | 40.6 | 42.0 | 42.4 | 43.0 |
| | no | 6.0 | 8.2 | 10.0 | 11.2 | 11.0 | 10.1 | 9.1 |
| | NO! | 3.8 | 4.9 | 5.4 | 5.3 | 5.5 | 4.6 | 4.2 |
| School lets parents know when do good | No Answer | 10.4 | 10.6 | 8.8 | 12.6 | 11.6 | 10.9 | 10.4 |
| | YES! | 28.4 | 23.8 | 19.6 | 14.7 | 13.0 | 12.7 | 11.9 |
| | yes | 26.3 | 24.6 | 24.8 | 22.0 | 21.8 | 22.2 | 22.4 |
| | no | 23.0 | 26.2 | 29.7 | 32.8 | 34.5 | 34.9 | 34.9 |
| | NO! | 11.9 | 14.8 | 17.2 | 17.9 | 19.1 | 19.4 | 20.4 |
| Teachers praise when do well in school | No Answer | 11.8 | 11.4 | 9.4 | 13.3 | 12.0 | 11.4 | 10.9 |
| | YES! | 28.9 | 24.6 | 21.5 | 16.5 | 14.6 | 14.6 | 13.6 |
| | yes | 30.5 | 30.3 | 31.6 | 30.9 | 31.1 | 32.2 | 34.4 |
| | no | 19.2 | 22.6 | 25.9 | 27.5 | 29.9 | 30.1 | 29.6 |
| | NO! | 9.6 | 11.2 | 11.6 | 11.8 | 12.4 | 11.7 | 11.5 |

Participation in Afterschool Activities by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Participation in ROCK, LEAD, SADD, etc. | No Answer | 14.8 | 15.1 | 12.5 | 16.3 | 15.0 | 13.9 | 13.5 |
| | Yes | 18.2 | 12.5 | 9.0 | 7.6 | 7.3 | 7.9 | 7.7 |
| | No | 66.9 | 72.4 | 78.5 | 76.1 | 77.7 | 78.1 | 78.8 |

Favorable Attitudes Toward Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Wrong to take a gun to school | No Answer | 10.7 | 11.2 | 8.3 | 12.3 | 10.9 | 10.1 | 9.9 |
| | Very wrong | 83.4 | 80.2 | 81.4 | 76.5 | 78.6 | 79.9 | 80.1 |
| | Wrong | 4.0 | 5.7 | 6.9 | 7.4 | 6.8 | 6.4 | 6.3 |
| | A little bit wrong | 1.2 | 1.8 | 2.2 | 2.2 | 2.1 | 1.9 | 2.1 |
| | Not at all wrong | .7 | 1.1 | 1.3 | 1.6 | 1.6 | 1.7 | 1.6 |
| Wrong to steal something more than \$5 | No Answer | 10.8 | 11.4 | 8.4 | 12.5 | 11.2 | 10.4 | 10.1 |
| | Very wrong | 65.6 | 59.6 | 56.2 | 50.2 | 52.2 | 55.1 | 56.8 |
| | Wrong | 17.4 | 19.6 | 23.3 | 24.3 | 24.4 | 23.4 | 23.8 |
| | A little bit wrong | 4.9 | 7.1 | 9.4 | 10.3 | 9.7 | 8.7 | 7.2 |
| | Not at all wrong | 1.3 | 2.2 | 2.7 | 2.7 | 2.5 | 2.4 | 2.0 |
| Wrong to attack someone | No Answer | 11.0 | 11.6 | 8.6 | 12.8 | 11.4 | 10.5 | 10.2 |
| | Very wrong | 76.3 | 70.5 | 68.0 | 61.9 | 63.4 | 65.3 | 67.1 |
| | Wrong | 8.1 | 11.4 | 14.7 | 16.1 | 16.7 | 16.0 | 15.5 |
| | A little bit wrong | 3.1 | 4.4 | 6.2 | 6.5 | 6.1 | 5.8 | 5.1 |
| | Not at all wrong | 1.5 | 2.2 | 2.5 | 2.7 | 2.5 | 2.3 | 2.0 |
| Wrong to pick a fight | No Answer | 11.1 | 11.8 | 8.7 | 12.9 | 11.4 | 10.7 | 10.3 |
| | Very wrong | 59.5 | 52.1 | 46.5 | 41.8 | 42.3 | 45.5 | 46.3 |
| | Wrong | 18.9 | 21.1 | 24.6 | 25.6 | 27.6 | 26.6 | 28.4 |
| | A little bit wrong | 7.7 | 10.8 | 14.9 | 14.9 | 14.3 | 13.4 | 11.8 |
| | Not at all wrong | 2.8 | 4.2 | 5.3 | 4.8 | 4.4 | 3.8 | 3.2 |
| Wrong to skip school | No Answer | 11.3 | 11.8 | 8.8 | 13.0 | 11.6 | 10.8 | 10.4 |
| | Very wrong | 75.1 | 66.4 | 60.5 | 49.8 | 47.8 | 46.4 | 43.6 |
| | Wrong | 9.5 | 14.4 | 19.8 | 22.8 | 25.0 | 25.8 | 26.5 |
| | A little bit wrong | 2.8 | 5.1 | 7.9 | 10.8 | 12.0 | 13.1 | 15.4 |
| | Not at all wrong | 1.4 | 2.3 | 3.0 | 3.6 | 3.6 | 3.9 | 4.0 |
| Wrong to drink alcohol regularly | No Answer | 11.2 | 11.8 | 8.9 | 13.1 | 11.6 | 10.9 | 10.4 |
| | Very wrong | 78.3 | 70.2 | 61.9 | 48.6 | 44.7 | 42.3 | 37.7 |
| | Wrong | 6.2 | 9.5 | 14.6 | 17.4 | 18.9 | 19.5 | 20.0 |
| | A little bit wrong | 2.9 | 5.6 | 10.1 | 14.4 | 17.2 | 18.6 | 21.6 |
| | Not at all wrong | 1.4 | 2.9 | 4.5 | 6.5 | 7.6 | 8.6 | 10.3 |
| Wrong to smoke cigarettes | No Answer | 11.4 | 11.9 | 8.9 | 13.2 | 11.7 | 10.9 | 10.5 |
| | Very wrong | 79.4 | 73.1 | 66.7 | 55.4 | 52.2 | 48.1 | 40.1 |
| | Wrong | 5.8 | 8.2 | 13.0 | 15.6 | 17.4 | 17.5 | 18.2 |
| | A little bit wrong | 2.0 | 4.0 | 7.4 | 9.9 | 11.5 | 14.0 | 16.4 |
| | Not at all wrong | 1.3 | 2.8 | 4.1 | 5.9 | 7.2 | 9.5 | 14.7 |

Favorable Attitudes Toward Drug Use by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Wrong to smoke marijuana | No Answer | 11.5 | 12.0 | 9.0 | 13.2 | 11.6 | 10.9 | 10.5 |
| | Very wrong | 82.8 | 76.2 | 69.4 | 56.4 | 51.7 | 48.5 | 44.7 |
| | Wrong | 3.0 | 5.4 | 9.5 | 12.6 | 14.4 | 15.1 | 16.3 |
| | A little bit wrong | 1.4 | 3.4 | 6.8 | 9.6 | 12.0 | 13.5 | 15.2 |
| | Not at all wrong | 1.4 | 2.9 | 5.3 | 8.3 | 10.3 | 11.9 | 13.3 |
| Wrong to use illegal drugs | No Answer | 11.7 | 12.2 | 9.1 | 13.3 | 11.9 | 11.0 | 10.6 |
| | Very wrong | 84.6 | 80.9 | 80.0 | 72.1 | 72.1 | 72.0 | 71.2 |
| | Wrong | 2.2 | 3.9 | 6.4 | 8.6 | 9.7 | 9.6 | 10.6 |
| | A little bit wrong | .7 | 1.5 | 2.6 | 3.4 | 3.7 | 4.4 | 4.4 |
| | Not at all wrong | .8 | 1.5 | 1.9 | 2.6 | 2.7 | 3.1 | 3.2 |

Rebelliousness by Indiana 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|----------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Like to see what can get away with | No Answer | 13.1 | 13.1 | 10.1 | 14.3 | 12.6 | 11.6 | 11.3 |
| | Very false | 60.0 | 51.2 | 44.8 | 39.3 | 38.2 | 38.2 | 38.3 |
| | Somewhat false | 14.8 | 18.3 | 21.5 | 21.1 | 23.0 | 22.6 | 23.1 |
| | Somewhat true | 9.0 | 13.1 | 17.7 | 19.0 | 20.0 | 21.3 | 21.3 |
| | Very true | 3.0 | 4.3 | 6.0 | 6.3 | 6.2 | 6.3 | 6.0 |
| Ignore rules | No Answer | 13.2 | 13.3 | 10.2 | 14.5 | 12.7 | 11.7 | 11.5 |
| | Very false | 62.7 | 55.8 | 50.6 | 44.2 | 43.8 | 44.1 | 42.3 |
| | Somewhat false | 14.5 | 17.3 | 20.6 | 21.9 | 23.2 | 23.7 | 25.0 |
| | Somewhat true | 6.8 | 9.6 | 13.4 | 14.4 | 15.3 | 15.6 | 16.3 |
| | Very true | 2.8 | 4.0 | 5.2 | 5.0 | 5.0 | 4.8 | 5.0 |
| Do opposite of what I am told | No Answer | 13.3 | 13.4 | 10.4 | 14.6 | 12.8 | 11.8 | 11.5 |
| | Very false | 63.6 | 58.2 | 55.0 | 51.2 | 53.3 | 55.7 | 56.4 |
| | Somewhat false | 13.6 | 16.1 | 19.3 | 19.2 | 20.1 | 19.6 | 19.6 |
| | Somewhat true | 6.8 | 8.7 | 11.0 | 10.8 | 10.1 | 9.5 | 9.3 |
| | Very true | 2.7 | 3.5 | 4.3 | 4.2 | 3.8 | 3.4 | 3.2 |

**Laws and Norms Favorable to Drug Use
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Caught by police alcohol | No Answer | 14.5 | 14.2 | 11.1 | 15.5 | 13.7 | 12.8 | 12.6 |
| | YES! | 27.2 | 21.4 | 15.3 | 10.4 | 8.7 | 8.1 | 7.7 |
| | yes | 27.0 | 24.9 | 21.8 | 17.9 | 15.4 | 14.5 | 13.9 |
| | no | 21.2 | 27.3 | 36.2 | 39.1 | 43.5 | 44.8 | 45.7 |
| | NO! | 10.1 | 12.3 | 15.7 | 17.1 | 18.6 | 19.8 | 20.1 |
| Caught by police marijuana | No Answer | 14.6 | 14.3 | 11.2 | 15.7 | 13.9 | 13.0 | 12.7 |
| | YES! | 37.7 | 29.0 | 20.4 | 12.8 | 10.1 | 8.8 | 8.1 |
| | yes | 25.2 | 25.5 | 25.0 | 20.3 | 17.6 | 16.6 | 15.8 |
| | no | 14.4 | 20.6 | 29.3 | 34.7 | 39.9 | 42.0 | 43.5 |
| | NO! | 8.2 | 10.6 | 14.1 | 16.4 | 18.5 | 19.6 | 20.0 |
| Caught by police gun | No Answer | 14.6 | 14.3 | 11.2 | 15.7 | 13.9 | 13.0 | 12.7 |
| | YES! | 41.9 | 34.2 | 27.9 | 20.3 | 17.6 | 16.4 | 14.8 |
| | yes | 22.9 | 25.2 | 27.4 | 26.7 | 26.7 | 26.8 | 26.6 |
| | no | 13.2 | 17.3 | 22.7 | 25.8 | 29.3 | 31.3 | 32.5 |
| | NO! | 7.5 | 9.0 | 10.8 | 11.5 | 12.5 | 12.5 | 13.3 |
| Caught by police smoke cigarettes | No Answer | 15.1 | 14.6 | 11.6 | 16.2 | 14.3 | 13.1 | 13.1 |
| | YES! | 24.5 | 19.3 | 13.9 | 9.4 | 7.7 | 7.2 | 6.2 |
| | yes | 22.5 | 20.6 | 17.8 | 14.5 | 12.0 | 10.8 | 9.6 |
| | no | 23.7 | 28.5 | 35.1 | 36.2 | 39.9 | 40.3 | 40.8 |
| | NO! | 14.2 | 17.0 | 21.6 | 23.7 | 26.1 | 28.6 | 30.3 |
| Neighborhood feel about marijuana | No Answer | 13.6 | 13.8 | 10.5 | 15.5 | 13.6 | 12.6 | 12.3 |
| | Very wrong | 75.6 | 69.6 | 63.7 | 51.7 | 48.3 | 46.5 | 44.4 |
| | Wrong | 7.4 | 11.0 | 16.1 | 19.8 | 22.6 | 24.0 | 25.5 |
| | A little bit wrong | 2.3 | 3.9 | 7.0 | 9.3 | 11.1 | 12.4 | 12.8 |
| | Not at all wrong | 1.1 | 1.8 | 2.8 | 3.7 | 4.4 | 4.5 | 5.1 |
| Neighborhood feel about alcohol | No Answer | 13.7 | 13.8 | 10.7 | 15.6 | 13.7 | 12.7 | 12.6 |
| | Very wrong | 67.3 | 60.3 | 53.3 | 41.3 | 37.2 | 34.4 | 30.5 |
| | Wrong | 11.9 | 15.1 | 20.3 | 23.0 | 25.4 | 25.9 | 25.9 |
| | A little bit wrong | 5.0 | 7.6 | 11.3 | 14.3 | 17.1 | 19.4 | 22.2 |
| | Not at all wrong | 2.0 | 3.2 | 4.4 | 5.8 | 6.6 | 7.6 | 8.8 |
| Neighborhood feel about cigarettes | No Answer | 13.9 | 14.0 | 10.8 | 15.8 | 13.8 | 12.8 | 12.6 |
| | Very wrong | 68.2 | 62.2 | 56.0 | 44.8 | 40.5 | 36.6 | 28.9 |
| | Wrong | 11.2 | 14.3 | 19.2 | 22.1 | 24.6 | 24.8 | 24.8 |
| | A little bit wrong | 4.6 | 6.4 | 9.5 | 11.6 | 14.5 | 17.4 | 20.6 |
| | Not at all wrong | 2.1 | 3.1 | 4.4 | 5.7 | 6.6 | 8.3 | 13.1 |

**Community Rewards for Prosocial Involvement
by Indiana 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Neighbors notice good job and let me know | No Answer | 14.9 | 14.3 | 11.6 | 16.2 | 14.4 | 13.3 | 13.1 |
| | YES! | 20.9 | 16.9 | 13.4 | 10.9 | 9.3 | 8.9 | 8.4 |
| | yes | 21.0 | 19.9 | 18.4 | 16.6 | 15.7 | 15.6 | 15.4 |
| | no | 23.1 | 26.1 | 29.0 | 29.2 | 30.6 | 31.0 | 31.0 |
| | NO! | 20.0 | 22.7 | 27.5 | 27.1 | 30.1 | 31.2 | 32.1 |
| | | | | | | | | |
| People in neighborhood who are proud of me | No Answer | 14.9 | 14.4 | 11.8 | 16.3 | 14.5 | 13.4 | 13.1 |
| | YES! | 25.3 | 20.5 | 16.4 | 13.0 | 11.7 | 10.9 | 10.9 |
| | yes | 24.0 | 24.7 | 24.1 | 22.5 | 22.6 | 23.5 | 24.2 |
| | no | 18.8 | 20.5 | 23.7 | 24.1 | 24.8 | 25.0 | 24.2 |
| | NO! | 16.9 | 19.8 | 24.0 | 24.1 | 26.4 | 27.3 | 27.6 |
| | | | | | | | | |
| People in neighborhood who encourage to do my best | No Answer | 15.0 | 14.6 | 12.0 | 16.5 | 14.7 | 13.5 | 13.4 |
| | YES! | 31.2 | 25.3 | 19.6 | 15.3 | 13.3 | 12.6 | 11.9 |
| | yes | 23.2 | 23.6 | 24.4 | 23.0 | 23.0 | 23.6 | 23.9 |
| | no | 14.9 | 17.8 | 21.0 | 21.9 | 23.4 | 23.6 | 23.8 |
| | NO! | 15.7 | 18.7 | 23.0 | 23.3 | 25.6 | 26.8 | 27.0 |
| | | | | | | | | |

**Personal and Family Information
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------|----------------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Hispanic | No Answer | 7.7 | 6.3 | 6.2 | 6.5 | 6.8 | 6.2 | 5.9 |
| | Hispanic | 8.7 | 9.2 | 9.2 | 8.7 | 7.6 | 8.2 | 7.1 |
| | Non-Hispanic | 83.7 | 84.5 | 84.6 | 84.8 | 85.6 | 85.6 | 87.1 |
| Race | No Answer | 3.5 | 4.0 | 4.1 | 4.1 | 3.7 | 3.1 | 3.1 |
| | White | 78.4 | 79.5 | 77.9 | 78.5 | 79.2 | 80.5 | 82.3 |
| | Black/African American | 6.4 | 5.4 | 6.5 | 6.0 | 7.2 | 6.3 | 5.8 |
| | Asian | 1.4 | 1.6 | 1.6 | 1.7 | 1.6 | 1.9 | 1.7 |
| | Native Hawaiian/Pacific Islander | .2 | .1 | .3 | .5 | .4 | .5 | .5 |
| | American Indian/Alaskan Native | .5 | .7 | .7 | .8 | .7 | .7 | .6 |
| | More than one race | 5.9 | 5.7 | 6.3 | 6.0 | 5.1 | 4.8 | 4.3 |
| | Race not known or other | 3.7 | 3.1 | 2.4 | 2.5 | 2.2 | 2.1 | 1.8 |
| Age | No Answer | 1.2 | 1.5 | 1.3 | 1.3 | 1.8 | 1.5 | 1.5 |
| | 10 years or younger | .4 | .0 | .0 | .0 | .0 | .0 | .0 |
| | 11 years | 19.9 | .2 | .0 | .0 | .0 | .0 | .0 |
| | 12 years | 66.0 | 18.0 | .1 | .0 | .0 | .0 | .0 |
| | 13 years | 11.9 | 69.4 | 19.2 | .1 | .0 | .0 | .0 |
| | 14 years | .7 | 10.4 | 67.1 | 18.1 | .1 | .0 | .0 |
| | 15 years | .0 | .5 | 11.5 | 67.9 | 16.0 | .3 | .0 |
| | 16 years | .0 | .0 | .7 | 11.9 | 70.6 | 17.7 | .2 |
| | 17 years | .0 | .0 | .0 | .7 | 10.9 | 71.4 | 17.2 |
| | 18 years or older | .0 | .0 | .0 | .0 | .6 | 9.2 | 81.1 |
| Grades | No Answer | 4.8 | 4.3 | 4.4 | 4.3 | 3.3 | 2.6 | 2.1 |
| | Mostly A's | 32.4 | 31.4 | 30.2 | 28.2 | 26.6 | 26.6 | 29.6 |
| | Mostly B's | 37.2 | 36.3 | 36.2 | 35.7 | 36.7 | 39.4 | 39.0 |
| | Mostly C's | 19.5 | 21.1 | 22.9 | 24.8 | 26.0 | 25.4 | 24.4 |
| | Mostly D's | 4.3 | 4.7 | 4.6 | 4.8 | 5.6 | 4.6 | 3.6 |
| | Mostly F's | 1.8 | 2.3 | 1.7 | 2.2 | 1.9 | 1.4 | 1.2 |
| Live with | No Answer | 1.8 | 1.8 | 1.8 | 1.9 | 1.4 | 1.4 | 1.1 |
| | Both parents | 60.4 | 60.6 | 57.8 | 57.4 | 57.2 | 57.1 | 58.1 |
| | Mother only | 13.6 | 12.4 | 14.1 | 14.5 | 14.8 | 15.5 | 15.4 |
| | Father only | 2.7 | 3.3 | 3.7 | 4.4 | 4.9 | 5.0 | 5.0 |
| | Mother & stepfather | 11.8 | 12.3 | 13.2 | 12.7 | 12.1 | 11.8 | 10.0 |
| | Father & stepmother | 2.9 | 3.6 | 3.8 | 3.8 | 4.2 | 3.6 | 3.3 |
| | Other | 6.7 | 6.1 | 5.6 | 5.3 | 5.4 | 5.6 | 7.1 |
| Employed - You | No Answer | 19.3 | 16.7 | 16.3 | 13.2 | 10.9 | 7.6 | 6.1 |
| | Yes, full time | 1.1 | 1.5 | 1.5 | 2.0 | 2.4 | 2.8 | 4.3 |
| | Yes, part time | 8.2 | 10.7 | 13.7 | 16.3 | 22.9 | 35.8 | 45.1 |
| | No | 71.4 | 71.1 | 68.5 | 68.5 | 63.8 | 53.9 | 44.6 |
| Employed - Mom | No Answer | 7.2 | 6.3 | 5.4 | 6.1 | 5.2 | 4.8 | 4.3 |
| | Yes, full time | 53.7 | 56.3 | 59.6 | 60.7 | 63.7 | 65.0 | 64.9 |
| | Yes, part time | 20.1 | 18.3 | 16.7 | 14.8 | 14.0 | 12.9 | 12.4 |
| | No | 18.9 | 19.1 | 18.3 | 18.4 | 17.1 | 17.3 | 18.3 |
| Employed - Dad | No Answer | 8.7 | 7.6 | 7.9 | 8.6 | 7.8 | 7.7 | 7.2 |
| | Yes, full time | 71.4 | 74.9 | 76.4 | 76.4 | 78.5 | 78.7 | 79.1 |
| | Yes, part time | 11.5 | 9.4 | 7.3 | 5.9 | 4.8 | 4.2 | 4.0 |
| | No | 8.4 | 8.1 | 8.4 | 9.2 | 8.8 | 9.4 | 9.7 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|-------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | | .4 | .3 | .5 | .4 | .5 | .6 | .6 |
| | Never | | 91.3 | 86.0 | 79.3 | 72.3 | 65.2 | 59.3 | 54.4 |
| | 1-5 times | | 6.3 | 8.7 | 10.9 | 12.0 | 13.9 | 14.8 | 14.3 |
| | 6-19 times | | .9 | 2.1 | 3.1 | 3.8 | 4.4 | 5.3 | 5.7 |
| | 20-40 times | | .4 | .8 | 1.7 | 2.1 | 3.0 | 2.9 | 3.9 |
| | 40+ times | | .7 | 2.1 | 4.5 | 9.4 | 13.0 | 17.0 | 21.2 |
| Smokeless Tobacco | No Answer | | .3 | .3 | .4 | .7 | .7 | .7 | .8 |
| | Never | | 95.8 | 91.9 | 86.6 | 79.6 | 72.6 | 68.1 | 62.5 |
| | 1-5 times | | 3.1 | 5.2 | 7.6 | 9.5 | 11.6 | 12.7 | 13.6 |
| | 6-19 times | | .5 | 1.1 | 1.8 | 3.2 | 4.3 | 4.4 | 5.5 |
| | 20-40 times | | .1 | .5 | .9 | 1.7 | 2.3 | 2.5 | 3.7 |
| | 40+ times | | .3 | .9 | 2.7 | 5.4 | 8.5 | 11.6 | 13.9 |
| Cigars | No Answer | | .4 | .4 | .6 | .7 | .9 | .9 | 1.2 |
| | Never | | 96.7 | 93.5 | 88.2 | 78.7 | 70.6 | 62.1 | 50.8 |
| | 1-5 times | | 2.3 | 4.4 | 7.0 | 11.3 | 13.5 | 15.3 | 17.2 |
| | 6-19 times | | .3 | .8 | 2.0 | 4.3 | 6.6 | 8.9 | 11.3 |
| | 20-40 times | | .1 | .3 | .8 | 2.2 | 3.6 | 4.8 | 7.8 |
| | 40+ times | | .2 | .5 | 1.3 | 2.8 | 4.8 | 8.0 | 11.8 |
| Pipe | No Answer | | .4 | .6 | .7 | 1.0 | 1.0 | 1.1 | 1.2 |
| | Never | | 97.9 | 95.3 | 92.1 | 86.5 | 80.7 | 74.6 | 66.3 |
| | 1-5 times | | 1.1 | 2.6 | 4.0 | 6.1 | 8.7 | 10.9 | 13.6 |
| | 6-19 times | | .2 | .7 | 1.3 | 2.4 | 3.7 | 5.1 | 7.4 |
| | 20-40 times | | .1 | .4 | .7 | 1.4 | 2.2 | 2.6 | 4.2 |
| | 40+ times | | .2 | .6 | 1.2 | 2.7 | 3.7 | 5.7 | 7.3 |
| Alcohol | No Answer | | .7 | .6 | .8 | .8 | 1.0 | 1.0 | 1.3 |
| | Never | | 79.4 | 72.0 | 62.7 | 53.0 | 44.3 | 38.3 | 33.1 |
| | 1-5 times | | 16.6 | 19.3 | 22.0 | 22.1 | 22.1 | 20.0 | 17.6 |
| | 6-19 times | | 2.3 | 4.6 | 8.1 | 11.9 | 13.7 | 14.6 | 13.7 |
| | 20-40 times | | .5 | 1.7 | 2.8 | 4.9 | 6.7 | 8.7 | 9.8 |
| | 40+ times | | .6 | 1.7 | 3.5 | 7.3 | 12.3 | 17.3 | 24.5 |
| Marijuana | No Answer | | .4 | .4 | .7 | .6 | 1.0 | 1.1 | 1.2 |
| | Never | | 95.7 | 91.3 | 83.9 | 74.4 | 66.2 | 60.1 | 56.6 |
| | 1-5 times | | 2.4 | 4.3 | 6.9 | 9.2 | 10.4 | 11.5 | 11.5 |
| | 6-19 times | | .6 | 1.3 | 2.7 | 4.1 | 5.3 | 5.7 | 6.5 |
| | 20-40 times | | .3 | .8 | 1.5 | 2.8 | 3.3 | 4.1 | 4.3 |
| | 40+ times | | .6 | 1.9 | 4.3 | 8.8 | 13.8 | 17.5 | 19.9 |
| Cocaine | No Answer | | .4 | .4 | .5 | .6 | .9 | .9 | 1.1 |
| | Never | | 98.9 | 98.1 | 97.2 | 95.7 | 94.2 | 92.0 | 90.8 |
| | 1-5 times | | .3 | .8 | 1.3 | 2.0 | 3.0 | 4.1 | 4.7 |
| | 6-19 times | | .1 | .2 | .4 | .6 | .7 | 1.2 | 1.5 |
| | 20-40 times | | .1 | .2 | .2 | .4 | .5 | .4 | .7 |
| | 40+ times | | .2 | .3 | .4 | .7 | .8 | 1.3 | 1.2 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | | Grade | | | | | | |
|------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Crack | No Answer | | .4 | .4 | .6 | .5 | .7 | .6 | .8 |
| | Never | | 98.9 | 98.2 | 97.7 | 97.1 | 96.4 | 95.9 | 95.5 |
| | 1-5 times | | .3 | .6 | 1.0 | 1.1 | 1.5 | 1.6 | 2.0 |
| | 6-19 times | | .1 | .2 | .3 | .5 | .4 | .6 | .6 |
| | 20-40 times | | .1 | .2 | .1 | .3 | .3 | .4 | .3 |
| | 40+ times | | .2 | .4 | .4 | .5 | .6 | .8 | .9 |
| Inhalants | No Answer | | 1.2 | .6 | .6 | .8 | .7 | .7 | .9 |
| | Never | | 96.6 | 96.2 | 94.2 | 92.3 | 91.4 | 90.2 | 89.6 |
| | 1-5 times | | 1.4 | 2.1 | 3.3 | 4.4 | 4.5 | 4.9 | 5.1 |
| | 6-19 times | | .4 | .5 | 1.0 | 1.3 | 1.9 | 2.2 | 2.1 |
| | 20-40 times | | .1 | .2 | .3 | .6 | .7 | 1.0 | 1.0 |
| | 40+ times | | .2 | .4 | .6 | .7 | .9 | 1.0 | 1.4 |
| Amphetamines | No Answer | | 1.8 | .9 | .7 | .7 | .9 | 1.0 | .9 |
| | Never | | 97.7 | 97.8 | 97.0 | 94.9 | 92.8 | 90.6 | 89.3 |
| | 1-5 times | | .3 | .7 | 1.1 | 2.1 | 2.7 | 3.4 | 3.5 |
| | 6-19 times | | .1 | .2 | .5 | 1.0 | 1.4 | 1.9 | 2.3 |
| | 20-40 times | | .0 | .1 | .3 | .6 | .9 | 1.1 | 1.5 |
| | 40+ times | | .1 | .2 | .5 | .8 | 1.4 | 2.0 | 2.5 |
| Methamphetamines | No Answer | | 1.0 | .6 | .6 | .7 | .8 | .8 | .8 |
| | Never | | 98.4 | 98.3 | 97.8 | 97.1 | 96.4 | 95.6 | 95.5 |
| | 1-5 times | | .2 | .6 | .7 | 1.0 | 1.3 | 1.5 | 1.6 |
| | 6-19 times | | .1 | .2 | .2 | .5 | .6 | .6 | .6 |
| | 20-40 times | | .1 | .1 | .2 | .3 | .3 | .5 | .5 |
| | 40+ times | | .1 | .2 | .5 | .4 | .7 | 1.0 | 1.0 |
| Heroin | No Answer | | 1.1 | .6 | .5 | .8 | .8 | .8 | 1.0 |
| | Never | | 98.4 | 98.4 | 98.1 | 97.2 | 96.6 | 95.9 | 95.6 |
| | 1-5 times | | .2 | .6 | .7 | 1.0 | 1.2 | 1.4 | 1.4 |
| | 6-19 times | | .1 | .2 | .2 | .3 | .4 | .5 | .6 |
| | 20-40 times | | .1 | .0 | .1 | .3 | .3 | .4 | .4 |
| | 40+ times | | .2 | .2 | .4 | .4 | .6 | 1.0 | 1.1 |
| Steroids | No Answer | | .9 | .6 | .6 | .8 | .7 | .7 | .9 |
| | Never | | 98.4 | 98.0 | 97.6 | 97.0 | 96.7 | 96.1 | 95.7 |
| | 1-5 times | | .5 | .8 | .9 | 1.0 | 1.2 | 1.3 | 1.4 |
| | 6-19 times | | .1 | .2 | .3 | .4 | .3 | .6 | .6 |
| | 20-40 times | | .1 | .0 | .1 | .2 | .3 | .3 | .4 |
| | 40+ times | | .2 | .3 | .5 | .6 | .7 | 1.0 | 1.0 |
| Ecstasy | No Answer | | 1.9 | .9 | .7 | 1.0 | .7 | .9 | 1.0 |
| | Never | | 97.5 | 97.8 | 96.9 | 95.2 | 93.8 | 91.9 | 90.6 |
| | 1-5 times | | .2 | .8 | 1.5 | 2.2 | 3.1 | 4.5 | 4.9 |
| | 6-19 times | | .1 | .2 | .3 | .7 | 1.0 | 1.2 | 1.8 |
| | 20-40 times | | .0 | .1 | .2 | .4 | .6 | .4 | .7 |
| | 40+ times | | .1 | .2 | .5 | .6 | .7 | 1.0 | 1.0 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--------------------------|-------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Tranquilizers | No Answer | 1.6 | .9 | .7 | 1.0 | .8 | 1.0 | 1.2 |
| | Never | 97.8 | 98.0 | 97.4 | 95.7 | 94.6 | 93.6 | 92.7 |
| | 1-5 times | .3 | .7 | 1.0 | 1.6 | 2.2 | 2.5 | 2.5 |
| | 6-19 times | .1 | .1 | .3 | .7 | .9 | 1.3 | 1.3 |
| | 20-40 times | .0 | .1 | .1 | .4 | .5 | .5 | .8 |
| | 40+ times | .1 | .1 | .4 | .7 | .9 | 1.2 | 1.5 |
| Hallucinogens | No Answer | 2.0 | 1.0 | .8 | 1.0 | 1.0 | 1.1 | 1.4 |
| | Never | 97.5 | 97.8 | 96.7 | 94.7 | 92.5 | 90.4 | 88.8 |
| | 1-5 times | .3 | .7 | 1.4 | 2.4 | 3.8 | 5.2 | 5.4 |
| | 6-19 times | .1 | .1 | .5 | .8 | 1.3 | 1.5 | 2.3 |
| | 20-40 times | .0 | .1 | .2 | .6 | .5 | .6 | .7 |
| | 40+ times | .1 | .3 | .5 | .6 | .9 | 1.2 | 1.5 |
| Prescription painkillers | No Answer | 1.1 | .7 | .9 | .8 | .9 | 1.0 | 1.1 |
| | Never | 97.7 | 96.5 | 93.9 | 89.6 | 85.8 | 82.1 | 79.9 |
| | 1-5 times | .7 | 1.5 | 2.7 | 4.6 | 5.4 | 6.6 | 6.5 |
| | 6-19 times | .2 | .6 | 1.1 | 2.3 | 3.5 | 4.0 | 4.7 |
| | 20-40 times | .1 | .3 | .6 | 1.0 | 1.8 | 2.0 | 2.8 |
| | 40+ times | .2 | .5 | .8 | 1.8 | 2.7 | 4.3 | 5.0 |
| Prescription drugs | No Answer | 1.2 | .8 | .9 | 1.0 | 1.0 | 1.1 | 1.3 |
| | Never | 97.6 | 96.8 | 94.7 | 90.9 | 87.5 | 84.6 | 82.2 |
| | 1-5 times | .6 | 1.4 | 2.3 | 3.6 | 4.8 | 5.2 | 5.9 |
| | 6-19 times | .2 | .4 | .7 | 1.8 | 2.5 | 3.4 | 3.8 |
| | 20-40 times | .1 | .2 | .6 | 1.0 | 1.4 | 1.9 | 2.0 |
| | 40+ times | .3 | .4 | .8 | 1.8 | 2.7 | 3.9 | 4.8 |
| Over the counter drugs | No Answer | 1.0 | .9 | .9 | 1.0 | 1.0 | 1.1 | 1.1 |
| | Never | 97.8 | 96.5 | 94.9 | 92.4 | 90.1 | 88.7 | 87.2 |
| | 1-5 times | .7 | 1.7 | 2.5 | 3.3 | 4.6 | 5.1 | 5.7 |
| | 6-19 times | .2 | .4 | .7 | 1.5 | 2.0 | 2.3 | 2.5 |
| | 20-40 times | .1 | .2 | .3 | .5 | .8 | 1.0 | 1.2 |
| | 40+ times | .2 | .4 | .7 | 1.3 | 1.5 | 2.0 | 2.3 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | | Grade | | | | | | |
|-------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | | 2.1 | 1.9 | 2.3 | 2.9 | 2.5 | 2.5 | 2.3 |
| | Never | | 95.2 | 92.2 | 88.2 | 82.9 | 79.0 | 75.5 | 69.8 |
| | 1-5 times | | 1.9 | 3.6 | 5.0 | 5.8 | 7.0 | 7.3 | 8.7 |
| | 6-19 times | | .4 | .9 | 1.6 | 2.5 | 3.3 | 3.2 | 4.3 |
| | 20-40 times | | .1 | .5 | 1.0 | 1.7 | 2.1 | 2.7 | 2.7 |
| | 40+ times | | .2 | .9 | 1.8 | 4.2 | 6.1 | 8.9 | 12.1 |
| Smokeless Tobacco | No Answer | | 2.4 | 2.1 | 2.8 | 3.1 | 3.0 | 2.9 | 2.6 |
| | Never | | 96.1 | 94.5 | 91.0 | 86.6 | 83.0 | 81.6 | 78.2 |
| | 1-5 times | | 1.0 | 2.2 | 3.4 | 5.1 | 5.8 | 5.8 | 6.5 |
| | 6-19 times | | .3 | .6 | 1.2 | 1.7 | 2.5 | 2.4 | 3.2 |
| | 20-40 times | | .1 | .3 | .5 | 1.2 | 1.5 | 1.9 | 2.4 |
| | 40+ times | | .1 | .3 | 1.1 | 2.4 | 4.2 | 5.5 | 7.0 |
| Cigars | No Answer | | 2.6 | 2.1 | 3.0 | 3.6 | 3.1 | 3.1 | 2.9 |
| | Never | | 96.4 | 95.7 | 92.6 | 87.9 | 84.1 | 80.2 | 72.0 |
| | 1-5 times | | .7 | 1.6 | 3.1 | 5.8 | 8.5 | 10.9 | 16.5 |
| | 6-19 times | | .1 | .3 | .7 | 1.6 | 2.4 | 3.3 | 4.8 |
| | 20-40 times | | .0 | .1 | .2 | .5 | .8 | 1.0 | 1.7 |
| | 40+ times | | .1 | .1 | .3 | .6 | 1.1 | 1.5 | 2.2 |
| Pipe | No Answer | | 2.7 | 2.3 | 3.3 | 3.8 | 3.3 | 3.3 | 3.2 |
| | Never | | 96.5 | 96.0 | 93.4 | 90.5 | 88.7 | 86.7 | 81.7 |
| | 1-5 times | | .5 | 1.1 | 2.0 | 3.3 | 4.5 | 5.8 | 9.5 |
| | 6-19 times | | .1 | .3 | .7 | 1.1 | 1.5 | 2.3 | 2.7 |
| | 20-40 times | | .1 | .2 | .3 | .5 | .9 | .7 | 1.0 |
| | 40+ times | | .1 | .1 | .3 | .8 | 1.1 | 1.2 | 1.8 |
| Alcohol | No Answer | | 2.9 | 2.3 | 3.2 | 3.4 | 3.1 | 3.1 | 2.6 |
| | Never | | 91.3 | 87.8 | 81.9 | 73.9 | 68.0 | 63.6 | 57.1 |
| | 1-5 times | | 4.9 | 7.7 | 11.2 | 15.8 | 19.6 | 21.2 | 24.2 |
| | 6-19 times | | .6 | 1.3 | 2.3 | 4.2 | 5.6 | 7.6 | 10.3 |
| | 20-40 times | | .1 | .4 | .7 | 1.2 | 1.9 | 2.2 | 2.8 |
| | 40+ times | | .1 | .4 | .7 | 1.4 | 1.8 | 2.3 | 3.1 |
| Marijuana | No Answer | | 2.8 | 2.2 | 3.2 | 3.6 | 3.3 | 3.2 | 3.1 |
| | Never | | 95.3 | 93.3 | 88.2 | 82.3 | 78.1 | 75.2 | 73.2 |
| | 1-5 times | | 1.2 | 2.4 | 4.6 | 6.4 | 7.7 | 8.6 | 8.6 |
| | 6-19 times | | .3 | .9 | 1.6 | 3.0 | 3.8 | 4.2 | 4.6 |
| | 20-40 times | | .2 | .6 | 1.0 | 1.6 | 2.4 | 2.9 | 3.2 |
| | 40+ times | | .3 | .6 | 1.5 | 3.1 | 4.8 | 5.9 | 7.4 |
| Cocaine | No Answer | | 3.0 | 2.4 | 3.3 | 3.8 | 3.4 | 3.3 | 3.3 |
| | Never | | 96.7 | 96.8 | 95.4 | 94.3 | 94.5 | 94.0 | 93.9 |
| | 1-5 times | | .2 | .5 | .7 | 1.1 | 1.2 | 1.5 | 1.7 |
| | 6-19 times | | .1 | .2 | .3 | .4 | .5 | .4 | .6 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .2 | .3 | .3 |
| | 40+ times | | .1 | .1 | .1 | .2 | .3 | .5 | .3 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------|-------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Crack | No Answer | 3.0 | 2.4 | 3.3 | 3.9 | 3.4 | 3.3 | 3.2 |
| | Never | 96.7 | 96.8 | 95.7 | 94.8 | 95.2 | 94.9 | 95.1 |
| | 1-5 times | .1 | .5 | .6 | .6 | .7 | 1.0 | .9 |
| | 6-19 times | .1 | .1 | .3 | .4 | .4 | .3 | .3 |
| | 20-40 times | .1 | .1 | .0 | .1 | .2 | .2 | .3 |
| | 40+ times | .1 | .1 | .1 | .2 | .2 | .3 | .2 |
| Inhalants | No Answer | 3.6 | 2.4 | 3.3 | 3.9 | 3.5 | 3.4 | 3.3 |
| | Never | 95.5 | 96.2 | 94.6 | 94.0 | 94.1 | 94.5 | 94.4 |
| | 1-5 times | .7 | 1.1 | 1.5 | 1.5 | 1.6 | 1.4 | 1.4 |
| | 6-19 times | .1 | .2 | .4 | .4 | .5 | .5 | .6 |
| | 20-40 times | .0 | .1 | .1 | .1 | .2 | .2 | .2 |
| | 40+ times | .1 | .0 | .1 | .1 | .1 | .1 | .2 |
| Amphetamines | No Answer | 4.0 | 2.7 | 3.4 | 3.8 | 3.6 | 3.5 | 3.4 |
| | Never | 95.7 | 96.7 | 95.3 | 94.3 | 93.8 | 93.8 | 93.2 |
| | 1-5 times | .2 | .5 | .8 | 1.1 | 1.6 | 1.6 | 2.0 |
| | 6-19 times | .1 | .2 | .3 | .5 | .6 | .6 | .8 |
| | 20-40 times | .0 | .0 | .1 | .2 | .2 | .3 | .3 |
| | 40+ times | .0 | .0 | .1 | .1 | .1 | .1 | .3 |
| Methamphetamines | No Answer | 3.6 | 2.5 | 3.4 | 3.9 | 3.6 | 3.4 | 3.4 |
| | Never | 96.1 | 96.9 | 95.6 | 95.1 | 95.2 | 94.7 | 94.9 |
| | 1-5 times | .2 | .4 | .6 | .5 | .6 | 1.0 | .8 |
| | 6-19 times | .1 | .1 | .2 | .3 | .4 | .5 | .5 |
| | 20-40 times | .0 | .0 | .1 | .1 | .1 | .1 | .2 |
| | 40+ times | .1 | .1 | .1 | .1 | .1 | .3 | .3 |
| Heroin | No Answer | 3.5 | 2.6 | 3.4 | 3.9 | 3.6 | 3.6 | 3.3 |
| | Never | 96.2 | 96.8 | 95.8 | 95.1 | 95.2 | 94.7 | 94.8 |
| | 1-5 times | .1 | .3 | .5 | .5 | .6 | .8 | .9 |
| | 6-19 times | .0 | .1 | .2 | .2 | .3 | .4 | .5 |
| | 20-40 times | .0 | .0 | .1 | .2 | .2 | .3 | .2 |
| | 40+ times | .1 | .0 | .1 | .2 | .2 | .3 | .2 |
| Steroids | No Answer | 3.3 | 2.6 | 3.4 | 3.9 | 3.6 | 3.5 | 3.4 |
| | Never | 96.3 | 96.7 | 95.6 | 94.9 | 95.1 | 94.9 | 94.8 |
| | 1-5 times | .2 | .4 | .5 | .5 | .6 | .7 | .8 |
| | 6-19 times | .1 | .1 | .3 | .3 | .3 | .4 | .5 |
| | 20-40 times | .0 | .1 | .1 | .1 | .2 | .2 | .2 |
| | 40+ times | .0 | .1 | .1 | .2 | .2 | .3 | .2 |
| Ecstasy | No Answer | 4.2 | 2.8 | 3.4 | 4.0 | 3.6 | 3.7 | 3.4 |
| | Never | 95.5 | 96.6 | 95.3 | 94.3 | 94.2 | 94.0 | 93.6 |
| | 1-5 times | .2 | .4 | .8 | 1.1 | 1.5 | 1.4 | 2.0 |
| | 6-19 times | .1 | .1 | .2 | .4 | .4 | .5 | .6 |
| | 20-40 times | .0 | .1 | .1 | .2 | .2 | .2 | .3 |
| | 40+ times | .0 | .0 | .1 | .1 | .1 | .2 | .2 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|--------------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Tranquilizers | No Answer | | 4.1 | 2.8 | 3.4 | 4.0 | 3.8 | 3.7 | 3.5 |
| | Never | | 95.6 | 96.7 | 95.6 | 94.4 | 94.1 | 94.0 | 94.2 |
| | 1-5 times | | .2 | .3 | .6 | .9 | 1.2 | 1.3 | 1.4 |
| | 6-19 times | | .0 | .1 | .3 | .4 | .4 | .6 | .6 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .1 | .2 | .2 |
| | 40+ times | | .1 | .1 | .1 | .2 | .4 | .2 | .2 |
| Hallucinogens | No Answer | | 4.3 | 2.9 | 3.6 | 4.1 | 3.7 | 3.7 | 3.5 |
| | Never | | 95.4 | 96.5 | 95.1 | 94.1 | 93.9 | 93.2 | 92.9 |
| | 1-5 times | | .2 | .4 | .9 | 1.2 | 1.6 | 2.2 | 2.4 |
| | 6-19 times | | .0 | .1 | .3 | .3 | .4 | .6 | .7 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .2 | .2 | .3 |
| | 40+ times | | .0 | .1 | .1 | .1 | .2 | .2 | .3 |
| Prescription painkillers | No Answer | | 3.7 | 2.7 | 3.6 | 3.9 | 3.5 | 3.7 | 3.5 |
| | Never | | 95.7 | 95.8 | 93.7 | 91.6 | 90.3 | 89.5 | 88.3 |
| | 1-5 times | | .4 | .9 | 1.8 | 2.9 | 3.6 | 3.8 | 4.7 |
| | 6-19 times | | .1 | .3 | .6 | 1.0 | 1.6 | 1.9 | 2.2 |
| | 20-40 times | | .0 | .1 | .2 | .4 | .5 | .6 | .8 |
| | 40+ times | | .1 | .1 | .2 | .3 | .4 | .5 | .6 |
| Prescription drugs | No Answer | | 3.7 | 2.8 | 3.7 | 4.2 | 3.7 | 3.8 | 3.5 |
| | Never | | 95.7 | 95.8 | 93.9 | 91.9 | 91.0 | 90.2 | 89.6 |
| | 1-5 times | | .4 | .7 | 1.6 | 2.4 | 3.2 | 3.4 | 4.1 |
| | 6-19 times | | .1 | .3 | .5 | .8 | 1.1 | 1.5 | 1.5 |
| | 20-40 times | | .1 | .2 | .1 | .4 | .6 | .5 | .5 |
| | 40+ times | | .1 | .2 | .2 | .3 | .4 | .6 | .8 |
| Over the counter drugs | No Answer | | 4.1 | 3.3 | 3.9 | 4.4 | 4.1 | 4.1 | 3.8 |
| | Never | | 95.2 | 95.3 | 94.0 | 92.6 | 92.3 | 92.4 | 92.2 |
| | 1-5 times | | .4 | 1.0 | 1.4 | 1.9 | 2.3 | 2.2 | 2.7 |
| | 6-19 times | | .2 | .2 | .4 | .6 | .7 | .7 | .8 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .3 | .3 | .3 |
| | 40+ times | | .1 | .1 | .2 | .3 | .3 | .3 | .3 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | 2.8 | 2.6 | 3.3 | 3.9 | 3.4 | 3.6 | 3.2 |
| | Never | 89.8 | 85.0 | 78.1 | 71.0 | 64.6 | 58.7 | 54.2 |
| | 10 years or younger | 3.9 | 4.8 | 4.9 | 5.1 | 5.6 | 4.6 | 4.2 |
| | 11 years | 2.1 | 2.9 | 3.0 | 2.9 | 2.7 | 2.7 | 1.9 |
| | 12 years | 1.2 | 2.8 | 3.6 | 4.0 | 3.7 | 3.7 | 3.0 |
| | 13 years | .2 | 1.7 | 4.8 | 4.9 | 5.0 | 5.1 | 4.0 |
| | 14 years | .0 | .2 | 1.9 | 5.6 | 5.1 | 5.1 | 4.4 |
| | 15 years | .0 | .0 | .3 | 2.5 | 6.7 | 6.8 | 6.5 |
| | 16 years | .0 | .0 | .0 | .1 | 2.9 | 6.9 | 8.6 |
| | 17 years or older | .0 | .0 | .0 | .0 | .3 | 2.8 | 9.9 |
| Smokeless Tobacco | No Answer | 3.0 | 2.8 | 3.7 | 4.3 | 4.0 | 4.0 | 3.9 |
| | Never | 93.7 | 90.7 | 84.9 | 77.9 | 71.5 | 67.2 | 61.8 |
| | 10 years or younger | 1.4 | 2.0 | 2.0 | 2.4 | 2.3 | 2.2 | 2.2 |
| | 11 years | 1.0 | 1.4 | 1.4 | 1.4 | 1.3 | 1.1 | .8 |
| | 12 years | .7 | 1.6 | 2.2 | 2.5 | 2.3 | 2.1 | 1.7 |
| | 13 years | .2 | 1.3 | 3.5 | 3.6 | 3.5 | 3.6 | 2.9 |
| | 14 years | .0 | .2 | 1.9 | 4.8 | 4.7 | 4.2 | 3.5 |
| | 15 years | .0 | .0 | .2 | 2.7 | 7.2 | 6.6 | 6.1 |
| | 16 years | .0 | .0 | .0 | .3 | 3.0 | 6.5 | 8.7 |
| | 17 years or older | .0 | .0 | .0 | .0 | .2 | 2.5 | 8.3 |
| Cigars | No Answer | 3.2 | 3.0 | 4.0 | 4.5 | 4.2 | 4.1 | 3.8 |
| | Never | 94.6 | 92.0 | 86.7 | 77.9 | 70.5 | 62.7 | 51.8 |
| | 10 years or younger | 1.0 | 1.4 | 1.5 | 1.7 | 1.9 | 1.9 | 1.7 |
| | 11 years | .6 | 1.0 | 1.3 | 1.3 | 1.3 | 1.1 | .8 |
| | 12 years | .6 | 1.4 | 1.8 | 2.4 | 2.1 | 2.1 | 1.6 |
| | 13 years | .1 | 1.0 | 3.0 | 3.9 | 3.6 | 3.4 | 2.7 |
| | 14 years | .0 | .2 | 1.5 | 5.3 | 4.6 | 4.7 | 4.2 |
| | 15 years | .0 | .0 | .2 | 2.8 | 8.1 | 8.0 | 7.1 |
| | 16 years | .0 | .0 | .0 | .2 | 3.5 | 8.7 | 11.1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .2 | 3.3 | 15.1 |
| Pipe | No Answer | 3.3 | 3.3 | 4.2 | 5.1 | 5.0 | 4.8 | 4.5 |
| | Never | 95.7 | 93.9 | 90.4 | 85.0 | 80.2 | 74.8 | 67.6 |
| | 10 years or younger | .4 | .8 | .8 | 1.0 | 1.1 | 1.0 | 1.1 |
| | 11 years | .2 | .6 | .8 | .7 | .7 | .5 | .4 |
| | 12 years | .3 | .8 | 1.0 | 1.3 | 1.1 | 1.3 | .7 |
| | 13 years | .0 | .6 | 1.7 | 2.3 | 2.0 | 2.0 | 1.5 |
| | 14 years | .0 | .1 | .9 | 2.8 | 3.0 | 2.5 | 2.1 |
| | 15 years | .0 | .0 | .2 | 1.7 | 4.6 | 4.8 | 4.1 |
| | 16 years | .0 | .0 | .0 | .2 | 2.1 | 5.9 | 6.9 |
| | 17 years or older | .0 | .0 | .0 | .0 | .2 | 2.3 | 11.2 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-----------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Alcohol | No Answer | 3.6 | 3.4 | 4.3 | 4.5 | 4.3 | 4.4 | 3.8 |
| | Never | 79.1 | 72.8 | 63.6 | 54.0 | 46.1 | 40.0 | 34.8 |
| | 10 years or younger | 9.7 | 8.8 | 8.2 | 6.7 | 5.9 | 5.7 | 4.7 |
| | 11 years | 4.4 | 5.5 | 4.6 | 3.7 | 2.7 | 1.8 | 1.6 |
| | 12 years | 2.7 | 5.8 | 6.3 | 5.7 | 4.9 | 4.4 | 3.4 |
| | 13 years | .4 | 3.1 | 8.4 | 8.9 | 7.5 | 6.8 | 5.7 |
| | 14 years | .0 | .5 | 4.0 | 10.2 | 9.9 | 8.4 | 7.5 |
| | 15 years | .0 | .0 | .5 | 5.7 | 13.3 | 11.9 | 11.9 |
| | 16 years | .0 | .0 | .1 | .5 | 5.0 | 12.2 | 13.8 |
| | 17 years or older | .0 | .0 | .0 | .1 | .5 | 4.4 | 12.8 |
| Marijuana | No Answer | 3.3 | 3.2 | 3.8 | 4.4 | 4.1 | 4.0 | 3.9 |
| | Never | 93.6 | 90.0 | 83.0 | 73.6 | 66.4 | 60.8 | 57.5 |
| | 10 years or younger | .9 | 1.4 | 1.8 | 1.9 | 2.1 | 1.8 | 1.5 |
| | 11 years | 1.0 | 1.5 | 1.4 | 1.6 | 1.5 | 1.4 | 1.0 |
| | 12 years | .9 | 1.9 | 2.4 | 3.0 | 2.7 | 2.6 | 2.1 |
| | 13 years | .2 | 1.7 | 4.7 | 5.1 | 4.1 | 3.7 | 3.2 |
| | 14 years | .0 | .3 | 2.6 | 6.5 | 6.0 | 5.4 | 4.7 |
| | 15 years | .0 | .0 | .3 | 3.6 | 8.9 | 8.3 | 7.3 |
| | 16 years | .0 | .0 | .0 | .3 | 3.8 | 9.0 | 9.3 |
| | 17 years or older | .0 | .0 | .0 | .1 | .3 | 3.1 | 9.5 |
| Cocaine | No Answer | 3.2 | 3.0 | 3.9 | 4.7 | 4.5 | 4.6 | 4.2 |
| | Never | 96.4 | 96.0 | 94.6 | 92.8 | 91.8 | 90.2 | 89.4 |
| | 10 years or younger | .2 | .3 | .3 | .3 | .4 | .5 | .4 |
| | 11 years | .1 | .2 | .2 | .2 | .1 | .2 | .2 |
| | 12 years | .1 | .2 | .2 | .3 | .3 | .2 | .2 |
| | 13 years | .0 | .2 | .4 | .4 | .3 | .3 | .4 |
| | 14 years | .0 | .0 | .3 | .6 | .6 | .6 | .5 |
| | 15 years | .0 | .0 | .1 | .5 | 1.2 | 1.0 | 1.0 |
| | 16 years | .0 | .0 | .0 | .1 | .7 | 1.5 | 1.5 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .9 | 2.2 |
| Crack | No Answer | 3.3 | 3.2 | 4.1 | 4.8 | 4.5 | 4.7 | 4.3 |
| | Never | 96.3 | 96.0 | 94.8 | 93.7 | 93.5 | 93.1 | 93.1 |
| | 10 years or younger | .2 | .3 | .2 | .3 | .4 | .3 | .4 |
| | 11 years | .1 | .2 | .1 | .2 | .1 | .2 | .1 |
| | 12 years | .1 | .2 | .2 | .2 | .2 | .2 | .2 |
| | 13 years | .0 | .2 | .3 | .2 | .3 | .2 | .2 |
| | 14 years | .0 | .0 | .2 | .3 | .3 | .2 | .3 |
| | 15 years | .0 | .0 | .1 | .3 | .5 | .4 | .4 |
| | 16 years | .0 | .0 | .0 | .1 | .3 | .4 | .4 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .2 | .5 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Inhalants | No Answer | 4.2 | 3.3 | 4.0 | 4.9 | 4.6 | 4.7 | 4.4 |
| | Never | 94.3 | 94.5 | 92.5 | 90.7 | 90.2 | 89.4 | 89.3 |
| | 10 years or younger | .8 | .7 | .6 | .5 | .5 | .6 | .6 |
| | 11 years | .3 | .4 | .5 | .4 | .2 | .3 | .3 |
| | 12 years | .3 | .5 | .7 | .6 | .5 | .6 | .3 |
| | 13 years | .1 | .4 | 1.0 | 1.1 | .8 | .8 | .8 |
| | 14 years | .0 | .1 | .5 | 1.1 | .9 | 1.0 | .7 |
| | 15 years | .0 | .0 | .1 | .6 | 1.6 | 1.3 | 1.1 |
| | 16 years | .0 | .0 | .0 | .1 | .7 | 1.1 | 1.1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .3 | 1.3 |
| Amphetamines | No Answer | 4.4 | 3.5 | 4.0 | 4.8 | 4.7 | 4.8 | 4.4 |
| | Never | 95.3 | 95.7 | 94.4 | 92.4 | 91.0 | 89.6 | 89.1 |
| | 10 years or younger | .2 | .2 | .3 | .3 | .3 | .3 | .3 |
| | 11 years | .1 | .2 | .2 | .1 | .1 | .2 | .1 |
| | 12 years | .0 | .2 | .3 | .4 | .3 | .4 | .4 |
| | 13 years | .0 | .2 | .4 | .5 | .5 | .6 | .5 |
| | 14 years | .0 | .0 | .3 | .8 | 1.0 | 1.0 | .9 |
| | 15 years | .0 | .0 | .1 | .7 | 1.4 | 1.2 | 1.2 |
| | 16 years | .0 | .0 | .0 | .0 | .7 | 1.5 | 1.8 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .4 | 1.4 |
| Methamphetamines | No Answer | 4.0 | 3.2 | 3.9 | 4.8 | 4.7 | 4.7 | 4.5 |
| | Never | 95.7 | 96.0 | 95.1 | 93.8 | 93.4 | 92.9 | 93.0 |
| | 10 years or younger | .2 | .2 | .3 | .3 | .2 | .4 | .3 |
| | 11 years | .1 | .2 | .1 | .2 | .1 | .1 | .1 |
| | 12 years | .0 | .2 | .1 | .1 | .2 | .1 | .1 |
| | 13 years | .0 | .2 | .3 | .2 | .2 | .1 | .2 |
| | 14 years | .0 | .0 | .1 | .3 | .4 | .2 | .2 |
| | 15 years | .0 | .0 | .1 | .3 | .4 | .5 | .4 |
| | 16 years | .0 | .0 | .0 | .1 | .4 | .6 | .5 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .4 | .8 |
| Heroin | No Answer | 4.0 | 3.4 | 4.1 | 4.8 | 4.7 | 4.6 | 4.4 |
| | Never | 95.7 | 96.1 | 95.0 | 94.0 | 93.5 | 93.0 | 93.1 |
| | 10 years or younger | .1 | .2 | .2 | .3 | .3 | .5 | .3 |
| | 11 years | .0 | .2 | .1 | .1 | .1 | .1 | .1 |
| | 12 years | .1 | .0 | .2 | .2 | .2 | .1 | .1 |
| | 13 years | .0 | .1 | .2 | .1 | .2 | .1 | .1 |
| | 14 years | .0 | .0 | .2 | .2 | .3 | .2 | .3 |
| | 15 years | .0 | .0 | .0 | .3 | .4 | .4 | .4 |
| | 16 years | .0 | .0 | .0 | .0 | .3 | .6 | .5 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .3 | .7 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|---------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Steroids | No Answer | 3.7 | 3.3 | 4.1 | 4.8 | 4.7 | 4.8 | 4.4 |
| | Never | 95.8 | 95.8 | 94.7 | 93.8 | 93.6 | 93.1 | 93.2 |
| | 10 years or younger | .2 | .3 | .3 | .2 | .3 | .3 | .4 |
| | 11 years | .1 | .2 | .2 | .1 | .1 | .1 | .0 |
| | 12 years | .1 | .2 | .1 | .2 | .1 | .1 | .1 |
| | 13 years | .0 | .2 | .3 | .2 | .2 | .1 | .2 |
| | 14 years | .0 | .0 | .1 | .3 | .2 | .2 | .2 |
| | 15 years | .0 | .0 | .1 | .3 | .4 | .4 | .3 |
| | 16 years | .0 | .0 | .0 | .0 | .4 | .6 | .5 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .3 | .7 |
| | | | | | | | | |
| Ecstasy | No Answer | 4.7 | 3.5 | 4.2 | 4.9 | 4.6 | 4.7 | 4.3 |
| | Never | 95.0 | 95.7 | 94.1 | 92.5 | 91.6 | 90.2 | 89.4 |
| | 10 years or younger | .1 | .2 | .3 | .3 | .3 | .4 | .4 |
| | 11 years | .1 | .2 | .2 | .2 | .1 | .2 | .1 |
| | 12 years | .1 | .2 | .3 | .3 | .1 | .1 | .2 |
| | 13 years | .0 | .2 | .4 | .5 | .4 | .3 | .4 |
| | 14 years | .0 | .0 | .5 | .7 | .7 | .5 | .4 |
| | 15 years | .0 | .0 | .1 | .5 | 1.4 | 1.2 | .9 |
| | 16 years | .0 | .0 | .0 | .1 | .8 | 1.7 | 1.9 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .7 | 2.2 |
| | | | | | | | | |
| Tranquilizers | No Answer | 4.3 | 3.7 | 4.3 | 5.0 | 4.8 | 4.7 | 4.5 |
| | Never | 95.4 | 95.7 | 94.4 | 92.8 | 92.1 | 91.4 | 91.2 |
| | 10 years or younger | .2 | .2 | .3 | .2 | .3 | .4 | .3 |
| | 11 years | .1 | .2 | .2 | .2 | .1 | .1 | .1 |
| | 12 years | .0 | .1 | .2 | .2 | .3 | .2 | .2 |
| | 13 years | .0 | .1 | .3 | .6 | .4 | .5 | .5 |
| | 14 years | .0 | .0 | .2 | .6 | .6 | .6 | .6 |
| | 15 years | .0 | .0 | .0 | .3 | .9 | .9 | .8 |
| | 16 years | .0 | .0 | .0 | .1 | .5 | .8 | 1.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .4 | .9 |
| | | | | | | | | |
| Hallucinogens | No Answer | 4.8 | 3.6 | 4.3 | 5.1 | 4.9 | 4.7 | 4.7 |
| | Never | 95.0 | 95.5 | 93.9 | 92.0 | 90.6 | 89.3 | 88.0 |
| | 10 years or younger | .1 | .2 | .2 | .3 | .3 | .4 | .4 |
| | 11 years | .0 | .2 | .2 | .2 | .2 | .1 | .1 |
| | 12 years | .1 | .2 | .3 | .4 | .2 | .2 | .2 |
| | 13 years | .0 | .2 | .6 | .6 | .5 | .4 | .3 |
| | 14 years | .0 | .0 | .4 | .8 | .7 | .7 | .5 |
| | 15 years | .0 | .0 | .1 | .6 | 1.5 | 1.1 | .9 |
| | 16 years | .0 | .0 | .0 | .1 | 1.0 | 2.1 | 1.9 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .9 | 2.9 |
| | | | | | | | | |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--------------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Prescription painkillers | No Answer | 4.0 | 3.4 | 4.2 | 5.0 | 4.6 | 4.8 | 4.5 |
| | Never | 95.3 | 94.6 | 91.8 | 87.7 | 84.8 | 82.0 | 80.3 |
| | 10 years or younger | .3 | .4 | .5 | .4 | .5 | .4 | .5 |
| | 11 years | .1 | .4 | .5 | .3 | .3 | .3 | .2 |
| | 12 years | .2 | .6 | .7 | .9 | .8 | .5 | .4 |
| | 13 years | .0 | .5 | 1.3 | 1.5 | 1.3 | 1.2 | .9 |
| | 14 years | .0 | .1 | .8 | 2.5 | 2.2 | 1.8 | 1.8 |
| | 15 years | .0 | .0 | .2 | 1.5 | 3.5 | 3.3 | 2.7 |
| | 16 years | .0 | .0 | .0 | .1 | 1.9 | 3.8 | 4.6 |
| | 17 years or older | .0 | .0 | .0 | .1 | .2 | 1.7 | 4.1 |
| Prescription drugs | No Answer | 4.3 | 3.4 | 4.2 | 4.9 | 4.6 | 4.7 | 4.5 |
| | Never | 95.0 | 94.8 | 92.2 | 88.6 | 86.1 | 83.9 | 82.1 |
| | 10 years or younger | .4 | .5 | .5 | .5 | .6 | .5 | .6 |
| | 11 years | .1 | .3 | .4 | .3 | .3 | .2 | .1 |
| | 12 years | .1 | .5 | .6 | .7 | .5 | .3 | .4 |
| | 13 years | .0 | .4 | 1.1 | 1.4 | 1.1 | 1.1 | .8 |
| | 14 years | .0 | .1 | .7 | 2.0 | 1.9 | 1.5 | 1.5 |
| | 15 years | .0 | .0 | .1 | 1.4 | 3.1 | 2.9 | 2.5 |
| | 16 years | .0 | .0 | .0 | .1 | 1.8 | 3.3 | 3.8 |
| | 17 years or older | .0 | .0 | .0 | .1 | .1 | 1.5 | 3.8 |
| Over the counter drugs | No Answer | 4.4 | 3.9 | 4.7 | 5.3 | 5.1 | 5.2 | 4.8 |
| | Never | 94.9 | 94.1 | 92.2 | 90.1 | 88.1 | 87.1 | 86.4 |
| | 10 years or younger | .3 | .5 | .4 | .5 | .5 | .5 | .4 |
| | 11 years | .2 | .3 | .4 | .3 | .3 | .1 | .2 |
| | 12 years | .2 | .6 | .6 | .6 | .5 | .4 | .4 |
| | 13 years | .1 | .5 | 1.1 | 1.0 | .8 | .7 | .6 |
| | 14 years | .0 | .1 | .6 | 1.2 | 1.5 | 1.2 | 1.0 |
| | 15 years | .0 | .0 | .1 | .8 | 2.2 | 2.0 | 1.4 |
| | 16 years | .0 | .0 | .0 | .1 | 1.0 | 2.0 | 2.5 |
| | 17 years or older | .0 | .0 | .0 | .1 | .1 | .8 | 2.3 |

**Perceived Availability of Drugs
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------------------|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Availability of cigarettes | No Answer | 10.3 | 9.2 | 8.5 | 7.8 | 5.9 | 5.3 | 4.6 |
| | Very difficult | 61.5 | 52.2 | 42.4 | 31.2 | 24.0 | 17.1 | 8.0 |
| | Fairly difficult | 10.2 | 12.5 | 13.2 | 12.2 | 10.9 | 7.9 | 2.8 |
| | Fairly easy | 8.6 | 12.9 | 16.9 | 20.0 | 21.9 | 20.8 | 8.5 |
| | Easy | 9.4 | 13.2 | 19.1 | 28.9 | 37.4 | 48.9 | 76.0 |
| Availability of alcohol | No Answer | 10.6 | 9.3 | 8.5 | 7.6 | 5.7 | 5.1 | 4.5 |
| | Very difficult | 51.6 | 42.2 | 32.6 | 23.6 | 18.0 | 14.5 | 11.7 |
| | Fairly difficult | 14.1 | 15.6 | 15.5 | 14.1 | 13.4 | 11.6 | 11.2 |
| | Fairly easy | 11.7 | 15.7 | 20.0 | 24.1 | 26.0 | 27.4 | 30.6 |
| | Easy | 12.0 | 17.2 | 23.4 | 30.5 | 36.9 | 41.5 | 42.0 |
| Availability of marijuana | No Answer | 11.3 | 9.8 | 9.1 | 8.3 | 6.3 | 5.7 | 5.0 |
| | Very difficult | 79.6 | 71.9 | 59.0 | 42.5 | 32.5 | 25.8 | 21.1 |
| | Fairly difficult | 3.8 | 6.6 | 9.8 | 11.3 | 11.6 | 11.6 | 11.5 |
| | Fairly easy | 2.3 | 5.1 | 9.5 | 14.5 | 17.9 | 19.3 | 22.0 |
| | Easy | 3.0 | 6.6 | 12.6 | 23.4 | 31.7 | 37.5 | 40.4 |
| Availability of drugs | No Answer | 11.7 | 10.2 | 9.7 | 9.2 | 7.3 | 6.6 | 5.6 |
| | Very difficult | 83.2 | 79.7 | 75.0 | 63.4 | 57.4 | 51.0 | 47.2 |
| | Fairly difficult | 2.4 | 5.0 | 7.6 | 13.0 | 16.8 | 19.1 | 21.7 |
| | Fairly easy | .9 | 1.9 | 3.3 | 6.5 | 9.1 | 11.1 | 12.1 |
| | Easy | 1.7 | 3.1 | 4.3 | 7.9 | 9.4 | 12.2 | 13.4 |

**Binge Drinking in the Past Two Weeks
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Binge drinking | No Answer | 3.7 | 3.2 | 2.8 | 3.0 | 2.7 | 3.1 | 2.9 |
| | None | 90.0 | 89.0 | 86.8 | 81.5 | 77.5 | 73.1 | 67.2 |
| | Once | 2.5 | 3.1 | 4.1 | 5.8 | 7.5 | 8.6 | 9.9 |
| | Twice | 1.4 | 1.7 | 2.3 | 3.7 | 4.5 | 5.5 | 7.4 |
| | 3-5 times | 1.4 | 1.6 | 2.2 | 3.0 | 4.5 | 5.5 | 7.5 |
| | 6-9 times | .4 | .7 | .8 | 1.3 | 1.5 | 2.1 | 2.5 |
| | 10+ times | .6 | .7 | 1.0 | 1.7 | 1.8 | 2.1 | 2.6 |

**Main Sources of Alcoholic Beverages in the Past Month
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------------|---|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Main sources of alcoholic beverages | No Answer | 7.3 | 6.9 | 5.8 | 6.5 | 6.9 | 6.9 | 7.8 |
| | No drink | 86.7 | 83.3 | 79.4 | 72.3 | 66.2 | 61.9 | 54.4 |
| | Restaurants/bars/clubs | .1 | .1 | .1 | .2 | .2 | .3 | .4 |
| | Public events | .0 | .0 | .1 | .1 | .2 | .1 | .1 |
| | Had someone else buy it | .4 | 1.0 | 1.9 | 4.4 | 7.5 | 10.9 | 14.4 |
| | Liquor stores/convenience stores/supermarkets | .0 | .1 | .1 | .4 | .5 | .8 | 1.4 |
| | Received from person 21 or older | 1.1 | 1.4 | 2.3 | 3.2 | 5.2 | 6.3 | 9.1 |
| | Took it from a store | .2 | .3 | .3 | .5 | .5 | .6 | .5 |
| | Family members | 2.2 | 3.1 | 4.3 | 4.6 | 4.3 | 3.9 | 4.0 |
| | Other ways | 2.1 | 3.8 | 5.7 | 7.8 | 8.6 | 8.2 | 8.0 |

**Most Important Reasons for Drinking
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|---|-----------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Did not drink alcoholic beverages | No answer | | 21.3 | 27.9 | 35.3 | 44.3 | 51.9 | 56.3 | 62.9 |
| | Yes | | 78.7 | 72.1 | 64.7 | 55.7 | 48.1 | 43.7 | 37.1 |
| To experiment | No answer | | 87.8 | 84.1 | 79.5 | 79.1 | 77.8 | 80.2 | 80.0 |
| | Yes | | 12.2 | 15.9 | 20.5 | 20.9 | 22.2 | 19.8 | 20.0 |
| To relax or relieve tension | No answer | | 98.7 | 96.9 | 93.8 | 89.7 | 85.1 | 81.0 | 77.3 |
| | Yes | | 1.3 | 3.1 | 6.2 | 10.3 | 14.9 | 19.0 | 22.7 |
| To feel good or get high | No answer | | 98.9 | 97.5 | 95.6 | 92.2 | 88.1 | 85.2 | 82.0 |
| | Yes | | 1.1 | 2.5 | 4.4 | 7.8 | 11.9 | 14.8 | 18.0 |
| To seek deeper insights and understanding | No answer | | 99.5 | 99.2 | 98.6 | 97.5 | 97.1 | 96.3 | 95.4 |
| | Yes | | .5 | .8 | 1.4 | 2.5 | 2.9 | 3.7 | 4.6 |
| To have a good time with friends | No answer | | 98.0 | 95.1 | 90.8 | 82.1 | 74.0 | 67.7 | 60.9 |
| | Yes | | 2.0 | 4.9 | 9.2 | 17.9 | 26.0 | 32.3 | 39.1 |
| To fit in with a group I like | No answer | | 99.1 | 98.6 | 98.1 | 98.0 | 97.4 | 97.0 | 97.0 |
| | Yes | | .9 | 1.4 | 1.9 | 2.0 | 2.6 | 3.0 | 3.0 |
| To get away from my problems | No answer | | 98.4 | 97.2 | 95.1 | 93.5 | 91.5 | 89.9 | 89.3 |
| | Yes | | 1.6 | 2.8 | 4.9 | 6.5 | 8.5 | 10.1 | 10.7 |
| Because of boredom | No answer | | 98.1 | 96.3 | 93.9 | 92.1 | 89.1 | 88.0 | 87.1 |
| | Yes | | 1.9 | 3.7 | 6.1 | 7.9 | 10.9 | 12.0 | 12.9 |
| Because of anger | No answer | | 98.6 | 97.8 | 96.1 | 95.2 | 94.2 | 93.4 | 93.1 |
| | Yes | | 1.4 | 2.2 | 3.9 | 4.8 | 5.8 | 6.6 | 6.9 |
| To get through the day | No answer | | 99.4 | 99.0 | 98.4 | 97.8 | 97.6 | 97.1 | 97.4 |
| | Yes | | .6 | 1.0 | 1.6 | 2.2 | 2.4 | 2.9 | 2.6 |
| To increase the effects of other drugs | No answer | | 99.7 | 99.4 | 98.9 | 98.1 | 97.4 | 97.0 | 96.4 |
| | Yes | | .3 | .6 | 1.1 | 1.9 | 2.6 | 3.0 | 3.6 |
| To decrease the effects of other drugs | No answer | | 99.8 | 99.7 | 99.6 | 99.4 | 99.6 | 99.6 | 99.3 |
| | Yes | | .2 | .3 | .4 | .6 | .4 | .4 | .7 |
| To get to sleep | No answer | | 99.2 | 99.0 | 98.5 | 97.7 | 97.4 | 96.7 | 95.8 |
| | Yes | | .8 | 1.0 | 1.5 | 2.3 | 2.6 | 3.3 | 4.2 |
| Because it tastes good | No answer | | 97.0 | 94.6 | 92.2 | 88.5 | 85.8 | 83.7 | 81.8 |
| | Yes | | 3.0 | 5.4 | 7.8 | 11.5 | 14.2 | 16.3 | 18.2 |
| Because I am hooked | No answer | | 99.4 | 99.3 | 99.1 | 98.8 | 98.8 | 98.5 | 98.2 |
| | Yes | | .6 | .7 | .9 | 1.2 | 1.2 | 1.5 | 1.8 |

**Consequences of Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-----------------------|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Had a hangover | No Answer | 4.6 | 4.2 | 3.7 | 4.4 | 4.1 | 4.4 | 4.0 |
| | Never | 91.2 | 87.2 | 82.3 | 74.4 | 68.6 | 64.3 | 58.5 |
| | Once | 2.1 | 3.7 | 5.6 | 7.7 | 9.4 | 10.5 | 10.6 |
| | Twice | .8 | 1.6 | 2.8 | 4.4 | 5.3 | 6.0 | 7.1 |
| | 3-5 times | .5 | 1.4 | 2.4 | 3.9 | 5.7 | 6.2 | 8.4 |
| | 6-10 times | .2 | .5 | 1.0 | 1.7 | 2.4 | 2.8 | 4.3 |
| | 11 or more times | .5 | 1.3 | 2.3 | 3.5 | 4.6 | 5.9 | 7.1 |
| | | | | | | | | |
| Had nausea or vomited | No Answer | 5.0 | 4.6 | 4.2 | 4.9 | 4.6 | 4.7 | 4.2 |
| | Never | 91.2 | 88.3 | 85.3 | 79.2 | 73.3 | 67.6 | 61.9 |
| | Once | 2.1 | 3.7 | 5.1 | 7.4 | 9.6 | 11.5 | 12.4 |
| | Twice | .7 | 1.3 | 2.0 | 2.9 | 4.5 | 5.2 | 7.3 |
| | 3-5 times | .4 | .9 | 1.8 | 2.9 | 4.2 | 5.7 | 7.3 |
| | 6-10 times | .2 | .4 | .6 | 1.1 | 1.8 | 2.3 | 3.2 |
| | 11 or more times | .4 | .8 | 1.1 | 1.6 | 2.1 | 3.0 | 3.8 |
| | | | | | | | | |
| Had a memory loss | No Answer | 5.3 | 5.0 | 4.3 | 5.1 | 4.8 | 5.0 | 4.5 |
| | Never | 92.2 | 90.5 | 88.5 | 83.1 | 78.1 | 74.3 | 70.2 |
| | Once | 1.4 | 2.0 | 3.2 | 4.5 | 6.5 | 6.9 | 8.2 |
| | Twice | .4 | .8 | 1.4 | 2.2 | 3.1 | 3.7 | 4.4 |
| | 3-5 times | .2 | .6 | 1.0 | 2.0 | 3.0 | 4.3 | 5.6 |
| | 6-10 times | .1 | .3 | .5 | 1.0 | 1.6 | 2.3 | 2.5 |
| | 11 or more times | .4 | .8 | 1.1 | 2.1 | 2.9 | 3.5 | 4.6 |
| | | | | | | | | |
| Poor on school test | No Answer | 5.3 | 4.8 | 4.3 | 5.2 | 4.9 | 5.0 | 4.5 |
| | Never | 91.3 | 90.7 | 89.7 | 87.5 | 87.3 | 86.9 | 86.9 |
| | Once | 1.5 | 1.6 | 2.0 | 2.3 | 2.5 | 2.4 | 2.5 |
| | Twice | .6 | .9 | 1.1 | 1.2 | 1.2 | 1.3 | 1.6 |
| | 3-5 times | .5 | .7 | 1.2 | 1.4 | 1.4 | 1.8 | 1.9 |
| | 6-10 times | .2 | .3 | .4 | .5 | .8 | .9 | .7 |
| | 11 or more times | .6 | .9 | 1.2 | 1.9 | 1.8 | 1.9 | 1.9 |
| | | | | | | | | |
| Missed school | No Answer | 5.5 | 4.9 | 4.5 | 5.2 | 5.0 | 4.9 | 4.5 |
| | Never | 91.5 | 91.5 | 90.5 | 88.9 | 88.9 | 88.6 | 86.8 |
| | Once | .9 | 1.2 | 1.7 | 1.7 | 1.9 | 1.9 | 2.4 |
| | Twice | .6 | .7 | 1.0 | 1.1 | 1.2 | 1.2 | 1.7 |
| | 3-5 times | .7 | .7 | 1.0 | 1.2 | 1.2 | 1.3 | 1.7 |
| | 6-10 times | .3 | .3 | .5 | .7 | .7 | .7 | .9 |
| | 11 or more times | .5 | .7 | .9 | 1.3 | 1.2 | 1.4 | 2.0 |
| | | | | | | | | |

**Consequences of Alcohol, Tobacco, and Other Drug Use
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Got into a fight | No Answer | 5.4 | 4.9 | 4.5 | 5.2 | 5.0 | 5.0 | 4.6 |
| | Never | 89.4 | 88.3 | 86.9 | 84.6 | 82.9 | 82.0 | 80.8 |
| | Once | 2.3 | 2.6 | 3.3 | 3.5 | 4.5 | 4.5 | 5.2 |
| | Twice | .8 | 1.2 | 1.5 | 2.0 | 2.5 | 2.6 | 3.2 |
| | 3-5 times | .7 | 1.2 | 1.6 | 1.8 | 2.2 | 2.6 | 2.9 |
| | 6-10 times | .4 | .5 | .7 | 1.1 | 1.0 | 1.3 | 1.2 |
| | 11 or more times | 1.0 | 1.4 | 1.6 | 1.9 | 1.9 | 2.1 | 2.1 |
| | | | | | | | | |
| Damaged property | No Answer | 5.5 | 5.1 | 4.4 | 5.2 | 5.0 | 5.0 | 4.6 |
| | Never | 92.6 | 91.9 | 91.4 | 89.4 | 88.9 | 88.4 | 88.0 |
| | Once | .8 | 1.1 | 1.5 | 1.8 | 2.0 | 2.0 | 2.5 |
| | Twice | .3 | .5 | .7 | .9 | 1.0 | 1.4 | 1.4 |
| | 3-5 times | .3 | .5 | .7 | 1.0 | 1.2 | 1.3 | 1.4 |
| | 6-10 times | .1 | .3 | .4 | .5 | .6 | .5 | .6 |
| | 11 or more times | .4 | .6 | .9 | 1.3 | 1.2 | 1.3 | 1.6 |
| | | | | | | | | |
| Driven by someone under the influence | No Answer | 4.1 | 3.8 | 3.3 | 3.8 | 3.6 | 3.7 | 3.6 |
| | Yes | 19.7 | 22.9 | 29.8 | 35.8 | 42.4 | 47.2 | 51.5 |
| | No | 76.2 | 73.3 | 66.9 | 60.4 | 54.0 | 49.1 | 44.9 |
| Used alcohol or drugs to relax or fit in | No Answer | 4.7 | 4.6 | 3.9 | 4.4 | 4.3 | 4.3 | 4.1 |
| | Yes | 3.2 | 5.7 | 9.2 | 13.9 | 18.5 | 21.5 | 25.1 |
| | No | 92.2 | 89.7 | 86.9 | 81.6 | 77.2 | 74.3 | 70.8 |
| Used alcohol or drugs alone | No Answer | 5.3 | 4.9 | 4.2 | 4.9 | 4.4 | 4.4 | 4.1 |
| | Yes | 3.7 | 8.0 | 12.2 | 18.2 | 22.8 | 24.4 | 26.8 |
| | No | 91.1 | 87.2 | 83.6 | 76.9 | 72.8 | 71.2 | 69.1 |
| Forgot things you did while high | No Answer | 8.1 | 7.3 | 5.9 | 6.4 | 5.5 | 5.0 | 4.8 |
| | Yes | 2.4 | 4.6 | 7.6 | 11.8 | 16.6 | 20.2 | 24.0 |
| | No | 89.5 | 88.1 | 86.5 | 81.8 | 77.9 | 74.8 | 71.2 |
| Had been told to cut down | No Answer | 9.7 | 8.6 | 6.6 | 7.2 | 5.7 | 5.4 | 4.9 |
| | Yes | 2.5 | 3.6 | 5.4 | 7.5 | 9.8 | 10.5 | 11.5 |
| | No | 87.7 | 87.9 | 88.1 | 85.3 | 84.4 | 84.1 | 83.6 |
| Got into trouble | No Answer | 9.9 | 8.6 | 6.4 | 7.1 | 5.7 | 5.3 | 4.9 |
| | Yes | 3.2 | 4.5 | 6.7 | 9.9 | 11.8 | 13.4 | 14.8 |
| | No | 87.0 | 86.9 | 86.9 | 83.0 | 82.5 | 81.3 | 80.3 |

**Gambling during the Last Twelve Months
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Played cards | No Answer | 3.7 | 3.2 | 3.3 | 4.1 | 3.9 | 3.9 | 3.9 |
| | Not at all | 76.9 | 72.4 | 68.9 | 66.4 | 66.4 | 65.0 | 63.3 |
| | Less than once a month | 9.7 | 11.8 | 13.6 | 14.4 | 14.8 | 15.3 | 16.4 |
| | About once a month | 5.1 | 6.8 | 7.9 | 8.1 | 8.3 | 9.0 | 9.5 |
| | About once a week | 3.2 | 3.8 | 4.6 | 4.9 | 4.5 | 4.6 | 4.8 |
| | Daily | 1.4 | 2.1 | 1.7 | 2.2 | 2.0 | 2.2 | 2.1 |
| Bet on games | No Answer | 3.9 | 3.6 | 3.6 | 4.2 | 4.1 | 4.0 | 4.1 |
| | Not at all | 76.1 | 71.0 | 67.7 | 64.8 | 65.0 | 64.4 | 63.6 |
| | Less than once a month | 9.0 | 10.9 | 11.5 | 12.0 | 12.1 | 11.9 | 13.0 |
| | About once a month | 5.1 | 6.4 | 7.6 | 8.5 | 8.4 | 8.9 | 8.9 |
| | About once a week | 3.3 | 4.4 | 5.5 | 6.1 | 6.0 | 6.3 | 6.3 |
| | Daily | 2.6 | 3.7 | 4.1 | 4.4 | 4.3 | 4.4 | 4.1 |
| Bet on sports | No Answer | 4.1 | 3.5 | 3.8 | 4.4 | 4.3 | 4.2 | 4.2 |
| | Not at all | 76.8 | 72.3 | 69.8 | 67.8 | 69.1 | 70.7 | 70.3 |
| | Less than once a month | 10.5 | 12.7 | 13.8 | 14.4 | 13.8 | 13.2 | 13.8 |
| | About once a month | 3.7 | 5.5 | 6.0 | 6.6 | 6.2 | 6.0 | 6.2 |
| | About once a week | 2.9 | 3.3 | 3.9 | 3.7 | 3.7 | 3.6 | 3.0 |
| | Daily | 1.9 | 2.7 | 2.7 | 3.2 | 2.8 | 2.4 | 2.4 |
| Bought lottery tickets | No Answer | 4.2 | 3.9 | 3.7 | 4.5 | 4.5 | 4.5 | 4.3 |
| | Not at all | 79.5 | 77.5 | 77.3 | 76.8 | 77.7 | 77.6 | 63.8 |
| | Less than once a month | 9.8 | 10.7 | 11.3 | 10.9 | 10.8 | 10.7 | 18.3 |
| | About once a month | 3.7 | 4.5 | 4.5 | 4.4 | 4.0 | 4.0 | 7.7 |
| | About once a week | 1.9 | 2.0 | 2.2 | 2.0 | 1.8 | 1.9 | 4.1 |
| | Daily | .9 | 1.3 | 1.0 | 1.3 | 1.2 | 1.4 | 1.8 |
| Gambled in a casino | No Answer | 4.7 | 3.9 | 4.0 | 4.6 | 4.4 | 4.3 | 4.1 |
| | Not at all | 94.4 | 94.4 | 94.3 | 93.2 | 93.4 | 93.0 | 92.0 |
| | Less than once a month | .4 | .8 | .8 | .7 | .9 | 1.1 | 1.9 |
| | About once a month | .1 | .2 | .3 | .3 | .3 | .5 | .6 |
| | About once a week | .1 | .2 | .2 | .2 | .2 | .3 | .4 |
| | Daily | .3 | .5 | .5 | .9 | .7 | .8 | 1.0 |
| Played online | No Answer | 4.5 | 3.7 | 3.8 | 4.5 | 4.5 | 4.4 | 4.1 |
| | Not at all | 84.0 | 82.9 | 81.7 | 82.1 | 83.4 | 84.2 | 84.8 |
| | Less than once a month | 3.3 | 3.9 | 4.2 | 3.5 | 3.6 | 3.1 | 3.3 |
| | About once a month | 1.6 | 2.0 | 2.2 | 2.2 | 2.1 | 2.2 | 2.1 |
| | About once a week | 2.3 | 2.6 | 2.9 | 2.7 | 2.3 | 2.2 | 2.3 |
| | Daily | 4.3 | 4.9 | 5.1 | 5.0 | 4.0 | 3.9 | 3.5 |
| Felt bad about the amount bet | No Answer | 11.9 | 11.2 | 10.7 | 10.4 | 10.5 | 9.7 | 9.0 |
| | Yes | 4.4 | 4.0 | 3.7 | 3.6 | 3.5 | 3.4 | 3.9 |
| | No | 19.2 | 22.7 | 25.9 | 28.2 | 28.8 | 29.0 | 32.8 |
| | I do not bet for money | 64.5 | 62.1 | 59.8 | 57.7 | 57.2 | 57.8 | 54.2 |
| Would like to stop betting but could not | No Answer | 12.9 | 11.8 | 11.1 | 10.9 | 10.8 | 9.8 | 9.2 |
| | Yes | 3.4 | 2.8 | 2.7 | 2.3 | 1.9 | 1.9 | 1.9 |
| | No | 19.6 | 23.4 | 26.3 | 28.7 | 29.7 | 29.8 | 34.3 |
| | I do not bet for money | 64.1 | 62.1 | 59.9 | 58.1 | 57.7 | 58.5 | 54.7 |

**Perceived Risk of Selected Substance Use Behavior
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|---------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Smoke 1+ pack cigarettes per day | No Answer | 4.5 | 4.1 | 3.6 | 4.9 | 4.3 | 4.2 | 4.5 |
| | No risk | 7.8 | 6.5 | 5.7 | 5.8 | 5.8 | 6.0 | 5.4 |
| | Slight risk | 10.5 | 8.3 | 8.9 | 8.8 | 7.8 | 8.4 | 8.4 |
| | Moderate risk | 24.9 | 24.0 | 26.1 | 25.4 | 24.8 | 26.0 | 24.7 |
| | Great risk | 52.2 | 57.1 | 55.6 | 55.1 | 57.2 | 55.4 | 57.0 |
| Smoke marijuana occasionally | No Answer | 4.7 | 4.3 | 3.9 | 5.0 | 4.4 | 4.4 | 4.6 |
| | No risk | 8.9 | 9.5 | 12.5 | 19.1 | 23.9 | 28.5 | 29.0 |
| | Slight risk | 10.7 | 12.9 | 17.8 | 21.8 | 24.6 | 25.8 | 27.8 |
| | Moderate risk | 27.5 | 28.1 | 27.6 | 24.9 | 22.8 | 20.4 | 19.7 |
| | Great risk | 48.2 | 45.2 | 38.2 | 29.1 | 24.3 | 21.0 | 18.9 |
| Smoke marijuana regularly | No Answer | 5.5 | 4.9 | 4.6 | 5.8 | 4.9 | 4.9 | 5.1 |
| | No risk | 8.3 | 7.8 | 9.0 | 12.8 | 14.6 | 16.6 | 17.2 |
| | Slight risk | 5.0 | 5.8 | 8.6 | 12.9 | 16.3 | 18.6 | 19.8 |
| | Moderate risk | 12.9 | 13.6 | 17.1 | 20.4 | 21.7 | 22.7 | 23.5 |
| | Great risk | 68.4 | 68.0 | 60.7 | 48.1 | 42.6 | 37.2 | 34.4 |
| Occasionally consume 1-2 alcoholic drinks | No Answer | 5.4 | 4.9 | 4.4 | 5.6 | 4.8 | 4.8 | 4.9 |
| | No risk | 21.7 | 22.0 | 24.8 | 29.9 | 32.3 | 37.0 | 39.1 |
| | Slight risk | 37.0 | 38.1 | 38.2 | 35.3 | 35.7 | 33.6 | 34.3 |
| | Moderate risk | 18.3 | 17.3 | 16.1 | 14.5 | 13.8 | 12.7 | 10.8 |
| | Great risk | 17.5 | 17.8 | 16.6 | 14.7 | 13.4 | 12.0 | 10.9 |
| Binge drink weekly | No Answer | 5.1 | 4.5 | 4.2 | 5.2 | 4.7 | 4.7 | 4.9 |
| | No risk | 9.7 | 8.5 | 7.9 | 8.6 | 9.0 | 10.0 | 9.5 |
| | Slight risk | 14.5 | 13.4 | 14.6 | 17.2 | 17.2 | 18.5 | 21.1 |
| | Moderate risk | 28.4 | 30.4 | 31.6 | 31.2 | 31.8 | 33.0 | 33.5 |
| | Great risk | 42.3 | 43.2 | 41.8 | 37.9 | 37.4 | 33.8 | 31.0 |

**Perceived Peer Approval of Using Selected Substances
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Smoke 1+ pack cigarettes per day | No Answer | 6.6 | 5.8 | 4.8 | 5.8 | 5.3 | 5.2 | 5.2 |
| | Strongly approve | 2.6 | 3.0 | 2.7 | 3.0 | 3.3 | 3.4 | 3.7 |
| | Approve | 1.2 | 1.9 | 3.0 | 4.8 | 5.8 | 7.1 | 8.2 |
| | Do not know | 14.5 | 15.9 | 18.7 | 20.7 | 21.1 | 21.5 | 21.9 |
| | Disapprove | 11.1 | 12.7 | 15.1 | 17.8 | 18.7 | 20.0 | 21.5 |
| | Strongly disapprove | 63.9 | 60.7 | 55.7 | 47.9 | 45.8 | 42.9 | 39.5 |
| Smoke marijuana occasionally | No Answer | 6.7 | 5.9 | 4.9 | 5.9 | 5.5 | 5.3 | 5.4 |
| | Strongly approve | 2.9 | 3.6 | 4.2 | 6.0 | 7.5 | 9.2 | 9.0 |
| | Approve | 1.7 | 3.2 | 7.0 | 11.4 | 15.6 | 17.8 | 19.0 |
| | Do not know | 12.6 | 14.7 | 17.9 | 20.7 | 20.7 | 21.4 | 21.6 |
| | Disapprove | 8.2 | 9.5 | 11.4 | 13.8 | 13.9 | 13.6 | 14.5 |
| | Strongly disapprove | 67.9 | 63.0 | 54.6 | 42.2 | 36.8 | 32.6 | 30.4 |
| Smoke marijuana regularly | No Answer | 7.0 | 6.1 | 5.2 | 6.2 | 5.8 | 5.5 | 5.5 |
| | Strongly approve | 3.0 | 3.8 | 4.3 | 6.0 | 7.7 | 9.0 | 9.0 |
| | Approve | 1.1 | 2.2 | 4.8 | 7.7 | 10.4 | 11.5 | 12.0 |
| | Do not know | 12.1 | 13.7 | 16.9 | 19.5 | 19.8 | 20.4 | 20.0 |
| | Disapprove | 6.1 | 7.1 | 8.9 | 11.8 | 12.3 | 13.5 | 14.6 |
| | Strongly disapprove | 70.6 | 67.1 | 59.9 | 48.9 | 44.1 | 40.0 | 38.8 |
| Occasionally consume 1-2 alcoholic drinks | No Answer | 7.0 | 6.2 | 5.1 | 6.0 | 5.6 | 5.4 | 5.5 |
| | Strongly approve | 3.0 | 3.6 | 4.0 | 5.3 | 7.1 | 8.8 | 10.3 |
| | Approve | 4.0 | 6.2 | 10.8 | 17.2 | 21.9 | 25.6 | 29.8 |
| | Do not know | 18.0 | 20.1 | 24.4 | 26.8 | 27.0 | 26.6 | 25.6 |
| | Disapprove | 12.4 | 13.1 | 13.2 | 13.3 | 12.6 | 11.8 | 10.2 |
| | Strongly disapprove | 55.7 | 50.9 | 42.5 | 31.4 | 25.8 | 21.8 | 18.6 |
| Binge drink weekly | No Answer | 7.1 | 6.2 | 5.2 | 6.2 | 5.7 | 5.5 | 5.6 |
| | Strongly approve | 2.9 | 3.4 | 3.5 | 4.6 | 5.8 | 6.9 | 7.8 |
| | Approve | 1.4 | 2.4 | 4.5 | 7.5 | 10.3 | 12.7 | 15.0 |
| | Do not know | 13.5 | 15.5 | 19.5 | 23.1 | 24.0 | 24.1 | 24.5 |
| | Disapprove | 8.9 | 10.0 | 12.2 | 15.0 | 15.7 | 16.2 | 16.1 |
| | Strongly disapprove | 66.3 | 62.5 | 55.0 | 43.6 | 38.6 | 34.6 | 30.9 |

**Parental Attitudes toward Drug Use and Antisocial Behavior
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Parents feel about alcohol | No Answer | 5.5 | 5.0 | 4.4 | 5.7 | 5.4 | 5.3 | 5.6 |
| | Very wrong | 82.5 | 79.7 | 75.4 | 67.1 | 61.8 | 55.4 | 47.0 |
| | Wrong | 7.3 | 8.7 | 11.2 | 14.3 | 16.7 | 17.6 | 19.9 |
| | A little bit wrong | 3.3 | 4.4 | 6.2 | 9.3 | 11.9 | 15.3 | 19.0 |
| | Not at all wrong | 1.4 | 2.1 | 2.8 | 3.5 | 4.2 | 6.4 | 8.6 |
| Parents feel about cigarettes | No Answer | 5.5 | 5.0 | 4.5 | 5.9 | 5.5 | 5.4 | 5.6 |
| | Very wrong | 87.2 | 86.1 | 83.1 | 77.7 | 74.1 | 67.5 | 58.0 |
| | Wrong | 5.0 | 6.0 | 7.8 | 10.3 | 12.6 | 15.5 | 18.8 |
| | A little bit wrong | 1.1 | 1.4 | 2.5 | 3.3 | 4.6 | 7.0 | 10.2 |
| | Not at all wrong | 1.1 | 1.5 | 2.0 | 2.7 | 3.2 | 4.7 | 7.3 |
| Parents feel about marijuana | No Answer | 5.8 | 5.1 | 4.6 | 5.9 | 5.6 | 5.5 | 5.7 |
| | Very wrong | 90.1 | 88.5 | 85.2 | 79.2 | 75.9 | 71.1 | 67.5 |
| | Wrong | 2.3 | 3.3 | 5.5 | 8.0 | 9.9 | 12.1 | 14.4 |
| | A little bit wrong | .6 | 1.3 | 2.4 | 3.8 | 5.0 | 6.3 | 7.4 |
| | Not at all wrong | 1.2 | 1.7 | 2.4 | 3.2 | 3.6 | 5.0 | 5.0 |
| Parents feel about stealing | No Answer | 5.9 | 5.2 | 4.6 | 6.1 | 5.6 | 5.5 | 5.8 |
| | Very wrong | 80.8 | 79.8 | 78.3 | 75.0 | 75.0 | 73.6 | 73.8 |
| | Wrong | 10.1 | 10.6 | 11.8 | 13.1 | 14.1 | 14.6 | 15.1 |
| | A little bit wrong | 2.0 | 2.6 | 3.1 | 3.6 | 3.1 | 3.8 | 3.1 |
| | Not at all wrong | 1.2 | 1.7 | 2.1 | 2.2 | 2.1 | 2.6 | 2.3 |
| Parents feel about graffiti | No Answer | 6.0 | 5.2 | 4.7 | 6.1 | 5.5 | 5.6 | 5.6 |
| | Very wrong | 83.5 | 82.9 | 79.5 | 76.5 | 74.8 | 72.3 | 71.3 |
| | Wrong | 6.9 | 7.7 | 10.0 | 11.3 | 13.5 | 14.6 | 15.7 |
| | A little bit wrong | 1.9 | 2.2 | 3.4 | 3.8 | 3.7 | 4.5 | 4.5 |
| | Not at all wrong | 1.7 | 1.9 | 2.4 | 2.4 | 2.5 | 3.0 | 3.0 |
| Parents feel about fighting | No Answer | 6.0 | 5.4 | 4.9 | 6.2 | 5.7 | 5.7 | 5.9 |
| | Very wrong | 64.1 | 61.2 | 55.8 | 53.0 | 51.1 | 50.1 | 49.1 |
| | Wrong | 18.0 | 18.8 | 20.6 | 21.4 | 23.1 | 23.3 | 25.7 |
| | A little bit wrong | 8.8 | 10.5 | 13.5 | 14.3 | 15.2 | 15.6 | 14.5 |
| | Not at all wrong | 3.1 | 4.0 | 5.2 | 5.1 | 5.0 | 5.2 | 4.7 |

**Family Management and Conflict
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Clear rules | No Answer | 4.5 | 4.5 | 4.3 | 6.3 | 6.4 | 6.8 | 6.9 |
| | YES! | 65.0 | 61.4 | 57.9 | 54.6 | 54.1 | 54.0 | 51.9 |
| | yes | 26.5 | 29.2 | 32.4 | 33.0 | 33.7 | 33.1 | 34.4 |
| | no | 2.7 | 3.3 | 3.8 | 4.2 | 4.2 | 4.3 | 4.7 |
| | NO! | 1.4 | 1.6 | 1.6 | 1.9 | 1.7 | 1.9 | 2.1 |
| Ask about homework | No Answer | 4.4 | 4.6 | 4.4 | 6.5 | 6.7 | 7.2 | 7.3 |
| | YES! | 68.9 | 62.5 | 55.8 | 49.1 | 42.6 | 37.9 | 30.8 |
| | yes | 21.6 | 24.9 | 29.5 | 31.2 | 33.8 | 34.2 | 34.9 |
| | no | 3.5 | 5.7 | 7.6 | 10.0 | 12.8 | 14.8 | 19.4 |
| | NO! | 1.6 | 2.2 | 2.7 | 3.1 | 4.1 | 5.9 | 7.6 |
| Parents know whereabouts | No Answer | 4.9 | 4.8 | 4.7 | 6.9 | 7.1 | 7.6 | 7.5 |
| | YES! | 71.8 | 66.9 | 60.6 | 53.3 | 49.3 | 45.2 | 38.5 |
| | yes | 17.1 | 20.9 | 26.0 | 30.3 | 33.1 | 35.2 | 38.0 |
| | no | 3.9 | 5.2 | 6.2 | 7.1 | 7.8 | 8.8 | 11.9 |
| | NO! | 2.2 | 2.2 | 2.4 | 2.4 | 2.7 | 3.3 | 4.1 |
| Clear rules about alcohol use | No Answer | 5.9 | 5.6 | 5.2 | 7.3 | 7.5 | 7.7 | 7.8 |
| | YES! | 79.5 | 76.5 | 69.8 | 61.3 | 57.4 | 53.5 | 47.6 |
| | yes | 9.5 | 11.8 | 17.0 | 21.7 | 24.5 | 26.2 | 30.6 |
| | no | 3.0 | 3.9 | 5.9 | 7.1 | 8.3 | 9.5 | 10.8 |
| | NO! | 2.2 | 2.2 | 2.2 | 2.5 | 2.3 | 3.0 | 3.2 |
| Parents ask before most family decisions made | No Answer | 8.7 | 7.4 | 6.4 | 8.6 | 8.7 | 8.8 | 8.6 |
| | YES! | 35.7 | 31.9 | 28.5 | 25.3 | 22.7 | 22.3 | 20.3 |
| | yes | 32.5 | 34.5 | 36.5 | 36.9 | 38.0 | 37.7 | 38.6 |
| | no | 13.5 | 15.8 | 18.7 | 20.0 | 21.2 | 21.7 | 23.3 |
| | NO! | 9.6 | 10.4 | 9.9 | 9.3 | 9.4 | 9.5 | 9.1 |
| Ask parents for help if problem | No Answer | 6.2 | 6.2 | 5.6 | 8.4 | 8.4 | 8.6 | 8.5 |
| | YES! | 66.0 | 57.6 | 50.4 | 43.8 | 40.9 | 39.9 | 38.7 |
| | yes | 19.1 | 23.9 | 28.3 | 30.7 | 32.3 | 33.9 | 35.1 |
| | no | 4.5 | 7.0 | 9.5 | 10.7 | 11.9 | 11.4 | 11.7 |
| | NO! | 4.1 | 5.3 | 6.3 | 6.5 | 6.5 | 6.2 | 6.0 |
| Parents give me chances for fun with them | No Answer | 6.6 | 6.7 | 6.0 | 8.7 | 8.8 | 8.7 | 8.8 |
| | YES! | 53.4 | 49.3 | 42.9 | 36.0 | 32.4 | 29.9 | 27.1 |
| | yes | 27.4 | 28.4 | 32.9 | 35.2 | 36.4 | 37.6 | 37.5 |
| | no | 8.9 | 11.0 | 13.1 | 14.5 | 16.8 | 17.9 | 20.2 |
| | NO! | 3.8 | 4.7 | 5.1 | 5.5 | 5.6 | 5.9 | 6.3 |

**Family Management and Conflict
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Parents know when home late | No Answer | 7.7 | 7.5 | 6.4 | 9.1 | 9.1 | 9.1 | 9.5 |
| | YES! | 65.9 | 61.1 | 56.5 | 48.3 | 45.7 | 43.1 | 35.3 |
| | yes | 18.6 | 22.3 | 26.8 | 30.4 | 32.2 | 32.5 | 35.6 |
| | no | 5.5 | 6.6 | 7.6 | 9.4 | 10.1 | 11.5 | 14.6 |
| | NO! | 2.3 | 2.5 | 2.6 | 2.9 | 2.9 | 3.7 | 5.0 |
| Parents catch you drinking | No Answer | 7.8 | 7.1 | 6.6 | 9.4 | 9.1 | 9.3 | 9.2 |
| | YES! | 67.7 | 57.2 | 45.2 | 33.1 | 26.2 | 23.2 | 18.4 |
| | yes | 13.5 | 18.2 | 22.4 | 22.8 | 22.7 | 20.4 | 19.5 |
| | no | 6.6 | 11.5 | 18.1 | 24.9 | 30.4 | 33.0 | 36.7 |
| | NO! | 4.4 | 5.9 | 7.7 | 9.7 | 11.6 | 14.1 | 16.1 |
| Parents catch you with gun | No Answer | 8.0 | 7.2 | 6.6 | 9.7 | 9.4 | 9.3 | 9.3 |
| | YES! | 74.2 | 67.9 | 59.5 | 50.1 | 45.2 | 41.1 | 35.1 |
| | yes | 10.4 | 13.9 | 18.8 | 21.6 | 22.9 | 24.3 | 25.2 |
| | no | 4.1 | 6.5 | 9.7 | 12.4 | 15.7 | 17.4 | 21.1 |
| | NO! | 3.3 | 4.4 | 5.4 | 6.1 | 6.9 | 7.9 | 9.2 |
| Parents catch you skip school | No Answer | 7.0 | 6.8 | 6.1 | 9.1 | 8.6 | 9.0 | 9.1 |
| | YES! | 76.9 | 71.7 | 66.3 | 57.1 | 53.1 | 48.4 | 41.5 |
| | yes | 10.7 | 14.1 | 18.3 | 23.1 | 24.7 | 27.7 | 29.7 |
| | no | 3.0 | 4.3 | 6.1 | 7.3 | 9.4 | 10.6 | 13.7 |
| | NO! | 2.4 | 3.1 | 3.3 | 3.4 | 4.0 | 4.4 | 6.0 |
| Family argue repetitively | No Answer | 7.0 | 6.1 | 5.8 | 7.8 | 7.9 | 8.0 | 8.1 |
| | YES! | 17.4 | 19.9 | 20.3 | 20.9 | 20.9 | 20.0 | 18.5 |
| | yes | 18.9 | 20.0 | 23.1 | 24.2 | 25.7 | 25.0 | 24.9 |
| | no | 31.9 | 33.6 | 34.1 | 33.7 | 32.6 | 34.7 | 35.0 |
| | NO! | 24.8 | 20.4 | 16.7 | 13.4 | 13.0 | 12.3 | 13.4 |
| Family have serious arguments | No Answer | 7.3 | 6.6 | 6.1 | 8.1 | 8.2 | 8.4 | 8.6 |
| | YES! | 15.0 | 15.8 | 16.5 | 16.6 | 16.3 | 15.9 | 13.6 |
| | yes | 14.8 | 15.7 | 17.5 | 18.8 | 19.2 | 19.4 | 19.1 |
| | no | 30.9 | 33.0 | 35.2 | 35.8 | 36.4 | 37.0 | 38.7 |
| | NO! | 32.0 | 28.8 | 24.7 | 20.7 | 19.9 | 19.4 | 20.0 |
| Family insult each other | No Answer | 8.4 | 7.4 | 6.8 | 8.7 | 9.0 | 8.8 | 9.3 |
| | YES! | 14.4 | 15.9 | 16.2 | 16.6 | 15.9 | 15.4 | 13.2 |
| | yes | 16.4 | 16.4 | 18.8 | 19.5 | 19.7 | 19.2 | 18.7 |
| | no | 26.4 | 29.2 | 30.6 | 31.2 | 32.4 | 33.4 | 34.9 |
| | NO! | 34.3 | 31.1 | 27.6 | 24.0 | 22.9 | 23.2 | 24.0 |

Interaction with Antisocial Peers
by Indiana Male 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Number of best friends suspended | No Answer | 6.5 | 6.6 | 6.1 | 9.3 | 8.9 | 8.8 | 9.0 |
| | None of my friends | 64.4 | 60.4 | 58.1 | 56.8 | 60.6 | 64.2 | 66.2 |
| | 1 of my friends | 17.9 | 18.6 | 19.1 | 17.9 | 16.5 | 14.8 | 13.8 |
| | 2 of my friends | 5.7 | 7.3 | 8.5 | 7.9 | 7.0 | 6.1 | 5.1 |
| | 3 of my friends | 1.9 | 2.5 | 2.8 | 2.4 | 2.3 | 2.1 | 1.9 |
| | 4 of my friends | 3.6 | 4.6 | 5.5 | 5.6 | 4.7 | 4.0 | 3.9 |
| Number of best friends carry guns | No Answer | 6.3 | 6.6 | 6.1 | 9.5 | 8.9 | 8.9 | 9.2 |
| | None of my friends | 88.7 | 87.0 | 85.4 | 80.9 | 80.3 | 79.6 | 77.0 |
| | 1 of my friends | 3.1 | 3.6 | 4.6 | 5.3 | 5.9 | 5.9 | 7.8 |
| | 2 of my friends | .9 | 1.0 | 1.5 | 1.7 | 2.0 | 2.3 | 2.7 |
| | 3 of my friends | .3 | .4 | .6 | .7 | .7 | 1.0 | .9 |
| | 4 of my friends | .7 | 1.3 | 1.7 | 1.9 | 2.3 | 2.2 | 2.5 |
| Number of best friends sold drugs | No Answer | 6.5 | 6.7 | 6.3 | 9.7 | 9.2 | 9.2 | 9.5 |
| | None of my friends | 89.5 | 84.3 | 79.6 | 69.3 | 65.8 | 64.6 | 62.1 |
| | 1 of my friends | 2.5 | 4.6 | 7.4 | 9.5 | 11.3 | 11.4 | 12.8 |
| | 2 of my friends | .7 | 1.7 | 2.7 | 5.0 | 6.1 | 6.6 | 7.0 |
| | 3 of my friends | .3 | .7 | 1.3 | 2.1 | 2.6 | 2.6 | 3.1 |
| | 4 of my friends | .6 | 1.9 | 2.7 | 4.5 | 5.1 | 5.6 | 5.4 |
| Number of best friends stolen vehicle | No Answer | 6.3 | 6.7 | 6.3 | 9.5 | 9.2 | 9.2 | 9.3 |
| | None of my friends | 90.6 | 88.7 | 87.5 | 82.6 | 82.4 | 83.6 | 84.0 |
| | 1 of my friends | 1.9 | 2.6 | 3.4 | 4.1 | 4.6 | 3.3 | 3.2 |
| | 2 of my friends | .4 | .7 | 1.0 | 1.3 | 1.6 | 1.6 | 1.4 |
| | 3 of my friends | .2 | .3 | .5 | .8 | .6 | .7 | .6 |
| | 4 of my friends | .5 | 1.0 | 1.4 | 1.7 | 1.7 | 1.6 | 1.5 |
| Number of best friends arrested | No Answer | 6.6 | 6.8 | 6.2 | 9.7 | 9.3 | 9.2 | 9.4 |
| | None of my friends | 85.2 | 81.6 | 78.1 | 72.4 | 71.9 | 71.4 | 69.7 |
| | 1 of my friends | 5.6 | 7.0 | 9.1 | 9.6 | 9.9 | 10.3 | 10.9 |
| | 2 of my friends | 1.3 | 1.8 | 2.9 | 3.7 | 4.0 | 4.1 | 4.8 |
| | 3 of my friends | .5 | 1.0 | 1.3 | 1.5 | 1.8 | 1.8 | 1.7 |
| | 4 of my friends | .8 | 1.8 | 2.4 | 3.2 | 3.1 | 3.2 | 3.5 |
| Number of best friends dropped out of school | No Answer | 6.7 | 6.9 | 6.4 | 9.9 | 9.2 | 9.3 | 9.3 |
| | None of my friends | 88.7 | 87.5 | 87.0 | 80.6 | 78.5 | 76.7 | 75.7 |
| | 1 of my friends | 3.2 | 3.4 | 3.8 | 5.4 | 7.7 | 8.1 | 8.5 |
| | 2 of my friends | .6 | .8 | 1.1 | 1.6 | 2.1 | 2.8 | 3.0 |
| | 3 of my friends | .2 | .3 | .4 | .8 | .9 | 1.1 | 1.4 |
| | 4 of my friends | .4 | 1.1 | 1.3 | 1.7 | 1.6 | 2.0 | 2.1 |
| Number of best friends been members of gangs | No Answer | 7.4 | 7.4 | 6.9 | 10.3 | 9.8 | 9.8 | 9.8 |
| | None of my friends | 83.8 | 81.5 | 80.1 | 77.2 | 78.4 | 79.6 | 80.4 |
| | 1 of my friends | 4.2 | 4.9 | 5.6 | 4.8 | 4.7 | 4.0 | 3.8 |
| | 2 of my friends | 1.7 | 2.0 | 2.2 | 2.2 | 2.2 | 2.0 | 2.0 |
| | 3 of my friends | .9 | .8 | 1.2 | 1.4 | 1.1 | 1.0 | .9 |
| | 4 of my friends | 2.0 | 3.4 | 4.0 | 4.0 | 3.8 | 3.6 | 3.1 |

**Interaction with Prosocial Peers
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Number of best friends participated in school activities | No Answer | 9.1 | 8.3 | 8.1 | 10.7 | 10.5 | 10.0 | 10.4 |
| | None of my friends | 26.4 | 26.1 | 25.9 | 25.0 | 24.0 | 24.8 | 23.6 |
| | 1 of my friends | 11.2 | 9.5 | 9.2 | 8.2 | 8.5 | 8.0 | 8.4 |
| | 2 of my friends | 14.4 | 13.2 | 12.8 | 11.9 | 12.5 | 12.2 | 12.1 |
| | 3 of my friends | 10.5 | 10.5 | 11.0 | 10.3 | 10.7 | 10.5 | 9.9 |
| | 4 of my friends | 28.4 | 32.4 | 33.1 | 34.0 | 33.9 | 34.5 | 35.7 |
| Number of best friends commitment to stay drug-free | No Answer | 10.2 | 9.2 | 8.6 | 11.6 | 10.8 | 10.5 | 10.5 |
| | None of my friends | 22.5 | 26.2 | 32.0 | 34.4 | 35.4 | 39.8 | 40.6 |
| | 1 of my friends | 4.5 | 5.4 | 7.2 | 9.2 | 11.0 | 11.3 | 11.9 |
| | 2 of my friends | 4.8 | 5.7 | 7.6 | 8.1 | 9.4 | 9.2 | 9.7 |
| | 3 of my friends | 5.6 | 6.3 | 7.1 | 7.7 | 7.9 | 6.9 | 7.2 |
| | 4 of my friends | 52.4 | 47.2 | 37.5 | 29.0 | 25.5 | 22.3 | 20.1 |
| Number of best friends liked school | No Answer | 9.3 | 8.8 | 8.2 | 10.9 | 10.6 | 10.4 | 10.6 |
| | None of my friends | 31.8 | 35.0 | 38.1 | 39.5 | 39.8 | 40.7 | 40.0 |
| | 1 of my friends | 9.9 | 10.3 | 10.3 | 9.5 | 10.7 | 10.3 | 10.1 |
| | 2 of my friends | 13.1 | 12.6 | 12.9 | 12.6 | 12.8 | 12.9 | 13.0 |
| | 3 of my friends | 10.6 | 10.7 | 10.2 | 9.1 | 9.3 | 8.4 | 8.4 |
| | 4 of my friends | 25.2 | 22.6 | 20.3 | 18.3 | 16.8 | 17.3 | 17.9 |
| Number of best friends regularly attended religious activities | No Answer | 11.4 | 9.8 | 8.8 | 11.6 | 10.6 | 10.2 | 10.6 |
| | None of my friends | 24.6 | 26.2 | 28.2 | 28.9 | 30.1 | 32.9 | 32.6 |
| | 1 of my friends | 14.4 | 14.3 | 15.9 | 16.2 | 16.9 | 16.0 | 17.2 |
| | 2 of my friends | 16.8 | 16.4 | 17.6 | 15.8 | 16.4 | 15.8 | 16.6 |
| | 3 of my friends | 12.4 | 12.6 | 11.4 | 11.1 | 10.4 | 9.9 | 8.8 |
| | 4 of my friends | 20.3 | 20.7 | 18.1 | 16.4 | 15.5 | 15.0 | 14.2 |
| Number of best friends tried to do well in school | No Answer | 9.7 | 8.8 | 8.6 | 11.1 | 10.7 | 10.0 | 10.8 |
| | None of my friends | 9.6 | 11.9 | 12.9 | 13.5 | 13.1 | 14.6 | 13.1 |
| | 1 of my friends | 4.7 | 4.9 | 6.1 | 6.6 | 6.1 | 5.8 | 6.0 |
| | 2 of my friends | 6.1 | 7.3 | 8.7 | 9.1 | 10.0 | 11.0 | 11.4 |
| | 3 of my friends | 9.9 | 12.0 | 13.4 | 13.0 | 14.3 | 13.7 | 14.6 |
| | 4 of my friends | 59.9 | 55.1 | 50.4 | 46.8 | 45.8 | 44.8 | 44.1 |

**Rewards for Antisocial Involvement
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---------------------------------|--------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cool if smoke cigarettes | No Answer | 8.1 | 8.0 | 7.2 | 10.8 | 10.1 | 9.8 | 9.9 |
| | No or very little chance | 76.3 | 71.1 | 67.2 | 60.9 | 60.1 | 59.9 | 58.1 |
| | Little chance | 6.8 | 9.2 | 12.0 | 12.6 | 14.4 | 14.6 | 15.6 |
| | Some chance | 4.0 | 5.9 | 7.8 | 9.0 | 9.5 | 9.2 | 10.3 |
| | Pretty good chance | 2.0 | 2.7 | 2.9 | 3.5 | 3.0 | 3.1 | 2.8 |
| | Very good chance | 2.8 | 3.1 | 2.9 | 3.1 | 2.8 | 3.5 | 3.3 |
| Cool if drink alcohol regularly | No Answer | 8.3 | 8.2 | 7.4 | 11.0 | 10.3 | 10.1 | 10.2 |
| | No or very little chance | 76.1 | 69.4 | 63.3 | 53.6 | 49.0 | 47.7 | 44.7 |
| | Little chance | 6.8 | 9.7 | 12.4 | 13.3 | 14.4 | 14.1 | 14.3 |
| | Some chance | 3.6 | 6.1 | 9.2 | 11.6 | 14.3 | 13.8 | 15.6 |
| | Pretty good chance | 2.2 | 3.2 | 4.2 | 6.3 | 7.6 | 8.9 | 9.0 |
| | Very good chance | 3.0 | 3.4 | 3.5 | 4.2 | 4.5 | 5.4 | 6.2 |
| Cool if smoke marijuana | No Answer | 8.3 | 8.2 | 7.4 | 11.0 | 10.3 | 9.9 | 10.2 |
| | No or very little chance | 79.0 | 72.8 | 65.8 | 55.9 | 51.9 | 51.4 | 49.5 |
| | Little chance | 4.4 | 6.6 | 9.2 | 10.2 | 12.3 | 12.3 | 13.9 |
| | Some chance | 2.6 | 4.6 | 7.6 | 10.1 | 12.0 | 12.3 | 13.4 |
| | Pretty good chance | 1.7 | 3.2 | 4.9 | 6.4 | 7.2 | 7.2 | 6.5 |
| | Very good chance | 4.0 | 4.7 | 5.1 | 6.4 | 6.3 | 6.9 | 6.5 |
| Cool if carry a gun | No Answer | 8.5 | 8.4 | 7.6 | 11.2 | 10.4 | 10.2 | 10.3 |
| | No or very little chance | 73.7 | 71.0 | 69.4 | 65.6 | 66.5 | 67.0 | 64.2 |
| | Little chance | 6.6 | 7.6 | 8.9 | 9.5 | 9.8 | 9.5 | 10.9 |
| | Some chance | 4.1 | 4.7 | 5.5 | 6.1 | 6.2 | 5.7 | 7.0 |
| | Pretty good chance | 2.1 | 3.0 | 3.4 | 2.7 | 2.5 | 2.7 | 2.7 |
| | Very good chance | 5.1 | 5.3 | 5.2 | 4.9 | 4.6 | 5.0 | 4.8 |

School Commitment
by Indiana Male 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|----------------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Days of school skipped during last month | No Answer | 6.4 | 6.4 | 6.1 | 9.0 | 8.6 | 8.2 | 8.5 |
| | None | 86.3 | 84.7 | 82.6 | 79.0 | 78.1 | 75.7 | 69.9 |
| | 1 | 3.4 | 3.9 | 5.3 | 5.2 | 5.8 | 6.9 | 9.0 |
| | 2 | 1.3 | 1.8 | 2.3 | 2.2 | 2.9 | 3.4 | 4.7 |
| | 3 | 1.2 | 1.1 | 1.4 | 1.8 | 1.6 | 2.2 | 3.3 |
| | 4-5 | .7 | .8 | 1.2 | 1.4 | 1.4 | 1.6 | 2.5 |
| | 6-10 | .3 | .4 | .4 | .5 | .6 | .8 | .9 |
| | 11 or more | .5 | .8 | .7 | .9 | 1.0 | 1.2 | 1.3 |
| Feel schoolwork is meaningful | No Answer | 6.7 | 7.0 | 6.1 | 9.3 | 8.8 | 8.5 | 8.6 |
| | Almost always | 38.7 | 30.1 | 25.3 | 20.3 | 16.8 | 16.5 | 14.6 |
| | Often | 20.9 | 22.4 | 23.5 | 21.4 | 20.1 | 19.2 | 17.6 |
| | Sometimes | 19.8 | 22.5 | 24.4 | 25.5 | 27.9 | 27.1 | 28.2 |
| | Seldom | 6.3 | 9.4 | 12.0 | 14.6 | 16.7 | 19.1 | 20.4 |
| | Never | 7.5 | 8.6 | 8.7 | 8.8 | 9.7 | 9.5 | 10.6 |
| How interesting are courses | No Answer | 10.0 | 8.6 | 7.1 | 10.2 | 9.5 | 9.0 | 9.0 |
| | Very interesting and stimulating | 19.1 | 13.5 | 11.2 | 9.2 | 8.3 | 8.9 | 8.7 |
| | Quite interesting | 24.7 | 22.3 | 20.2 | 19.6 | 17.9 | 18.4 | 17.0 |
| | Fairly interesting | 26.9 | 29.9 | 31.9 | 32.6 | 33.1 | 33.2 | 32.2 |
| | Slightly dull | 10.5 | 13.6 | 16.3 | 15.8 | 16.9 | 16.7 | 17.7 |
| | Very dull | 8.7 | 12.1 | 13.4 | 12.7 | 14.3 | 13.8 | 15.3 |
| Importance of school to later life | No Answer | 8.0 | 8.0 | 7.0 | 10.3 | 9.7 | 9.3 | 9.2 |
| | Very important | 59.1 | 47.9 | 41.8 | 31.1 | 25.9 | 24.0 | 20.8 |
| | Quite important | 15.9 | 20.0 | 20.5 | 21.1 | 20.4 | 18.8 | 18.5 |
| | Fairly important | 9.2 | 12.5 | 15.6 | 18.3 | 19.4 | 20.6 | 22.3 |
| | Slightly important | 5.1 | 7.7 | 10.7 | 13.3 | 17.0 | 19.9 | 20.6 |
| | Not at all important | 2.7 | 3.8 | 4.5 | 5.8 | 7.6 | 7.5 | 8.6 |
| Grades better than other students | No Answer | 8.0 | 8.2 | 7.1 | 10.5 | 9.8 | 9.5 | 9.3 |
| | YES! | 22.3 | 22.1 | 21.3 | 20.6 | 20.0 | 20.8 | 21.6 |
| | yes | 44.4 | 43.2 | 44.6 | 41.9 | 42.0 | 42.5 | 43.1 |
| | no | 19.7 | 20.6 | 21.5 | 21.4 | 22.6 | 22.1 | 21.4 |
| | NO! | 5.6 | 5.9 | 5.5 | 5.7 | 5.5 | 5.2 | 4.6 |
| How often did you enjoy school | No Answer | 7.6 | 8.0 | 6.7 | 10.3 | 9.8 | 9.5 | 9.2 |
| | Never | 10.8 | 12.1 | 12.0 | 12.6 | 12.9 | 12.5 | 12.2 |
| | Seldom | 10.3 | 12.6 | 13.8 | 15.2 | 16.6 | 17.9 | 17.5 |
| | Sometimes | 25.4 | 26.0 | 28.4 | 28.9 | 29.7 | 29.9 | 30.1 |
| | Often | 24.3 | 24.6 | 25.0 | 22.4 | 21.7 | 21.5 | 21.7 |
| | A lot | 21.6 | 16.7 | 14.1 | 10.6 | 9.3 | 8.6 | 9.2 |

School Commitment
by Indiana Male 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| How often did you hate school | No Answer | 7.7 | 7.9 | 6.9 | 10.4 | 9.9 | 9.6 | 9.2 |
| | Never | 19.6 | 14.0 | 11.4 | 9.2 | 7.9 | 7.7 | 7.7 |
| | Seldom | 23.7 | 24.8 | 24.1 | 21.5 | 20.5 | 19.4 | 19.8 |
| | Sometimes | 23.0 | 24.3 | 27.1 | 27.2 | 28.0 | 28.9 | 29.2 |
| | Often | 11.4 | 12.9 | 14.3 | 16.3 | 17.1 | 18.1 | 19.2 |
| | A lot | 14.5 | 16.0 | 16.1 | 15.4 | 16.5 | 16.3 | 14.9 |
| How often did you do your best work | No Answer | 7.7 | 8.1 | 6.9 | 10.5 | 10.0 | 9.6 | 9.3 |
| | Never | 2.1 | 3.0 | 3.0 | 3.4 | 3.9 | 3.4 | 3.8 |
| | Seldom | 2.7 | 3.9 | 5.1 | 5.5 | 6.3 | 6.8 | 8.4 |
| | Sometimes | 9.0 | 11.9 | 16.0 | 18.6 | 20.7 | 23.0 | 23.9 |
| | Often | 18.7 | 23.5 | 27.4 | 28.5 | 29.6 | 29.6 | 30.6 |
| | A lot | 59.8 | 49.6 | 41.6 | 33.6 | 29.5 | 27.6 | 24.0 |

**Opportunities for Prosocial Involvement
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Students have chances to help decide activities and rules | No Answer | 10.6 | 10.7 | 8.9 | 13.7 | 12.8 | 12.3 | 11.9 |
| | YES! | 28.2 | 23.2 | 20.7 | 20.2 | 18.7 | 17.5 | 16.3 |
| | yes | 31.1 | 30.9 | 31.8 | 31.3 | 30.9 | 31.8 | 31.5 |
| | no | 18.9 | 22.6 | 25.4 | 23.6 | 25.5 | 26.0 | 26.6 |
| | NO! | 11.2 | 12.6 | 13.2 | 11.2 | 12.1 | 12.5 | 13.6 |
| Teachers ask to work on special class projects | No Answer | 11.2 | 11.0 | 9.2 | 13.9 | 13.0 | 12.5 | 12.1 |
| | YES! | 22.6 | 19.0 | 15.6 | 12.8 | 11.5 | 10.9 | 10.4 |
| | yes | 30.6 | 29.1 | 30.6 | 27.8 | 28.1 | 29.2 | 30.0 |
| | no | 24.3 | 28.3 | 31.6 | 33.9 | 35.3 | 35.2 | 35.5 |
| | NO! | 11.4 | 12.6 | 13.0 | 11.5 | 12.0 | 12.1 | 12.0 |
| Chances for kids to get involved in activities outside of class | No Answer | 11.1 | 11.3 | 9.5 | 14.4 | 13.3 | 13.1 | 12.5 |
| | YES! | 63.6 | 62.2 | 57.3 | 50.7 | 47.9 | 45.1 | 43.5 |
| | yes | 19.3 | 20.1 | 26.1 | 26.6 | 30.1 | 32.5 | 34.0 |
| | no | 3.8 | 3.9 | 4.5 | 5.5 | 5.8 | 6.2 | 7.0 |
| | NO! | 2.2 | 2.5 | 2.6 | 2.7 | 2.9 | 3.1 | 2.9 |
| Chances for kids to talk with teacher one-on-one | No Answer | 11.8 | 11.8 | 10.1 | 14.9 | 13.7 | 13.5 | 12.7 |
| | YES! | 48.1 | 42.6 | 38.3 | 32.0 | 29.7 | 29.0 | 28.7 |
| | yes | 28.0 | 30.5 | 34.0 | 35.4 | 37.6 | 39.6 | 41.0 |
| | no | 8.8 | 10.4 | 12.6 | 12.9 | 14.3 | 13.5 | 13.1 |
| | NO! | 3.4 | 4.7 | 5.0 | 4.8 | 4.7 | 4.4 | 4.5 |
| Chances to be a part of class discussion and activities | No Answer | 12.8 | 12.6 | 10.6 | 15.4 | 13.9 | 13.7 | 13.2 |
| | YES! | 42.9 | 37.1 | 33.1 | 26.5 | 24.4 | 23.8 | 23.1 |
| | yes | 30.8 | 34.1 | 39.3 | 40.4 | 43.3 | 45.4 | 47.7 |
| | no | 9.4 | 11.0 | 12.0 | 13.0 | 13.7 | 12.5 | 12.1 |
| | NO! | 4.1 | 5.1 | 5.0 | 4.8 | 4.8 | 4.6 | 4.0 |

**Rewards for Prosocial Involvement
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Teachers notice when doing a good job and let me know | No Answer | 11.6 | 11.6 | 9.9 | 14.5 | 13.6 | 13.1 | 12.5 |
| | YES! | 40.0 | 32.1 | 27.1 | 20.9 | 18.4 | 17.6 | 15.5 |
| | yes | 31.5 | 34.4 | 38.4 | 37.3 | 38.7 | 40.4 | 41.8 |
| | no | 11.4 | 14.6 | 16.9 | 19.4 | 21.4 | 21.1 | 22.0 |
| | NO! | 5.4 | 7.3 | 7.6 | 7.9 | 7.8 | 7.8 | 8.2 |
| Feel safe at school | No Answer | 11.3 | 11.4 | 9.8 | 14.7 | 13.5 | 13.2 | 12.6 |
| | YES! | 52.2 | 44.6 | 39.3 | 29.8 | 30.4 | 31.4 | 32.8 |
| | yes | 25.5 | 30.3 | 35.5 | 38.6 | 39.7 | 40.9 | 40.9 |
| | no | 6.5 | 8.3 | 9.6 | 11.2 | 10.4 | 9.5 | 9.0 |
| | NO! | 4.5 | 5.4 | 5.9 | 5.7 | 5.9 | 4.9 | 4.6 |
| School lets parents know when do good | No Answer | 12.1 | 12.2 | 10.4 | 15.0 | 13.8 | 13.5 | 12.9 |
| | YES! | 29.3 | 25.3 | 21.0 | 16.2 | 14.5 | 13.4 | 12.0 |
| | yes | 25.0 | 24.2 | 25.5 | 23.1 | 22.7 | 23.1 | 23.4 |
| | no | 21.8 | 23.8 | 26.8 | 29.1 | 31.1 | 32.3 | 32.7 |
| | NO! | 11.8 | 14.5 | 16.3 | 16.6 | 17.9 | 17.7 | 19.0 |
| Teachers praise when do well in school | No Answer | 13.2 | 12.8 | 10.7 | 15.5 | 14.0 | 13.9 | 13.2 |
| | YES! | 27.4 | 24.2 | 22.0 | 17.4 | 15.7 | 14.7 | 13.4 |
| | yes | 29.9 | 30.1 | 32.0 | 30.9 | 31.1 | 33.0 | 34.0 |
| | no | 19.4 | 21.2 | 24.1 | 25.0 | 27.2 | 27.2 | 28.3 |
| | NO! | 10.1 | 11.7 | 11.3 | 11.2 | 12.1 | 11.1 | 11.0 |

**Participation in Afterschool Activities
by Indiana 6 Male th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Participation in ROCK, LEAD, SADD, etc. | No Answer | 17.1 | 17.8 | 15.5 | 20.0 | 18.7 | 17.6 | 16.8 |
| | Yes | 16.4 | 10.9 | 7.4 | 5.8 | 5.3 | 5.3 | 5.0 |
| | No | 66.5 | 71.2 | 77.2 | 74.2 | 76.0 | 77.1 | 78.2 |

**Favorable Attitudes Toward Drug Use
by Indiana Male 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Wrong to take a gun to school | No Answer | 11.8 | 12.5 | 9.5 | 14.0 | 12.5 | 12.0 | 11.8 |
| | Very wrong | 81.2 | 77.5 | 78.6 | 72.4 | 74.3 | 74.3 | 74.4 |
| | Wrong | 4.6 | 6.1 | 7.5 | 8.3 | 8.0 | 8.1 | 8.2 |
| | A little bit wrong | 1.5 | 2.3 | 2.6 | 2.8 | 2.8 | 2.8 | 3.1 |
| | Not at all wrong | .9 | 1.5 | 1.8 | 2.5 | 2.4 | 2.8 | 2.5 |
| Wrong to steal something more than \$5 | No Answer | 12.0 | 12.7 | 9.6 | 14.4 | 12.8 | 12.3 | 12.1 |
| | Very wrong | 60.7 | 55.1 | 52.2 | 46.2 | 46.9 | 48.5 | 50.4 |
| | Wrong | 19.5 | 21.3 | 24.3 | 24.5 | 25.8 | 25.1 | 25.9 |
| | A little bit wrong | 6.2 | 8.0 | 10.4 | 11.3 | 11.1 | 10.5 | 8.9 |
| | Not at all wrong | 1.7 | 2.9 | 3.6 | 3.6 | 3.4 | 3.6 | 2.7 |
| Wrong to attack someone | No Answer | 12.2 | 12.9 | 9.7 | 14.7 | 13.0 | 12.6 | 12.2 |
| | Very wrong | 72.3 | 66.6 | 63.6 | 56.3 | 56.0 | 55.9 | 57.6 |
| | Wrong | 9.8 | 13.0 | 16.1 | 17.8 | 19.7 | 19.9 | 19.8 |
| | A little bit wrong | 3.8 | 4.8 | 7.2 | 7.6 | 7.8 | 8.0 | 7.4 |
| | Not at all wrong | 1.8 | 2.8 | 3.4 | 3.7 | 3.5 | 3.6 | 2.9 |
| Wrong to pick a fight | No Answer | 12.3 | 13.1 | 9.8 | 14.8 | 13.0 | 12.6 | 12.3 |
| | Very wrong | 55.8 | 48.9 | 44.1 | 39.1 | 38.1 | 39.3 | 40.1 |
| | Wrong | 19.8 | 21.9 | 24.2 | 24.4 | 26.9 | 26.3 | 28.4 |
| | A little bit wrong | 8.5 | 11.3 | 15.3 | 15.6 | 15.8 | 16.1 | 14.7 |
| | Not at all wrong | 3.6 | 4.8 | 6.5 | 6.0 | 6.2 | 5.7 | 4.6 |
| Wrong to skip school | No Answer | 12.4 | 13.0 | 10.0 | 14.9 | 13.3 | 12.8 | 12.4 |
| | Very wrong | 72.6 | 64.4 | 59.5 | 48.5 | 45.2 | 42.3 | 39.8 |
| | Wrong | 10.4 | 15.1 | 19.8 | 22.5 | 25.1 | 26.4 | 26.6 |
| | A little bit wrong | 2.9 | 5.0 | 7.3 | 10.0 | 12.0 | 13.4 | 16.2 |
| | Not at all wrong | 1.7 | 2.5 | 3.5 | 4.1 | 4.4 | 5.0 | 5.0 |
| Wrong to drink alcohol regularly | No Answer | 12.4 | 13.2 | 10.1 | 15.0 | 13.3 | 12.8 | 12.4 |
| | Very wrong | 75.9 | 68.4 | 61.2 | 47.4 | 42.2 | 38.6 | 34.0 |
| | Wrong | 6.9 | 9.8 | 14.9 | 17.4 | 18.7 | 18.8 | 18.7 |
| | A little bit wrong | 3.0 | 5.3 | 9.1 | 13.3 | 16.7 | 18.6 | 21.7 |
| | Not at all wrong | 1.7 | 3.3 | 4.8 | 6.9 | 9.1 | 11.2 | 13.2 |
| Wrong to smoke cigarettes | No Answer | 12.6 | 13.2 | 10.1 | 15.2 | 13.4 | 12.8 | 12.6 |
| | Very wrong | 77.2 | 71.4 | 65.7 | 53.6 | 50.1 | 44.5 | 36.5 |
| | Wrong | 6.5 | 8.6 | 13.5 | 16.1 | 17.5 | 18.0 | 17.7 |
| | A little bit wrong | 2.1 | 3.7 | 6.7 | 8.9 | 11.0 | 13.9 | 16.0 |
| | Not at all wrong | 1.6 | 3.1 | 4.1 | 6.1 | 8.0 | 10.8 | 17.2 |

Favorable Attitudes Toward Drug Use
by Indiana Male 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|----------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Wrong to smoke marijuana | No Answer | 12.7 | 13.3 | 10.3 | 15.2 | 13.2 | 12.9 | 12.5 |
| | Very wrong | 80.3 | 73.8 | 67.6 | 53.2 | 47.8 | 43.2 | 39.0 |
| | Wrong | 3.5 | 5.8 | 9.8 | 12.9 | 14.7 | 15.3 | 16.3 |
| | A little bit wrong | 1.6 | 3.4 | 6.4 | 9.1 | 11.7 | 13.5 | 15.0 |
| | Not at all wrong | 1.9 | 3.6 | 5.9 | 9.7 | 12.5 | 15.2 | 17.3 |
| Wrong to use illegal drugs | No Answer | 12.9 | 13.7 | 10.3 | 15.3 | 13.7 | 13.0 | 12.8 |
| | Very wrong | 82.6 | 79.0 | 78.2 | 69.1 | 68.3 | 67.3 | 65.4 |
| | Wrong | 2.5 | 3.9 | 6.7 | 8.8 | 10.4 | 10.4 | 11.7 |
| | A little bit wrong | .8 | 1.5 | 2.5 | 3.5 | 4.2 | 5.0 | 5.7 |
| | Not at all wrong | 1.2 | 1.9 | 2.3 | 3.3 | 3.5 | 4.3 | 4.5 |

Rebelliousness
by Indiana Male 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|----------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Like to see what can get away with | No Answer | 14.7 | 14.6 | 11.1 | 16.2 | 14.2 | 13.5 | 13.3 |
| | Very false | 56.9 | 49.3 | 44.6 | 39.2 | 37.6 | 35.8 | 34.8 |
| | Somewhat false | 15.9 | 18.9 | 22.0 | 21.0 | 23.7 | 23.5 | 23.4 |
| | Somewhat true | 9.4 | 12.8 | 16.5 | 17.6 | 18.5 | 20.8 | 21.7 |
| | Very true | 3.2 | 4.4 | 5.8 | 5.9 | 5.9 | 6.4 | 6.8 |
| Ignore rules | No Answer | 14.7 | 14.8 | 11.3 | 16.4 | 14.4 | 13.7 | 13.5 |
| | Very false | 59.1 | 53.6 | 49.5 | 42.7 | 40.8 | 39.7 | 36.4 |
| | Somewhat false | 15.8 | 17.9 | 21.5 | 22.5 | 24.5 | 24.9 | 26.2 |
| | Somewhat true | 7.3 | 9.7 | 12.6 | 13.4 | 15.3 | 16.2 | 17.6 |
| | Very true | 3.0 | 4.0 | 5.1 | 5.0 | 5.1 | 5.6 | 6.4 |
| Do opposite of what I am told | No Answer | 14.9 | 15.1 | 11.6 | 16.6 | 14.6 | 13.8 | 13.6 |
| | Very false | 61.3 | 56.5 | 54.2 | 50.1 | 51.2 | 52.3 | 51.6 |
| | Somewhat false | 14.4 | 16.7 | 20.0 | 19.6 | 21.0 | 20.9 | 21.4 |
| | Somewhat true | 6.5 | 8.3 | 10.1 | 9.7 | 9.4 | 9.3 | 9.7 |
| | Very true | 2.9 | 3.5 | 4.1 | 4.1 | 3.8 | 3.6 | 3.7 |

**Laws and Norms Favorable to Drug Use
by Indiana Male 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Caught by police alcohol | No Answer | 15.9 | 15.6 | 12.5 | 17.5 | 15.7 | 14.8 | 14.7 |
| | YES! | 26.3 | 21.1 | 15.8 | 10.7 | 9.2 | 8.5 | 7.7 |
| | yes | 25.5 | 23.9 | 21.6 | 17.6 | 15.0 | 13.9 | 13.2 |
| | no | 21.6 | 25.9 | 34.2 | 36.8 | 41.3 | 42.7 | 43.4 |
| | NO! | 10.7 | 13.4 | 16.0 | 17.4 | 18.9 | 20.1 | 21.0 |
| Caught by police marijuana | No Answer | 15.9 | 15.7 | 12.5 | 17.6 | 15.8 | 15.1 | 15.0 |
| | YES! | 36.9 | 28.4 | 20.7 | 12.9 | 10.6 | 9.0 | 7.8 |
| | yes | 23.8 | 24.1 | 23.9 | 19.1 | 16.1 | 15.5 | 14.4 |
| | no | 14.7 | 20.1 | 28.3 | 33.5 | 38.2 | 40.4 | 41.7 |
| | NO! | 8.7 | 11.7 | 14.6 | 16.9 | 19.3 | 20.1 | 21.2 |
| Caught by police gun | No Answer | 16.0 | 15.8 | 12.6 | 17.7 | 15.8 | 15.2 | 14.9 |
| | YES! | 40.8 | 33.3 | 27.5 | 19.8 | 17.0 | 15.9 | 13.9 |
| | yes | 21.6 | 23.6 | 26.0 | 24.3 | 24.2 | 24.3 | 24.1 |
| | no | 13.4 | 17.0 | 22.0 | 25.5 | 29.0 | 30.9 | 32.3 |
| | NO! | 8.1 | 10.3 | 11.9 | 12.7 | 13.9 | 13.7 | 14.9 |
| Caught by police smoke cigarettes | No Answer | 16.7 | 16.1 | 12.9 | 18.2 | 16.2 | 15.3 | 15.4 |
| | YES! | 24.3 | 19.6 | 14.7 | 10.0 | 8.4 | 7.6 | 6.3 |
| | yes | 21.9 | 20.4 | 18.1 | 14.4 | 11.7 | 10.6 | 9.4 |
| | no | 22.8 | 26.5 | 33.2 | 34.4 | 37.5 | 38.2 | 38.5 |
| | NO! | 14.3 | 17.4 | 21.2 | 23.0 | 26.3 | 28.2 | 30.4 |
| Neighborhood feel about marijuana | No Answer | 14.8 | 15.4 | 11.7 | 17.4 | 15.5 | 14.7 | 14.5 |
| | Very wrong | 73.5 | 67.6 | 62.7 | 49.4 | 45.6 | 42.6 | 40.8 |
| | Wrong | 7.8 | 11.0 | 15.9 | 19.7 | 22.7 | 24.6 | 26.0 |
| | A little bit wrong | 2.4 | 3.8 | 6.6 | 9.0 | 11.2 | 12.5 | 12.6 |
| | Not at all wrong | 1.4 | 2.3 | 3.2 | 4.5 | 4.9 | 5.5 | 6.1 |
| Neighborhood feel about alcohol | No Answer | 15.0 | 15.4 | 12.0 | 17.6 | 15.6 | 14.9 | 14.7 |
| | Very wrong | 65.1 | 58.7 | 52.9 | 40.2 | 35.2 | 31.3 | 28.1 |
| | Wrong | 12.3 | 14.8 | 19.8 | 22.4 | 25.1 | 25.4 | 25.3 |
| | A little bit wrong | 5.2 | 7.6 | 10.6 | 13.6 | 16.8 | 19.6 | 22.0 |
| | Not at all wrong | 2.4 | 3.5 | 4.8 | 6.2 | 7.3 | 8.7 | 10.0 |
| Neighborhood feel about cigarettes | No Answer | 15.3 | 15.7 | 12.1 | 17.8 | 15.8 | 15.1 | 14.8 |
| | Very wrong | 66.7 | 61.2 | 56.0 | 43.9 | 38.8 | 33.7 | 25.9 |
| | Wrong | 11.1 | 13.8 | 18.6 | 21.3 | 24.5 | 24.6 | 24.1 |
| | A little bit wrong | 4.6 | 5.9 | 8.7 | 10.7 | 14.0 | 17.7 | 20.5 |
| | Not at all wrong | 2.4 | 3.3 | 4.6 | 6.3 | 7.0 | 8.9 | 14.7 |

Community Rewards for Prosocial Involvement
by Indiana Male 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Neighbors notice good job and let me know | No Answer | 16.2 | 15.8 | 12.9 | 18.2 | 16.3 | 15.4 | 15.2 |
| | YES! | 20.1 | 17.4 | 14.5 | 12.2 | 10.6 | 9.9 | 9.3 |
| | yes | 20.7 | 19.9 | 20.0 | 18.3 | 17.1 | 17.1 | 17.0 |
| | no | 22.8 | 25.0 | 27.4 | 27.5 | 28.9 | 30.7 | 29.7 |
| | NO! | 20.2 | 21.9 | 25.2 | 23.8 | 27.1 | 26.8 | 28.8 |
| People in neighborhood who are proud of me | No Answer | 16.2 | 15.9 | 13.1 | 18.3 | 16.4 | 15.5 | 15.3 |
| | YES! | 23.1 | 19.8 | 16.4 | 13.5 | 12.2 | 11.1 | 10.9 |
| | yes | 23.6 | 24.0 | 24.3 | 22.8 | 23.0 | 23.5 | 24.1 |
| | no | 19.3 | 20.9 | 23.5 | 23.8 | 24.3 | 26.0 | 24.3 |
| | NO! | 17.8 | 19.5 | 22.7 | 21.7 | 24.1 | 23.9 | 25.4 |
| People in neighborhood who encourage to do my best | No Answer | 16.4 | 16.1 | 13.3 | 18.6 | 16.7 | 15.7 | 15.6 |
| | YES! | 28.9 | 24.1 | 19.4 | 15.7 | 13.6 | 13.0 | 12.1 |
| | yes | 23.2 | 23.1 | 25.0 | 23.2 | 23.3 | 23.7 | 24.1 |
| | no | 15.2 | 18.4 | 20.4 | 21.4 | 22.9 | 23.9 | 23.5 |
| | NO! | 16.3 | 18.3 | 21.9 | 21.2 | 23.4 | 23.7 | 24.8 |

**Personal and Family Information
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------|----------------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Hispanic | No Answer | 6.7 | 5.6 | 5.8 | 6.0 | 6.5 | 5.5 | 4.8 |
| | Hispanic | 7.6 | 8.5 | 8.1 | 8.1 | 7.2 | 6.9 | 6.3 |
| | Non-Hispanic | 85.7 | 85.8 | 86.1 | 85.9 | 86.4 | 87.6 | 88.9 |
| Race | No Answer | 3.0 | 3.5 | 3.4 | 4.0 | 3.3 | 2.7 | 3.0 |
| | White | 79.2 | 80.9 | 79.5 | 80.3 | 79.4 | 81.9 | 84.0 |
| | Black/African American | 6.7 | 4.8 | 6.0 | 5.3 | 7.0 | 5.9 | 5.8 |
| | Asian | 1.6 | 1.5 | 1.6 | 1.7 | 1.8 | 1.9 | 1.5 |
| | Native Hawaiian/Pacific Islander | .1 | .2 | .3 | .4 | .4 | .3 | .3 |
| | American Indian/Alaskan Native | .4 | .4 | .5 | .5 | .6 | .5 | .4 |
| | More than one race | 5.9 | 5.9 | 6.4 | 5.7 | 5.6 | 5.0 | 3.8 |
| | Race not known or other | 3.0 | 2.9 | 2.2 | 2.2 | 1.9 | 1.7 | 1.2 |
| Age | No Answer | .8 | 1.0 | .9 | 1.0 | 1.1 | 1.2 | .8 |
| | 10 years or younger | .3 | .0 | .0 | .0 | .0 | .0 | .0 |
| | 11 years | 25.6 | .1 | .0 | .0 | .0 | .0 | .0 |
| | 12 years | 65.4 | 25.2 | .1 | .0 | .0 | .0 | .0 |
| | 13 years | 7.6 | 66.7 | 24.4 | .1 | .0 | .0 | .0 |
| | 14 years | .3 | 6.7 | 66.8 | 23.8 | .1 | .0 | .0 |
| | 15 years | .0 | .3 | 7.4 | 67.5 | 21.3 | .3 | .0 |
| | 16 years | .0 | .0 | .3 | 7.2 | 70.6 | 23.0 | .4 |
| | 17 years | .0 | .0 | .0 | .3 | 6.7 | 70.1 | 24.3 |
| | 18 years or older | .0 | .0 | .0 | .0 | .3 | 5.5 | 74.5 |
| Grades | No Answer | 4.7 | 4.1 | 3.8 | 3.3 | 2.6 | 2.0 | 1.5 |
| | Mostly A's | 42.7 | 42.3 | 40.5 | 38.6 | 36.4 | 36.0 | 42.3 |
| | Mostly B's | 35.0 | 34.4 | 34.6 | 36.6 | 39.2 | 41.5 | 39.0 |
| | Mostly C's | 13.8 | 14.8 | 17.1 | 17.5 | 18.2 | 17.9 | 15.6 |
| | Mostly D's | 2.6 | 3.1 | 2.9 | 3.0 | 2.7 | 2.1 | 1.3 |
| | Mostly F's | 1.2 | 1.4 | 1.1 | 1.1 | .8 | .5 | .3 |
| Live with | No Answer | 1.7 | 1.3 | 1.4 | 1.3 | .9 | .9 | .7 |
| | Both parents | 58.5 | 58.7 | 54.7 | 55.1 | 54.0 | 55.0 | 55.2 |
| | Mother only | 14.6 | 14.6 | 16.9 | 16.9 | 18.0 | 17.0 | 17.2 |
| | Father only | 2.4 | 2.8 | 2.9 | 3.0 | 3.6 | 4.1 | 3.8 |
| | Mother & stepfather | 13.8 | 14.1 | 15.5 | 15.0 | 14.9 | 13.8 | 12.2 |
| | Father & stepmother | 2.9 | 3.0 | 3.3 | 3.6 | 3.6 | 3.5 | 2.7 |
| | Other | 6.0 | 5.5 | 5.2 | 5.0 | 5.1 | 5.9 | 8.2 |
| Employed - You | No Answer | 17.8 | 14.1 | 13.7 | 12.0 | 9.4 | 6.3 | 4.5 |
| | Yes, full time | .7 | .5 | .8 | .9 | 1.2 | 1.6 | 2.5 |
| | Yes, part time | 4.8 | 6.0 | 7.9 | 10.3 | 18.2 | 36.9 | 49.6 |
| | No | 76.8 | 79.4 | 77.5 | 76.8 | 71.2 | 55.2 | 43.4 |
| Employed - Mom | No Answer | 6.4 | 5.0 | 4.7 | 4.6 | 4.1 | 3.6 | 3.5 |
| | Yes, full time | 51.1 | 54.4 | 57.0 | 58.7 | 61.4 | 63.7 | 64.0 |
| | Yes, part time | 22.4 | 21.0 | 18.0 | 16.8 | 15.2 | 14.4 | 13.4 |
| | No | 20.1 | 19.6 | 20.3 | 19.9 | 19.4 | 18.4 | 19.1 |
| Employed - Dad | No Answer | 8.4 | 7.1 | 7.6 | 7.8 | 7.9 | 7.4 | 7.1 |
| | Yes, full time | 69.6 | 72.8 | 74.1 | 75.6 | 76.9 | 78.2 | 77.5 |
| | Yes, part time | 13.3 | 10.4 | 8.8 | 7.4 | 5.5 | 4.7 | 4.0 |
| | No | 8.8 | 9.7 | 9.5 | 9.3 | 9.8 | 9.7 | 11.4 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|-------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | | .2 | .2 | .3 | .3 | .4 | .4 | .4 |
| | Never | | 93.4 | 87.1 | 78.8 | 72.6 | 68.2 | 63.3 | 60.4 |
| | 1-5 times | | 4.8 | 8.5 | 12.2 | 13.2 | 13.9 | 14.5 | 15.3 |
| | 6-19 times | | .8 | 1.9 | 3.1 | 4.1 | 4.9 | 5.2 | 5.5 |
| | 20-40 times | | .2 | .9 | 1.9 | 2.5 | 2.9 | 2.7 | 3.3 |
| | 40+ times | | .6 | 1.5 | 3.7 | 7.4 | 9.7 | 13.8 | 15.1 |
| Smokeless Tobacco | No Answer | | .1 | .1 | .2 | .3 | .3 | .3 | .4 |
| | Never | | 98.9 | 97.4 | 95.4 | 93.1 | 91.4 | 89.8 | 88.9 |
| | 1-5 times | | .7 | 2.1 | 3.4 | 5.1 | 6.2 | 7.0 | 8.0 |
| | 6-19 times | | .1 | .2 | .4 | .8 | 1.1 | 1.3 | 1.2 |
| | 20-40 times | | .0 | .1 | .3 | .4 | .5 | .6 | .7 |
| | 40+ times | | .1 | .1 | .3 | .4 | .5 | 1.0 | .9 |
| Cigars | No Answer | | .2 | .2 | .3 | .4 | .5 | .6 | .5 |
| | Never | | 98.9 | 97.3 | 94.4 | 90.2 | 85.7 | 81.4 | 75.6 |
| | 1-5 times | | .6 | 2.0 | 3.8 | 6.4 | 9.2 | 11.1 | 14.0 |
| | 6-19 times | | .1 | .3 | .8 | 1.7 | 2.5 | 3.6 | 5.6 |
| | 20-40 times | | .0 | .1 | .3 | .8 | 1.0 | 1.5 | 2.1 |
| | 40+ times | | .1 | .1 | .3 | .5 | 1.1 | 1.7 | 2.2 |
| Pipe | No Answer | | .3 | .4 | .6 | .7 | .8 | .8 | 1.1 |
| | Never | | 98.8 | 97.3 | 94.7 | 91.0 | 87.2 | 82.9 | 77.2 |
| | 1-5 times | | .6 | 1.6 | 3.0 | 4.9 | 6.7 | 8.5 | 10.8 |
| | 6-19 times | | .1 | .3 | .8 | 1.5 | 2.3 | 3.6 | 5.4 |
| | 20-40 times | | .1 | .2 | .5 | .9 | 1.3 | 1.8 | 2.5 |
| | 40+ times | | .1 | .2 | .5 | 1.0 | 1.6 | 2.4 | 3.1 |
| Alcohol | No Answer | | .5 | .6 | .6 | .9 | .8 | .8 | 1.1 |
| | Never | | 84.6 | 74.6 | 60.6 | 50.0 | 42.9 | 38.7 | 33.8 |
| | 1-5 times | | 12.6 | 17.8 | 24.0 | 25.6 | 26.0 | 24.0 | 21.8 |
| | 6-19 times | | 1.3 | 4.6 | 8.3 | 12.7 | 14.6 | 16.3 | 16.3 |
| | 20-40 times | | .4 | 1.3 | 3.4 | 5.3 | 7.4 | 8.7 | 11.1 |
| | 40+ times | | .5 | 1.0 | 3.1 | 5.6 | 8.4 | 11.5 | 15.8 |
| Marijuana | No Answer | | .3 | .3 | .4 | .7 | .7 | .8 | .9 |
| | Never | | 97.4 | 93.4 | 85.7 | 78.2 | 71.3 | 66.8 | 64.0 |
| | 1-5 times | | 1.5 | 3.7 | 7.2 | 9.3 | 11.4 | 12.1 | 13.6 |
| | 6-19 times | | .4 | 1.1 | 2.7 | 3.8 | 5.4 | 6.2 | 6.7 |
| | 20-40 times | | .2 | .7 | 1.4 | 2.5 | 3.5 | 4.1 | 4.5 |
| | 40+ times | | .3 | .8 | 2.6 | 5.5 | 7.8 | 9.9 | 10.3 |
| Cocaine | No Answer | | .2 | .2 | .3 | .4 | .5 | .5 | .7 |
| | Never | | 99.3 | 99.1 | 98.0 | 97.4 | 96.9 | 96.2 | 95.5 |
| | 1-5 times | | .2 | .5 | 1.1 | 1.4 | 1.8 | 2.3 | 2.3 |
| | 6-19 times | | .1 | .1 | .3 | .3 | .5 | .5 | .6 |
| | 20-40 times | | .0 | .1 | .2 | .2 | .1 | .2 | .4 |
| | 40+ times | | .1 | .1 | .2 | .3 | .2 | .3 | .5 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Crack | No Answer | | .2 | .2 | .3 | .4 | .4 | .4 | .4 |
| | Never | | 99.3 | 99.0 | 98.0 | 98.1 | 98.3 | 98.1 | 98.1 |
| | 1-5 times | | .3 | .6 | 1.1 | 1.0 | .8 | .9 | .9 |
| | 6-19 times | | .1 | .1 | .1 | .2 | .2 | .3 | .2 |
| | 20-40 times | | .0 | .1 | .2 | .2 | .1 | .1 | .1 |
| | 40+ times | | .1 | .0 | .2 | .2 | .2 | .3 | .3 |
| Inhalants | No Answer | | 1.1 | .5 | .4 | .5 | .5 | .5 | .5 |
| | Never | | 97.5 | 96.3 | 93.9 | 92.7 | 92.6 | 92.6 | 93.5 |
| | 1-5 times | | .9 | 2.2 | 3.9 | 4.5 | 4.6 | 4.3 | 3.7 |
| | 6-19 times | | .3 | .6 | 1.0 | 1.4 | 1.4 | 1.4 | 1.4 |
| | 20-40 times | | .1 | .2 | .4 | .4 | .5 | .7 | .4 |
| | 40+ times | | .2 | .2 | .4 | .5 | .5 | .6 | .5 |
| Amphetamines | No Answer | | 1.4 | .6 | .5 | .5 | .6 | .4 | .6 |
| | Never | | 98.3 | 98.6 | 97.2 | 95.4 | 93.9 | 92.4 | 92.5 |
| | 1-5 times | | .1 | .4 | 1.5 | 2.0 | 2.6 | 3.2 | 2.8 |
| | 6-19 times | | .1 | .1 | .4 | 1.0 | 1.4 | 1.7 | 1.7 |
| | 20-40 times | | .0 | .1 | .2 | .6 | .8 | 1.1 | 1.0 |
| | 40+ times | | .1 | .1 | .2 | .6 | .8 | 1.2 | 1.3 |
| Methamphetamines | No Answer | | .9 | .4 | .4 | .5 | .5 | .4 | .5 |
| | Never | | 98.6 | 98.8 | 98.2 | 98.0 | 98.0 | 97.7 | 97.6 |
| | 1-5 times | | .2 | .5 | .9 | .9 | .9 | 1.1 | .9 |
| | 6-19 times | | .0 | .1 | .2 | .2 | .3 | .2 | .4 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .1 | .2 | .2 |
| | 40+ times | | .1 | .1 | .2 | .2 | .2 | .4 | .5 |
| Heroin | No Answer | | .8 | .4 | .4 | .5 | .5 | .4 | .6 |
| | Never | | 99.0 | 99.1 | 98.6 | 98.3 | 98.3 | 98.3 | 98.0 |
| | 1-5 times | | .1 | .3 | .6 | .6 | .7 | .7 | .6 |
| | 6-19 times | | .0 | .1 | .2 | .2 | .2 | .2 | .2 |
| | 20-40 times | | .0 | .1 | .1 | .2 | .1 | .1 | .2 |
| | 40+ times | | .1 | .1 | .1 | .1 | .2 | .2 | .3 |
| Steroids | No Answer | | .5 | .4 | .4 | .5 | .6 | .4 | .6 |
| | Never | | 99.2 | 99.0 | 98.8 | 98.7 | 98.6 | 98.7 | 98.5 |
| | 1-5 times | | .2 | .4 | .5 | .5 | .5 | .4 | .4 |
| | 6-19 times | | .0 | .1 | .2 | .1 | .2 | .1 | .2 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| | 40+ times | | .1 | .1 | .1 | .1 | .1 | .2 | .2 |
| Ecstasy | No Answer | | 1.5 | .7 | .4 | .5 | .6 | .4 | .6 |
| | Never | | 98.2 | 98.5 | 97.7 | 96.5 | 95.9 | 95.2 | 94.5 |
| | 1-5 times | | .1 | .6 | 1.3 | 2.1 | 2.5 | 3.0 | 3.5 |
| | 6-19 times | | .0 | .1 | .4 | .5 | .4 | .7 | .6 |
| | 20-40 times | | .0 | .0 | .1 | .2 | .3 | .3 | .4 |
| | 40+ times | | .1 | .0 | .2 | .2 | .3 | .3 | .4 |

**Lifetime Use of Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--------------------------|-------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Tranquilizers | No Answer | 1.5 | .6 | .5 | .7 | .6 | .6 | .7 |
| | Never | 98.2 | 98.6 | 97.5 | 96.1 | 95.4 | 94.4 | 94.5 |
| | 1-5 times | .2 | .6 | 1.3 | 1.8 | 2.0 | 2.5 | 2.3 |
| | 6-19 times | .0 | .1 | .3 | .8 | 1.0 | 1.2 | 1.3 |
| | 20-40 times | .0 | .1 | .1 | .3 | .4 | .6 | .5 |
| | 40+ times | .1 | .1 | .2 | .3 | .6 | .7 | .8 |
| Hallucinogens | No Answer | 1.6 | .6 | .6 | .8 | .7 | .7 | .8 |
| | Never | 98.0 | 98.8 | 97.3 | 96.3 | 95.5 | 94.6 | 94.6 |
| | 1-5 times | .2 | .4 | 1.4 | 1.8 | 2.6 | 3.2 | 3.0 |
| | 6-19 times | .1 | .1 | .2 | .5 | .6 | .9 | .7 |
| | 20-40 times | .0 | .0 | .2 | .3 | .2 | .2 | .3 |
| | 40+ times | .1 | .0 | .2 | .2 | .4 | .4 | .5 |
| Prescription painkillers | No Answer | .7 | .4 | .5 | .5 | .6 | .6 | .7 |
| | Never | 98.3 | 97.1 | 93.7 | 90.2 | 87.5 | 86.2 | 86.2 |
| | 1-5 times | .6 | 1.5 | 3.5 | 4.9 | 6.0 | 5.9 | 5.8 |
| | 6-19 times | .1 | .6 | 1.2 | 2.1 | 2.7 | 3.2 | 3.0 |
| | 20-40 times | .1 | .2 | .6 | 1.1 | 1.3 | 1.9 | 1.7 |
| | 40+ times | .2 | .2 | .6 | 1.2 | 1.8 | 2.2 | 2.6 |
| Prescription drugs | No Answer | .7 | .4 | .6 | .7 | .7 | .7 | .8 |
| | Never | 98.4 | 97.3 | 94.3 | 91.0 | 88.4 | 86.9 | 86.5 |
| | 1-5 times | .5 | 1.6 | 2.9 | 4.3 | 5.3 | 5.3 | 5.6 |
| | 6-19 times | .1 | .4 | 1.0 | 1.7 | 2.3 | 2.8 | 2.7 |
| | 20-40 times | .1 | .2 | .6 | .9 | 1.3 | 1.9 | 1.6 |
| | 40+ times | .2 | .2 | .6 | 1.3 | 1.9 | 2.4 | 2.8 |
| Over the counter drugs | No Answer | .6 | .6 | .5 | .7 | .7 | .6 | .7 |
| | Never | 98.3 | 96.5 | 93.8 | 91.5 | 90.6 | 90.7 | 91.5 |
| | 1-5 times | .6 | 2.2 | 3.7 | 4.6 | 4.8 | 4.4 | 4.1 |
| | 6-19 times | .1 | .3 | .9 | 1.6 | 1.7 | 2.0 | 1.8 |
| | 20-40 times | .0 | .3 | .5 | .7 | .9 | 1.0 | .7 |
| | 40+ times | .2 | .2 | .5 | 1.0 | 1.2 | 1.4 | 1.2 |

Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana **Female** 6th - 12th Graders, 2011

(Values expressed as percentages)

| | | | Grade | | | | | | |
|-------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | | 1.2 | 1.1 | 1.3 | 1.3 | 1.4 | 1.2 | 1.4 |
| | Never | | 96.2 | 92.7 | 88.1 | 84.2 | 82.5 | 79.2 | 77.2 |
| | 1-5 times | | 1.9 | 4.0 | 6.3 | 7.1 | 6.9 | 6.7 | 6.9 |
| | 6-19 times | | .3 | 1.1 | 1.7 | 2.5 | 2.5 | 3.1 | 2.9 |
| | 20-40 times | | .1 | .4 | .9 | 1.5 | 1.7 | 2.4 | 2.4 |
| | 40+ times | | .2 | .6 | 1.6 | 3.4 | 5.0 | 7.4 | 9.2 |
| Smokeless Tobacco | No Answer | | 1.4 | 1.2 | 1.7 | 1.4 | 1.5 | 1.5 | 1.5 |
| | Never | | 98.2 | 97.7 | 96.7 | 96.4 | 96.2 | 95.8 | 96.0 |
| | 1-5 times | | .3 | .9 | 1.2 | 1.6 | 1.6 | 1.6 | 1.6 |
| | 6-19 times | | .0 | .1 | .2 | .3 | .3 | .5 | .4 |
| | 20-40 times | | .0 | .0 | .1 | .2 | .2 | .3 | .2 |
| | 40+ times | | .1 | .1 | .1 | .1 | .2 | .4 | .3 |
| Cigars | No Answer | | 1.7 | 1.2 | 1.8 | 1.6 | 1.6 | 1.6 | 1.7 |
| | Never | | 97.9 | 97.7 | 96.1 | 94.8 | 93.5 | 92.4 | 89.8 |
| | 1-5 times | | .2 | .8 | 1.6 | 2.6 | 3.7 | 4.5 | 6.3 |
| | 6-19 times | | .1 | .1 | .3 | .5 | .7 | .9 | 1.2 |
| | 20-40 times | | .0 | .0 | .1 | .3 | .2 | .3 | .4 |
| | 40+ times | | .0 | .0 | .1 | .1 | .3 | .3 | .5 |
| Pipe | No Answer | | 1.7 | 1.3 | 1.9 | 1.7 | 1.8 | 1.9 | 2.0 |
| | Never | | 97.8 | 97.5 | 95.4 | 94.4 | 92.9 | 91.4 | 88.1 |
| | 1-5 times | | .4 | .9 | 1.9 | 2.7 | 3.4 | 4.6 | 7.4 |
| | 6-19 times | | .1 | .1 | .5 | .7 | .9 | 1.2 | 1.5 |
| | 20-40 times | | .0 | .1 | .1 | .3 | .5 | .4 | .4 |
| | 40+ times | | .0 | .1 | .1 | .3 | .4 | .5 | .6 |
| Alcohol | No Answer | | 1.9 | 1.4 | 2.0 | 1.8 | 1.7 | 1.7 | 1.8 |
| | Never | | 93.2 | 88.0 | 79.7 | 73.2 | 70.4 | 68.0 | 62.8 |
| | 1-5 times | | 4.2 | 8.8 | 14.4 | 19.3 | 21.0 | 22.3 | 25.9 |
| | 6-19 times | | .4 | 1.2 | 2.8 | 4.0 | 4.7 | 5.9 | 6.8 |
| | 20-40 times | | .1 | .3 | .7 | .9 | 1.2 | 1.2 | 1.6 |
| | 40+ times | | .2 | .2 | .5 | .7 | 1.0 | .9 | 1.1 |
| Marijuana | No Answer | | 1.8 | 1.4 | 1.9 | 1.7 | 1.8 | 1.8 | 1.9 |
| | Never | | 96.9 | 94.9 | 90.1 | 86.5 | 83.9 | 82.4 | 82.2 |
| | 1-5 times | | .9 | 2.5 | 5.0 | 6.7 | 7.9 | 8.8 | 8.6 |
| | 6-19 times | | .3 | .8 | 1.5 | 2.3 | 2.8 | 3.1 | 2.9 |
| | 20-40 times | | .1 | .3 | .8 | 1.3 | 1.6 | 1.7 | 1.5 |
| | 40+ times | | .1 | .2 | .8 | 1.5 | 2.1 | 2.3 | 2.9 |
| Cocaine | No Answer | | 2.0 | 1.4 | 1.9 | 1.8 | 1.8 | 1.8 | 1.9 |
| | Never | | 97.8 | 98.3 | 97.3 | 97.3 | 97.2 | 97.1 | 97.0 |
| | 1-5 times | | .1 | .2 | .5 | .6 | .7 | .7 | .8 |
| | 6-19 times | | .0 | .0 | .1 | .2 | .2 | .2 | .2 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .0 | .1 | .1 |
| | 40+ times | | .0 | .0 | .0 | .1 | .2 | .0 | .1 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------|-------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Crack | No Answer | 1.9 | 1.5 | 2.0 | 1.9 | 1.7 | 1.8 | 1.9 |
| | Never | 97.8 | 98.1 | 97.3 | 97.5 | 97.7 | 97.6 | 97.5 |
| | 1-5 times | .2 | .3 | .5 | .4 | .3 | .3 | .3 |
| | 6-19 times | .0 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 20-40 times | .0 | .1 | .1 | .1 | .0 | .1 | .0 |
| | 40+ times | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| Inhalants | No Answer | 2.6 | 1.6 | 2.1 | 1.9 | 1.7 | 1.9 | 1.9 |
| | Never | 96.6 | 96.9 | 95.4 | 95.9 | 96.4 | 96.6 | 97.1 |
| | 1-5 times | .5 | 1.2 | 1.8 | 1.7 | 1.4 | 1.1 | .7 |
| | 6-19 times | .2 | .2 | .4 | .3 | .3 | .3 | .2 |
| | 20-40 times | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 40+ times | .0 | .1 | .1 | .1 | .1 | .1 | .1 |
| Amphetamines | No Answer | 3.0 | 1.8 | 2.1 | 1.9 | 1.9 | 1.8 | 1.8 |
| | Never | 96.9 | 97.8 | 96.7 | 96.2 | 96.0 | 95.7 | 96.2 |
| | 1-5 times | .1 | .3 | .9 | 1.1 | 1.4 | 1.5 | 1.3 |
| | 6-19 times | .0 | .1 | .2 | .4 | .4 | .5 | .5 |
| | 20-40 times | .0 | .0 | .1 | .2 | .1 | .3 | .2 |
| | 40+ times | .0 | .0 | .0 | .1 | .2 | .2 | .2 |
| Methamphetamines | No Answer | 2.6 | 1.6 | 2.2 | 1.9 | 1.8 | 1.8 | 1.8 |
| | Never | 97.1 | 98.0 | 97.1 | 97.4 | 97.6 | 97.5 | 97.4 |
| | 1-5 times | .1 | .2 | .5 | .3 | .4 | .4 | .4 |
| | 6-19 times | .0 | .1 | .1 | .2 | .1 | .2 | .2 |
| | 20-40 times | .0 | .0 | .0 | .1 | .1 | .0 | .1 |
| | 40+ times | .0 | .0 | .0 | .1 | .1 | .1 | .1 |
| Heroin | No Answer | 2.5 | 1.5 | 2.2 | 1.9 | 1.9 | 1.9 | 1.9 |
| | Never | 97.4 | 98.1 | 97.3 | 97.5 | 97.6 | 97.5 | 97.5 |
| | 1-5 times | .1 | .2 | .4 | .3 | .3 | .4 | .2 |
| | 6-19 times | .0 | .0 | .1 | .2 | .1 | .1 | .2 |
| | 20-40 times | .0 | .0 | .0 | .1 | .1 | .1 | .1 |
| | 40+ times | .0 | .1 | .0 | .1 | .1 | .1 | .0 |
| Steroids | No Answer | 2.1 | 1.5 | 2.2 | 2.0 | 1.9 | 1.9 | 1.9 |
| | Never | 97.7 | 98.2 | 97.4 | 97.6 | 97.8 | 97.7 | 97.7 |
| | 1-5 times | .1 | .2 | .3 | .2 | .2 | .3 | .2 |
| | 6-19 times | .0 | .1 | .1 | .1 | .1 | .1 | .2 |
| | 20-40 times | .0 | .0 | .0 | .1 | .0 | .0 | .1 |
| | 40+ times | .0 | .0 | .0 | .0 | .0 | .0 | .0 |
| Ecstasy | No Answer | 3.1 | 1.8 | 2.2 | 1.9 | 2.0 | 1.9 | 2.0 |
| | Never | 96.7 | 97.8 | 96.9 | 96.9 | 96.7 | 96.9 | 96.9 |
| | 1-5 times | .2 | .3 | .7 | .9 | 1.1 | .9 | .9 |
| | 6-19 times | .0 | .1 | .2 | .2 | .1 | .2 | .1 |
| | 20-40 times | .0 | .0 | .0 | .1 | .1 | .1 | .1 |
| | 40+ times | .0 | .0 | .0 | .1 | .0 | .0 | .1 |

**Monthly Use of Alcohol, Tobacco, and Other Drugs
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|--------------------------|-------------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Tranquilizers | No Answer | | 2.9 | 1.7 | 2.2 | 2.1 | 2.4 | 1.9 | 1.9 |
| | Never | | 96.9 | 97.8 | 96.8 | 96.3 | 95.8 | 96.4 | 96.5 |
| | 1-5 times | | .2 | .3 | .8 | 1.1 | 1.1 | 1.2 | 1.1 |
| | 6-19 times | | .1 | .0 | .2 | .3 | .3 | .4 | .3 |
| | 20-40 times | | .0 | .0 | .0 | .1 | .1 | .1 | .1 |
| | 40+ times | | .0 | .0 | .0 | .1 | .2 | .0 | .1 |
| Hallucinogens | No Answer | | 3.2 | 1.8 | 2.3 | 2.0 | 2.1 | 2.0 | 2.0 |
| | Never | | 96.6 | 97.8 | 96.7 | 96.8 | 96.6 | 96.6 | 96.7 |
| | 1-5 times | | .1 | .3 | .6 | .8 | 1.0 | 1.1 | .8 |
| | 6-19 times | | .0 | .1 | .2 | .3 | .2 | .2 | .3 |
| | 20-40 times | | .0 | .0 | .1 | .1 | .1 | .1 | .1 |
| | 40+ times | | .0 | .0 | .1 | .0 | .1 | .0 | .1 |
| Prescription painkillers | No Answer | | 2.2 | 1.6 | 2.3 | 2.1 | 1.9 | 1.9 | 1.9 |
| | Never | | 97.2 | 97.0 | 94.4 | 93.0 | 92.6 | 93.0 | 93.3 |
| | 1-5 times | | .4 | 1.0 | 2.4 | 3.3 | 4.0 | 3.4 | 3.0 |
| | 6-19 times | | .1 | .2 | .7 | .9 | .9 | .9 | 1.1 |
| | 20-40 times | | .1 | .1 | .2 | .4 | .4 | .5 | .4 |
| | 40+ times | | .1 | .0 | .1 | .3 | .2 | .3 | .3 |
| Prescription drugs | No Answer | | 2.3 | 1.6 | 2.4 | 2.1 | 1.9 | 2.0 | 2.0 |
| | Never | | 97.1 | 97.0 | 94.6 | 93.3 | 93.0 | 92.8 | 93.1 |
| | 1-5 times | | .4 | 1.1 | 2.1 | 3.0 | 3.5 | 3.5 | 3.2 |
| | 6-19 times | | .1 | .2 | .6 | .9 | 1.0 | 1.0 | 1.0 |
| | 20-40 times | | .1 | .1 | .2 | .3 | .3 | .5 | .4 |
| | 40+ times | | .1 | .1 | .1 | .3 | .3 | .3 | .3 |
| Over the counter drugs | No Answer | | 2.4 | 1.9 | 2.5 | 2.4 | 2.2 | 2.0 | 2.3 |
| | Never | | 96.9 | 96.3 | 94.3 | 93.5 | 94.3 | 94.8 | 95.3 |
| | 1-5 times | | .5 | 1.4 | 2.4 | 2.9 | 2.5 | 2.2 | 1.6 |
| | 6-19 times | | .1 | .3 | .5 | .6 | .6 | .5 | .4 |
| | 20-40 times | | .0 | .1 | .2 | .4 | .2 | .3 | .2 |
| | 40+ times | | .0 | .1 | .1 | .2 | .2 | .2 | .2 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cigarettes | No Answer | 1.7 | 1.7 | 1.7 | 1.8 | 2.1 | 1.8 | 2.0 |
| | Never | 92.4 | 86.6 | 78.9 | 73.0 | 68.1 | 63.5 | 60.6 |
| | 10 years or younger | 2.9 | 3.7 | 4.2 | 3.9 | 3.7 | 3.8 | 2.6 |
| | 11 years | 1.9 | 2.7 | 3.0 | 2.4 | 2.4 | 1.9 | 1.5 |
| | 12 years | .9 | 3.2 | 3.7 | 3.7 | 3.4 | 3.2 | 2.4 |
| | 13 years | .2 | 1.9 | 5.8 | 5.7 | 4.8 | 4.0 | 3.4 |
| | 14 years | .0 | .2 | 2.4 | 6.5 | 5.3 | 5.0 | 4.3 |
| | 15 years | .0 | .0 | .2 | 2.7 | 7.4 | 6.8 | 6.3 |
| | 16 years | .0 | .0 | .0 | .3 | 2.6 | 7.7 | 7.8 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 2.3 | 9.2 |
| | | | | | | | | |
| Smokeless Tobacco | No Answer | 2.0 | 1.8 | 2.1 | 2.1 | 2.4 | 2.2 | 2.6 |
| | Never | 97.3 | 96.1 | 94.3 | 92.2 | 90.3 | 88.9 | 87.8 |
| | 10 years or younger | .4 | .5 | .6 | .4 | .5 | .5 | .3 |
| | 11 years | .3 | .4 | .6 | .3 | .2 | .2 | .2 |
| | 12 years | .1 | .7 | .7 | .6 | .5 | .4 | .3 |
| | 13 years | .0 | .5 | 1.2 | 1.1 | 1.1 | .8 | .6 |
| | 14 years | .0 | .1 | .5 | 2.1 | 1.7 | 1.3 | 1.0 |
| | 15 years | .0 | .0 | .1 | 1.0 | 2.4 | 2.0 | 1.8 |
| | 16 years | .0 | .0 | .0 | .1 | 1.0 | 2.7 | 2.7 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .9 | 2.7 |
| | | | | | | | | |
| Cigars | No Answer | 2.1 | 1.9 | 2.1 | 2.4 | 2.7 | 2.6 | 2.7 |
| | Never | 97.3 | 96.2 | 93.6 | 89.7 | 85.6 | 81.8 | 76.2 |
| | 10 years or younger | .3 | .5 | .6 | .5 | .5 | .6 | .4 |
| | 11 years | .2 | .4 | .6 | .6 | .4 | .4 | .3 |
| | 12 years | .1 | .5 | .8 | .8 | .7 | .3 | .6 |
| | 13 years | .1 | .4 | 1.5 | 1.7 | 1.6 | 1.5 | 1.0 |
| | 14 years | .0 | .0 | .7 | 2.7 | 2.6 | 2.0 | 1.6 |
| | 15 years | .0 | .0 | .1 | 1.4 | 4.2 | 3.9 | 3.5 |
| | 16 years | .0 | .0 | .0 | .1 | 1.6 | 5.3 | 6.1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.6 | 7.6 |
| | | | | | | | | |
| Pipe | No Answer | 2.3 | 2.1 | 2.6 | 2.9 | 3.1 | 3.1 | 3.3 |
| | Never | 97.3 | 96.4 | 94.1 | 90.9 | 87.5 | 83.6 | 78.4 |
| | 10 years or younger | .2 | .2 | .2 | .3 | .3 | .3 | .3 |
| | 11 years | .1 | .3 | .4 | .3 | .2 | .2 | .2 |
| | 12 years | .1 | .6 | .7 | .5 | .5 | .5 | .3 |
| | 13 years | .0 | .3 | 1.2 | 1.5 | 1.1 | .8 | .7 |
| | 14 years | .0 | .0 | .7 | 2.3 | 2.2 | 1.7 | 1.4 |
| | 15 years | .0 | .0 | .1 | 1.2 | 3.4 | 3.4 | 2.5 |
| | 16 years | .0 | .0 | .0 | .1 | 1.6 | 4.9 | 4.4 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.5 | 8.5 |
| | | | | | | | | |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-----------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Alcohol | No Answer | 2.7 | 2.5 | 2.8 | 2.7 | 2.8 | 2.7 | 2.6 |
| | Never | 84.1 | 75.5 | 61.9 | 51.7 | 44.4 | 40.0 | 35.1 |
| | 10 years or younger | 6.8 | 6.6 | 6.4 | 5.1 | 4.2 | 3.8 | 2.4 |
| | 11 years | 4.0 | 5.1 | 4.7 | 3.3 | 2.3 | 1.7 | 1.3 |
| | 12 years | 2.0 | 6.6 | 7.8 | 5.9 | 4.3 | 3.9 | 2.6 |
| | 13 years | .3 | 3.4 | 11.3 | 11.4 | 8.7 | 6.3 | 5.6 |
| | 14 years | .0 | .3 | 4.7 | 13.6 | 12.1 | 9.7 | 8.4 |
| | 15 years | .0 | .0 | .4 | 5.8 | 15.9 | 15.3 | 13.9 |
| | 16 years | .0 | .0 | .0 | .5 | 5.2 | 13.0 | 16.2 |
| | 17 years or older | .0 | .0 | .0 | .0 | .2 | 3.6 | 12.0 |
| Marijuana | No Answer | 2.1 | 1.9 | 2.3 | 2.4 | 2.6 | 2.3 | 2.6 |
| | Never | 96.1 | 92.9 | 85.5 | 79.0 | 72.1 | 67.7 | 65.4 |
| | 10 years or younger | .4 | .7 | 1.0 | .8 | .9 | .8 | .5 |
| | 11 years | .7 | 1.0 | 1.2 | 1.0 | .8 | .6 | .6 |
| | 12 years | .5 | 1.9 | 2.4 | 2.0 | 1.8 | 1.6 | 1.4 |
| | 13 years | .2 | 1.4 | 5.0 | 4.5 | 3.9 | 3.0 | 2.5 |
| | 14 years | .0 | .2 | 2.4 | 7.1 | 5.5 | 4.8 | 3.4 |
| | 15 years | .0 | .0 | .3 | 3.2 | 9.0 | 7.8 | 6.2 |
| | 16 years | .0 | .0 | .0 | .2 | 3.4 | 8.9 | 8.8 |
| | 17 years or older | .0 | .0 | .0 | .0 | .2 | 2.4 | 8.7 |
| Cocaine | No Answer | 2.2 | 1.9 | 2.1 | 2.3 | 2.6 | 2.4 | 2.5 |
| | Never | 97.6 | 97.8 | 96.6 | 96.4 | 95.5 | 95.1 | 94.6 |
| | 10 years or younger | .1 | .0 | .2 | .1 | .1 | .1 | .1 |
| | 11 years | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .1 | .3 | .2 | .1 | .1 | .0 |
| | 13 years | .0 | .1 | .3 | .2 | .2 | .2 | .2 |
| | 14 years | .0 | .0 | .3 | .5 | .4 | .3 | .3 |
| | 15 years | .0 | .0 | .1 | .2 | .5 | .6 | .4 |
| | 16 years | .0 | .0 | .0 | .0 | .4 | .8 | .7 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .4 | 1.0 |
| Crack | No Answer | 2.3 | 1.9 | 2.2 | 2.4 | 2.6 | 2.3 | 2.5 |
| | Never | 97.4 | 97.7 | 96.8 | 96.6 | 96.5 | 96.6 | 96.4 |
| | 10 years or younger | .1 | .1 | .2 | .1 | .1 | .1 | .1 |
| | 11 years | .1 | .1 | .2 | .1 | .0 | .1 | .1 |
| | 12 years | .0 | .1 | .1 | .1 | .1 | .1 | .0 |
| | 13 years | .0 | .1 | .2 | .2 | .1 | .1 | .1 |
| | 14 years | .0 | .0 | .2 | .3 | .2 | .2 | .2 |
| | 15 years | .0 | .0 | .0 | .1 | .3 | .2 | .2 |
| | 16 years | .0 | .0 | .0 | .0 | .1 | .2 | .2 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .1 | .2 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Inhalants | No Answer | 3.0 | 2.2 | 2.2 | 2.4 | 2.6 | 2.3 | 2.6 |
| | Never | 96.0 | 95.6 | 93.7 | 93.2 | 92.5 | 92.7 | 93.0 |
| | 10 years or younger | .5 | .4 | .4 | .3 | .3 | .3 | .2 |
| | 11 years | .3 | .6 | .6 | .4 | .3 | .3 | .2 |
| | 12 years | .2 | .7 | .9 | .6 | .4 | .4 | .4 |
| | 13 years | .0 | .5 | 1.4 | 1.1 | .9 | .8 | .5 |
| | 14 years | .0 | .0 | .6 | 1.4 | 1.2 | .9 | .6 |
| | 15 years | .0 | .0 | .1 | .5 | 1.3 | 1.1 | .7 |
| | 16 years | .0 | .0 | .0 | .0 | .5 | 1.0 | .9 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .2 | .8 |
| Amphetamines | No Answer | 3.3 | 2.3 | 2.3 | 2.5 | 2.5 | 2.3 | 2.6 |
| | Never | 96.5 | 97.2 | 96.0 | 94.8 | 93.5 | 92.4 | 92.4 |
| | 10 years or younger | .1 | .0 | .1 | .1 | .1 | .2 | .2 |
| | 11 years | .0 | .1 | .2 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .2 | .3 | .2 | .2 | .2 | .2 |
| | 13 years | .0 | .2 | .6 | .7 | .7 | .6 | .4 |
| | 14 years | .0 | .0 | .4 | 1.1 | 1.0 | .8 | .6 |
| | 15 years | .0 | .0 | .1 | .4 | 1.3 | 1.4 | 1.1 |
| | 16 years | .0 | .0 | .0 | .1 | .5 | 1.3 | 1.4 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .4 | 1.1 |
| Methamphetamines | No Answer | 3.0 | 2.1 | 2.2 | 2.4 | 2.6 | 2.4 | 2.5 |
| | Never | 96.9 | 97.4 | 96.8 | 96.7 | 96.3 | 96.2 | 96.0 |
| | 10 years or younger | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 11 years | .0 | .1 | .1 | .1 | .0 | .1 | .1 |
| | 12 years | .0 | .1 | .2 | .1 | .1 | .0 | .0 |
| | 13 years | .0 | .1 | .3 | .2 | .1 | .1 | .1 |
| | 14 years | .0 | .0 | .2 | .3 | .2 | .2 | .1 |
| | 15 years | .0 | .0 | .0 | .2 | .3 | .3 | .2 |
| | 16 years | .0 | .0 | .0 | .0 | .2 | .4 | .2 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .2 | .6 |
| Heroin | No Answer | 2.8 | 2.0 | 2.3 | 2.5 | 2.6 | 2.4 | 2.5 |
| | Never | 97.0 | 97.6 | 97.0 | 96.8 | 96.6 | 96.7 | 96.4 |
| | 10 years or younger | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 11 years | .0 | .1 | .1 | .1 | .1 | .1 | .0 |
| | 12 years | .0 | .1 | .1 | .1 | .1 | .1 | .0 |
| | 13 years | .0 | .1 | .2 | .2 | .1 | .1 | .1 |
| | 14 years | .0 | .0 | .1 | .1 | .1 | .1 | .2 |
| | 15 years | .0 | .0 | .0 | .1 | .2 | .2 | .2 |
| | 16 years | .0 | .0 | .0 | .0 | .2 | .2 | .2 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .1 | .3 |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|---------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Steroids | No Answer | 2.5 | 2.1 | 2.4 | 2.4 | 2.8 | 2.5 | 2.6 |
| | Never | 97.4 | 97.6 | 97.1 | 97.2 | 96.7 | 97.0 | 96.9 |
| | 10 years or younger | .1 | .1 | .1 | .0 | .1 | .1 | .1 |
| | 11 years | .0 | .1 | .1 | .0 | .0 | .1 | .0 |
| | 12 years | .0 | .1 | .1 | .0 | .0 | .1 | .0 |
| | 13 years | .0 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 14 years | .0 | .0 | .1 | .1 | .1 | .0 | .1 |
| | 15 years | .0 | .0 | .0 | .1 | .1 | .1 | .1 |
| | 16 years | .0 | .0 | .0 | .0 | .1 | .1 | .1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .0 | .1 |
| | | | | | | | | |
| Ecstasy | No Answer | 3.6 | 2.3 | 2.4 | 2.5 | 2.7 | 2.5 | 2.5 |
| | Never | 96.3 | 97.2 | 96.2 | 95.4 | 94.6 | 94.2 | 93.7 |
| | 10 years or younger | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 11 years | .0 | .1 | .2 | .1 | .1 | .1 | .0 |
| | 12 years | .0 | .2 | .2 | .2 | .1 | .1 | .1 |
| | 13 years | .0 | .2 | .5 | .4 | .3 | .2 | .3 |
| | 14 years | .0 | .0 | .4 | .8 | .5 | .4 | .2 |
| | 15 years | .0 | .0 | .1 | .5 | 1.0 | .7 | .6 |
| | 16 years | .0 | .0 | .0 | .0 | .5 | 1.2 | 1.3 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .5 | 1.2 |
| | | | | | | | | |
| Tranquilizers | No Answer | 3.3 | 2.2 | 2.5 | 2.7 | 2.7 | 2.6 | 2.7 |
| | Never | 96.5 | 97.2 | 96.2 | 94.9 | 94.2 | 93.8 | 93.7 |
| | 10 years or younger | .1 | .1 | .2 | .1 | .1 | .2 | .1 |
| | 11 years | .0 | .1 | .2 | .1 | .2 | .1 | .1 |
| | 12 years | .1 | .1 | .2 | .2 | .2 | .2 | .2 |
| | 13 years | .0 | .2 | .4 | .6 | .5 | .3 | .2 |
| | 14 years | .0 | .0 | .3 | 1.0 | .7 | .6 | .5 |
| | 15 years | .0 | .0 | .0 | .3 | 1.0 | 1.3 | .7 |
| | 16 years | .0 | .0 | .0 | .0 | .3 | .7 | 1.0 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .2 | .7 |
| | | | | | | | | |
| Hallucinogens | No Answer | 3.7 | 2.3 | 2.6 | 2.7 | 2.9 | 2.7 | 2.9 |
| | Never | 96.2 | 97.2 | 96.0 | 95.3 | 94.4 | 93.9 | 93.8 |
| | 10 years or younger | .1 | .1 | .1 | .1 | .1 | .2 | .1 |
| | 11 years | .0 | .1 | .1 | .1 | .1 | .1 | .1 |
| | 12 years | .0 | .1 | .2 | .2 | .2 | .1 | .1 |
| | 13 years | .0 | .1 | .4 | .4 | .3 | .2 | .2 |
| | 14 years | .0 | .0 | .4 | .8 | .6 | .3 | .3 |
| | 15 years | .0 | .0 | .1 | .4 | .9 | .8 | .4 |
| | 16 years | .0 | .0 | .0 | .0 | .5 | 1.1 | .8 |
| | 17 years or older | .0 | .0 | .0 | .0 | .0 | .6 | 1.3 |
| | | | | | | | | |

**Age Distribution of First Time Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--------------------------|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Prescription painkillers | No Answer | 2.8 | 2.0 | 2.4 | 2.6 | 2.7 | 2.5 | 2.7 |
| | Never | 96.6 | 96.1 | 92.9 | 89.8 | 87.4 | 86.3 | 86.5 |
| | 10 years or younger | .3 | .3 | .3 | .3 | .2 | .3 | .1 |
| | 11 years | .2 | .4 | .5 | .4 | .3 | .2 | .2 |
| | 12 years | .2 | .7 | .8 | .6 | .6 | .5 | .3 |
| | 13 years | .0 | .4 | 1.9 | 1.7 | 1.4 | 1.2 | .7 |
| | 14 years | .0 | .0 | 1.0 | 2.9 | 2.2 | 1.8 | 1.3 |
| | 15 years | .0 | .0 | .2 | 1.6 | 3.5 | 3.2 | 2.2 |
| | 16 years | .0 | .0 | .0 | .1 | 1.5 | 3.0 | 3.2 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.0 | 2.7 |
| Prescription drugs | No Answer | 2.7 | 2.3 | 2.5 | 2.7 | 2.8 | 2.5 | 2.6 |
| | Never | 96.5 | 95.9 | 93.4 | 90.4 | 88.1 | 86.7 | 86.5 |
| | 10 years or younger | .4 | .3 | .2 | .3 | .2 | .3 | .1 |
| | 11 years | .1 | .4 | .5 | .4 | .2 | .2 | .2 |
| | 12 years | .1 | .6 | .6 | .6 | .5 | .4 | .2 |
| | 13 years | .1 | .5 | 1.7 | 1.5 | 1.2 | 1.0 | .6 |
| | 14 years | .0 | .0 | .9 | 2.8 | 2.0 | 1.8 | 1.4 |
| | 15 years | .0 | .0 | .1 | 1.3 | 3.4 | 2.6 | 2.2 |
| | 16 years | .0 | .0 | .0 | .1 | 1.4 | 3.2 | 3.1 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | 1.1 | 3.0 |
| Over the counter drugs | No Answer | 2.9 | 2.5 | 2.7 | 3.0 | 3.0 | 2.7 | 2.9 |
| | Never | 96.3 | 95.3 | 92.7 | 90.9 | 90.1 | 90.2 | 90.7 |
| | 10 years or younger | .3 | .3 | .4 | .5 | .2 | .3 | .2 |
| | 11 years | .2 | .3 | .5 | .3 | .3 | .1 | .2 |
| | 12 years | .3 | .8 | .9 | .7 | .6 | .4 | .3 |
| | 13 years | .0 | .7 | 1.7 | 1.4 | 1.2 | .8 | .6 |
| | 14 years | .0 | .1 | .9 | 2.1 | 1.6 | 1.3 | 1.0 |
| | 15 years | .0 | .0 | .1 | 1.1 | 2.1 | 1.8 | 1.5 |
| | 16 years | .0 | .0 | .0 | .1 | .9 | 1.8 | 1.5 |
| | 17 years or older | .0 | .0 | .0 | .0 | .1 | .5 | 1.2 |

**Perceived Availability of Drugs
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------------------|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Availability of cigarettes | No Answer | 8.4 | 6.9 | 6.1 | 5.2 | 4.2 | 3.8 | 3.4 |
| | Very difficult | 63.3 | 54.1 | 42.6 | 33.5 | 27.6 | 21.2 | 10.5 |
| | Fairly difficult | 10.1 | 13.0 | 13.7 | 13.1 | 11.9 | 9.5 | 4.0 |
| | Fairly easy | 9.4 | 13.7 | 19.1 | 23.1 | 24.1 | 23.4 | 12.4 |
| | Easy | 8.7 | 12.3 | 18.6 | 25.0 | 32.2 | 42.1 | 69.7 |
| Availability of alcohol | No Answer | 8.7 | 7.0 | 6.1 | 5.1 | 4.0 | 3.7 | 3.5 |
| | Very difficult | 55.1 | 43.4 | 32.4 | 23.4 | 18.5 | 16.0 | 12.0 |
| | Fairly difficult | 14.3 | 16.7 | 15.9 | 14.5 | 12.6 | 10.1 | 10.3 |
| | Fairly easy | 11.5 | 17.6 | 21.9 | 26.3 | 29.2 | 30.3 | 34.3 |
| | Easy | 10.2 | 15.3 | 23.7 | 30.7 | 35.8 | 39.8 | 40.0 |
| Availability of marijuana | No Answer | 9.4 | 7.7 | 7.0 | 6.0 | 5.0 | 4.4 | 4.1 |
| | Very difficult | 83.1 | 77.0 | 63.6 | 48.5 | 39.1 | 33.2 | 27.6 |
| | Fairly difficult | 3.5 | 5.8 | 9.3 | 11.6 | 11.6 | 11.3 | 12.7 |
| | Fairly easy | 1.7 | 4.5 | 9.6 | 15.5 | 19.0 | 21.1 | 24.4 |
| | Easy | 2.4 | 4.9 | 10.5 | 18.3 | 25.4 | 30.1 | 31.2 |
| Availability of drugs | No Answer | 9.5 | 7.8 | 7.2 | 6.6 | 5.5 | 4.9 | 4.6 |
| | Very difficult | 85.9 | 83.0 | 75.8 | 66.4 | 60.7 | 56.0 | 51.7 |
| | Fairly difficult | 2.3 | 4.8 | 8.8 | 14.1 | 17.2 | 18.9 | 21.6 |
| | Fairly easy | 1.0 | 2.3 | 4.7 | 7.5 | 9.9 | 12.1 | 13.1 |
| | Easy | 1.3 | 2.1 | 3.5 | 5.5 | 6.7 | 8.0 | 8.9 |

**Binge Drinking in the Past Two Weeks
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|----------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Binge drinking | No Answer | 2.6 | 1.9 | 1.5 | 1.9 | 1.6 | 1.8 | 1.6 |
| | None | 92.0 | 90.1 | 86.7 | 83.0 | 81.8 | 80.1 | 75.4 |
| | Once | 2.5 | 3.3 | 4.7 | 6.3 | 6.7 | 7.7 | 9.8 |
| | Twice | 1.1 | 1.9 | 3.0 | 3.7 | 4.2 | 4.1 | 5.5 |
| | 3-5 times | 1.0 | 1.6 | 2.7 | 3.4 | 3.6 | 4.1 | 5.3 |
| | 6-9 times | .4 | .5 | .7 | .9 | 1.2 | 1.2 | 1.5 |
| | 10+ times | .4 | .6 | .6 | .8 | .9 | 1.0 | .9 |

**Main Sources of Alcoholic Beverages in the Past Month
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------------|---|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Main sources of alcoholic beverages | No Answer | 5.8 | 4.6 | 4.5 | 5.0 | 5.2 | 5.5 | 6.2 |
| | No drink | 89.4 | 85.2 | 78.0 | 71.4 | 68.5 | 66.7 | 60.7 |
| | Restaurants/bars/clubs | .0 | .0 | .1 | .0 | .1 | .2 | .2 |
| | Public events | .0 | .0 | .0 | .0 | .0 | .0 | .1 |
| | Had someone else buy it | .3 | .8 | 1.9 | 4.2 | 6.3 | 7.7 | 10.9 |
| | Liquor stores/convenience stores/supermarkets | .1 | .0 | .0 | .1 | .1 | .2 | .4 |
| | Received from person 21 or older | .6 | 1.4 | 2.4 | 4.1 | 5.6 | 7.5 | 10.8 |
| | Took it from a store | .1 | .1 | .2 | .2 | .1 | .3 | .2 |
| | Family members | 1.9 | 3.7 | 5.6 | 6.2 | 5.7 | 4.8 | 4.3 |
| | Other ways | 1.9 | 4.2 | 7.2 | 8.8 | 8.4 | 7.0 | 6.4 |

**Most Important Reasons for Drinking
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | | Grade | | | | | | |
|---|-----------|--|-------|------|------|------|------|------|------|
| | | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Did not drink alcoholic beverages | No answer | | 15.8 | 24.8 | 37.8 | 47.8 | 53.5 | 56.9 | 62.0 |
| | Yes | | 84.2 | 75.2 | 62.2 | 52.2 | 46.5 | 43.1 | 38.0 |
| To experiment | No answer | | 90.5 | 83.8 | 76.3 | 73.2 | 73.1 | 74.8 | 76.4 |
| | Yes | | 9.5 | 16.2 | 23.7 | 26.8 | 26.9 | 25.2 | 23.6 |
| To relax or relieve tension | No answer | | 98.8 | 96.2 | 91.9 | 87.6 | 84.5 | 81.8 | 79.9 |
| | Yes | | 1.2 | 3.8 | 8.1 | 12.4 | 15.5 | 18.2 | 20.1 |
| To feel good or get high | No answer | | 99.0 | 97.2 | 94.0 | 90.0 | 87.7 | 85.6 | 83.7 |
| | Yes | | 1.0 | 2.8 | 6.0 | 10.0 | 12.3 | 14.4 | 16.3 |
| To seek deeper insights and understanding | No answer | | 99.5 | 99.1 | 98.5 | 97.9 | 98.2 | 98.1 | 97.9 |
| | Yes | | .5 | .9 | 1.5 | 2.1 | 1.8 | 1.9 | 2.1 |
| To have a good time with friends | No answer | | 98.0 | 93.6 | 86.4 | 76.7 | 70.6 | 65.8 | 59.6 |
| | Yes | | 2.0 | 6.4 | 13.6 | 23.3 | 29.4 | 34.2 | 40.4 |
| To fit in with a group I like | No answer | | 99.2 | 98.2 | 97.6 | 97.2 | 97.3 | 97.4 | 97.2 |
| | Yes | | .8 | 1.8 | 2.4 | 2.8 | 2.7 | 2.6 | 2.8 |
| To get away from my problems | No answer | | 98.0 | 94.7 | 91.0 | 87.8 | 87.1 | 87.4 | 88.2 |
| | Yes | | 2.0 | 5.3 | 9.0 | 12.2 | 12.9 | 12.6 | 11.8 |
| Because of boredom | No answer | | 98.1 | 95.2 | 91.6 | 89.6 | 88.3 | 88.2 | 87.7 |
| | Yes | | 1.9 | 4.8 | 8.4 | 10.4 | 11.7 | 11.8 | 12.3 |
| Because of anger | No answer | | 98.3 | 96.3 | 93.9 | 92.4 | 92.2 | 92.4 | 93.0 |
| | Yes | | 1.7 | 3.7 | 6.1 | 7.6 | 7.8 | 7.6 | 7.0 |
| To get through the day | No answer | | 99.4 | 98.6 | 97.7 | 97.2 | 97.5 | 97.8 | 97.9 |
| | Yes | | .6 | 1.4 | 2.3 | 2.8 | 2.5 | 2.2 | 2.1 |
| To increase the effects of other drugs | No answer | | 99.8 | 99.5 | 99.1 | 98.5 | 98.1 | 98.0 | 98.1 |
| | Yes | | .2 | .5 | .9 | 1.5 | 1.9 | 2.0 | 1.9 |
| To decrease the effects of other drugs | No answer | | 99.9 | 99.9 | 99.7 | 99.8 | 99.7 | 99.7 | 99.6 |
| | Yes | | .1 | .1 | .3 | .2 | .3 | .3 | .4 |
| To get to sleep | No answer | | 99.5 | 99.1 | 98.2 | 97.9 | 97.6 | 97.5 | 97.0 |
| | Yes | | .5 | .9 | 1.8 | 2.1 | 2.4 | 2.5 | 3.0 |
| Because it tastes good | No answer | | 97.5 | 93.5 | 88.2 | 84.4 | 83.3 | 81.9 | 80.5 |
| | Yes | | 2.5 | 6.5 | 11.8 | 15.6 | 16.7 | 18.1 | 19.5 |
| Because I am hooked | No answer | | 99.6 | 99.2 | 98.9 | 98.8 | 98.8 | 99.0 | 99.1 |
| | Yes | | .4 | .8 | 1.1 | 1.2 | 1.2 | 1.0 | .9 |

**Consequences of Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-----------------------|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Had a hangover | No Answer | 3.3 | 2.8 | 2.1 | 2.7 | 2.5 | 2.1 | 2.1 |
| | Never | 94.2 | 89.4 | 82.2 | 73.8 | 68.9 | 65.2 | 60.4 |
| | Once | 1.5 | 3.7 | 6.6 | 9.4 | 10.9 | 12.7 | 12.8 |
| | Twice | .4 | 1.9 | 3.5 | 5.2 | 6.3 | 6.9 | 7.6 |
| | 3-5 times | .4 | 1.3 | 2.7 | 4.8 | 6.1 | 7.1 | 9.3 |
| | 6-10 times | .1 | .3 | 1.2 | 1.7 | 2.5 | 2.6 | 3.9 |
| | 11 or more times | .2 | .7 | 1.6 | 2.4 | 2.7 | 3.3 | 4.0 |
| | | | | | | | | |
| Had nausea or vomited | No Answer | 3.4 | 3.0 | 2.3 | 2.8 | 2.7 | 2.4 | 2.4 |
| | Never | 93.8 | 90.4 | 85.4 | 78.5 | 73.4 | 68.6 | 63.6 |
| | Once | 1.7 | 3.7 | 6.3 | 9.0 | 11.2 | 12.8 | 13.5 |
| | Twice | .5 | 1.4 | 2.5 | 3.8 | 4.8 | 5.9 | 7.5 |
| | 3-5 times | .4 | .7 | 2.0 | 3.5 | 4.7 | 6.3 | 8.2 |
| | 6-10 times | .1 | .4 | .8 | 1.4 | 1.7 | 2.4 | 2.7 |
| | 11 or more times | .1 | .4 | .7 | 1.1 | 1.5 | 1.7 | 2.1 |
| | | | | | | | | |
| Had a memory loss | No Answer | 3.5 | 3.2 | 2.5 | 3.1 | 2.9 | 2.5 | 2.5 |
| | Never | 94.8 | 92.7 | 88.6 | 82.8 | 78.2 | 76.4 | 73.3 |
| | Once | 1.0 | 2.3 | 4.1 | 6.3 | 7.8 | 8.4 | 8.9 |
| | Twice | .3 | .7 | 1.8 | 2.7 | 3.7 | 4.1 | 5.0 |
| | 3-5 times | .3 | .6 | 1.6 | 2.7 | 3.9 | 4.6 | 5.6 |
| | 6-10 times | .0 | .2 | .7 | 1.1 | 1.6 | 2.0 | 2.3 |
| | 11 or more times | .1 | .3 | .8 | 1.3 | 1.9 | 2.0 | 2.4 |
| | | | | | | | | |
| Poor on school test | No Answer | 3.7 | 3.2 | 2.7 | 3.0 | 3.0 | 2.6 | 2.5 |
| | Never | 94.2 | 92.6 | 91.6 | 90.4 | 90.1 | 90.9 | 91.9 |
| | Once | 1.1 | 2.0 | 2.4 | 2.4 | 2.2 | 2.2 | 1.7 |
| | Twice | .4 | .7 | 1.0 | 1.2 | 1.4 | 1.2 | 1.1 |
| | 3-5 times | .3 | .7 | 1.0 | 1.4 | 1.5 | 1.5 | 1.2 |
| | 6-10 times | .1 | .2 | .5 | .6 | .7 | .7 | .6 |
| | 11 or more times | .2 | .5 | .8 | 1.0 | 1.2 | 1.0 | .9 |
| | | | | | | | | |
| Missed school | No Answer | 3.7 | 3.3 | 2.6 | 3.0 | 3.0 | 2.6 | 2.5 |
| | Never | 94.3 | 93.5 | 92.7 | 91.3 | 90.7 | 90.9 | 90.9 |
| | Once | .8 | 1.2 | 1.7 | 1.9 | 2.2 | 2.4 | 2.3 |
| | Twice | .4 | .6 | .9 | 1.1 | 1.3 | 1.1 | 1.2 |
| | 3-5 times | .4 | .7 | .9 | 1.0 | 1.3 | 1.3 | 1.4 |
| | 6-10 times | .1 | .2 | .5 | .6 | .7 | .7 | .7 |
| | 11 or more times | .3 | .5 | .7 | .9 | .9 | .9 | 1.0 |
| | | | | | | | | |

**Consequences of Alcohol, Tobacco, and Other Drug Use
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|--|------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Got into a fight | No Answer | 3.7 | 3.3 | 2.8 | 3.1 | 3.0 | 2.6 | 2.7 |
| | Never | 92.8 | 90.6 | 88.0 | 85.6 | 84.9 | 85.2 | 84.0 |
| | Once | 1.6 | 2.6 | 3.8 | 4.9 | 4.8 | 4.7 | 5.6 |
| | Twice | .7 | 1.2 | 1.9 | 2.2 | 2.7 | 2.7 | 3.1 |
| | 3-5 times | .5 | 1.0 | 1.6 | 1.9 | 2.2 | 2.7 | 2.6 |
| | 6-10 times | .2 | .5 | .8 | .9 | .9 | .8 | 1.0 |
| | 11 or more times | .4 | .9 | 1.1 | 1.3 | 1.4 | 1.2 | 1.1 |
| | | | | | | | | |
| Damaged property | No Answer | 3.8 | 3.3 | 2.6 | 2.9 | 2.9 | 2.5 | 2.4 |
| | Never | 95.4 | 94.9 | 94.2 | 93.8 | 94.0 | 94.9 | 94.6 |
| | Once | .4 | .8 | 1.4 | 1.2 | 1.3 | 1.1 | 1.2 |
| | Twice | .2 | .3 | .5 | .7 | .5 | .6 | .6 |
| | 3-5 times | .1 | .3 | .7 | .5 | .6 | .4 | .7 |
| | 6-10 times | .1 | .1 | .3 | .3 | .2 | .2 | .2 |
| | 11 or more times | .1 | .3 | .4 | .5 | .4 | .4 | .4 |
| | | | | | | | | |
| Driven by someone under the influence | No Answer | 3.1 | 2.3 | 1.8 | 2.2 | 2.1 | 2.1 | 1.8 |
| | Yes | 20.1 | 26.1 | 35.2 | 41.4 | 46.9 | 48.8 | 50.0 |
| | No | 76.9 | 71.5 | 63.0 | 56.4 | 51.0 | 49.1 | 48.2 |
| Used alcohol or drugs to relax or fit in | No Answer | 3.6 | 2.8 | 2.2 | 2.3 | 2.3 | 2.4 | 2.0 |
| | Yes | 2.6 | 6.0 | 10.9 | 15.7 | 18.5 | 20.2 | 21.5 |
| | No | 93.9 | 91.1 | 86.9 | 81.9 | 79.1 | 77.4 | 76.5 |
| Used alcohol or drugs alone | No Answer | 3.9 | 3.3 | 2.5 | 2.7 | 2.4 | 2.5 | 2.1 |
| | Yes | 3.0 | 7.2 | 12.5 | 16.3 | 18.1 | 18.7 | 17.5 |
| | No | 93.1 | 89.5 | 85.0 | 81.0 | 79.5 | 78.8 | 80.3 |
| Forgot things you did while high | No Answer | 6.8 | 4.7 | 3.8 | 3.7 | 3.3 | 3.3 | 2.7 |
| | Yes | 1.8 | 4.7 | 9.5 | 15.7 | 19.4 | 22.2 | 25.1 |
| | No | 91.4 | 90.6 | 86.7 | 80.6 | 77.4 | 74.5 | 72.2 |
| Had been told to cut down | No Answer | 8.2 | 5.9 | 4.3 | 3.9 | 3.3 | 3.2 | 2.6 |
| | Yes | 1.7 | 3.1 | 4.8 | 6.7 | 6.9 | 7.1 | 6.4 |
| | No | 90.1 | 91.0 | 90.9 | 89.4 | 89.8 | 89.7 | 91.0 |
| Got into trouble | No Answer | 8.7 | 5.9 | 4.1 | 3.8 | 3.3 | 3.2 | 2.6 |
| | Yes | 1.9 | 3.8 | 6.7 | 8.9 | 10.6 | 11.7 | 10.8 |
| | No | 89.4 | 90.4 | 89.2 | 87.3 | 86.1 | 85.1 | 86.5 |

**Gambling during the Last Twelve Months
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Played cards | No Answer | 2.0 | 1.9 | 1.9 | 2.1 | 2.4 | 2.2 | 2.1 |
| | Not at all | 87.6 | 85.2 | 83.3 | 83.1 | 83.1 | 84.7 | 83.3 |
| | Less than once a month | 5.9 | 7.4 | 8.5 | 8.5 | 8.3 | 7.6 | 8.4 |
| | About once a month | 2.5 | 3.3 | 3.9 | 4.0 | 3.9 | 3.5 | 3.7 |
| | About once a week | 1.4 | 1.6 | 1.8 | 1.7 | 1.7 | 1.5 | 1.8 |
| | Daily | .6 | .6 | .6 | .6 | .6 | .6 | .7 |
| Bet on games | No Answer | 2.3 | 2.1 | 2.1 | 2.2 | 2.7 | 2.4 | 2.3 |
| | Not at all | 87.3 | 84.5 | 82.8 | 82.1 | 83.2 | 84.9 | 84.0 |
| | Less than once a month | 5.4 | 6.6 | 7.5 | 7.7 | 6.9 | 6.4 | 7.0 |
| | About once a month | 2.5 | 3.5 | 4.2 | 4.3 | 3.9 | 3.4 | 3.6 |
| | About once a week | 1.4 | 2.0 | 2.2 | 2.1 | 2.1 | 1.9 | 1.9 |
| | Daily | 1.1 | 1.4 | 1.2 | 1.5 | 1.3 | 1.0 | 1.1 |
| Bet on sports | No Answer | 2.7 | 2.2 | 2.4 | 2.4 | 2.8 | 2.5 | 2.4 |
| | Not at all | 88.7 | 86.7 | 85.3 | 86.6 | 87.9 | 89.8 | 89.9 |
| | Less than once a month | 5.6 | 7.0 | 7.6 | 7.0 | 6.0 | 4.9 | 5.0 |
| | About once a month | 1.6 | 2.3 | 2.6 | 2.1 | 1.6 | 1.3 | 1.4 |
| | About once a week | .9 | 1.2 | 1.2 | .9 | .8 | .6 | .6 |
| | Daily | .6 | .7 | .9 | 1.0 | .9 | .9 | .7 |
| Bought lottery tickets | No Answer | 2.6 | 2.2 | 2.4 | 2.5 | 2.8 | 2.6 | 2.3 |
| | Not at all | 85.5 | 83.9 | 82.9 | 82.9 | 83.6 | 83.8 | 71.0 |
| | Less than once a month | 7.9 | 9.0 | 9.5 | 9.7 | 9.2 | 9.1 | 18.6 |
| | About once a month | 2.4 | 3.1 | 3.4 | 3.3 | 3.0 | 3.2 | 5.6 |
| | About once a week | 1.2 | 1.3 | 1.4 | 1.2 | 1.0 | 1.0 | 1.9 |
| | Daily | .4 | .5 | .4 | .5 | .4 | .4 | .6 |
| Gambled in a casino | No Answer | 2.7 | 2.2 | 2.4 | 2.5 | 2.8 | 2.5 | 2.3 |
| | Not at all | 96.8 | 97.3 | 96.9 | 96.8 | 96.5 | 96.6 | 96.3 |
| | Less than once a month | .2 | .3 | .4 | .3 | .4 | .4 | .8 |
| | About once a month | .1 | .1 | .1 | .1 | .1 | .1 | .1 |
| | About once a week | .1 | .0 | .1 | .1 | .1 | .1 | .1 |
| | Daily | .1 | .1 | .1 | .2 | .2 | .3 | .3 |
| Played online | No Answer | 2.8 | 2.2 | 2.5 | 2.5 | 2.9 | 2.5 | 2.3 |
| | Not at all | 90.5 | 90.3 | 90.1 | 91.1 | 91.4 | 92.8 | 92.6 |
| | Less than once a month | 1.9 | 2.3 | 2.3 | 2.1 | 1.8 | 1.4 | 1.5 |
| | About once a month | .7 | .9 | 1.0 | .9 | 1.0 | .8 | .8 |
| | About once a week | 1.4 | 1.5 | 1.4 | 1.2 | 1.1 | .9 | 1.0 |
| | Daily | 2.6 | 2.7 | 2.6 | 2.2 | 1.8 | 1.7 | 1.8 |
| Felt bad about the amount bet | No Answer | 9.4 | 8.7 | 8.1 | 7.7 | 7.3 | 6.7 | 5.8 |
| | Yes | 1.7 | 1.8 | 1.6 | 1.4 | 1.2 | 1.2 | 1.3 |
| | No | 11.0 | 12.6 | 14.0 | 13.3 | 12.1 | 11.2 | 14.7 |
| | I do not bet for money | 77.9 | 76.9 | 76.3 | 77.7 | 79.4 | 80.9 | 78.1 |
| Would like to stop betting but could not | No Answer | 9.7 | 9.0 | 8.4 | 7.8 | 7.6 | 6.9 | 5.9 |
| | Yes | 1.4 | 1.3 | 1.3 | 1.1 | .9 | 1.0 | .9 |
| | No | 11.1 | 12.6 | 13.9 | 13.0 | 11.8 | 11.1 | 14.4 |
| | I do not bet for money | 77.7 | 77.1 | 76.5 | 78.0 | 79.7 | 81.0 | 78.7 |

**Perceived Risk of Selected Substance Use Behavior
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|---|---------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Smoke 1+ pack cigarettes per day | No Answer | 3.1 | 2.5 | 2.2 | 2.7 | 3.0 | 2.8 | 2.6 |
| | No risk | 6.1 | 4.7 | 4.1 | 3.7 | 3.3 | 3.3 | 3.2 |
| | Slight risk | 10.0 | 7.7 | 8.4 | 8.0 | 7.1 | 6.9 | 6.1 |
| | Moderate risk | 21.8 | 21.8 | 23.1 | 22.4 | 20.8 | 20.3 | 20.1 |
| | Great risk | 59.1 | 63.3 | 62.1 | 63.3 | 65.9 | 66.7 | 67.9 |
| Smoke marijuana occasionally | No Answer | 3.4 | 2.9 | 2.5 | 2.9 | 3.1 | 3.0 | 2.7 |
| | No risk | 6.1 | 6.3 | 9.0 | 12.5 | 14.2 | 16.7 | 17.2 |
| | Slight risk | 9.5 | 11.5 | 16.3 | 20.7 | 23.9 | 24.8 | 26.4 |
| | Moderate risk | 31.4 | 31.6 | 32.0 | 30.6 | 29.7 | 28.6 | 28.2 |
| | Great risk | 49.5 | 47.7 | 40.3 | 33.3 | 29.1 | 26.8 | 25.4 |
| Smoke marijuana regularly | No Answer | 4.2 | 3.6 | 3.2 | 3.6 | 3.7 | 3.6 | 3.4 |
| | No risk | 5.9 | 5.0 | 6.0 | 7.1 | 7.6 | 8.4 | 8.9 |
| | Slight risk | 4.2 | 4.4 | 7.1 | 9.9 | 11.7 | 12.7 | 13.4 |
| | Moderate risk | 11.7 | 11.4 | 14.6 | 18.3 | 20.6 | 22.6 | 23.6 |
| | Great risk | 74.1 | 75.5 | 69.0 | 61.1 | 56.3 | 52.7 | 50.6 |
| Occasionally consume 1-2 alcoholic drinks | No Answer | 4.2 | 3.6 | 3.2 | 3.5 | 3.7 | 3.4 | 3.1 |
| | No risk | 18.4 | 20.0 | 23.7 | 28.6 | 27.4 | 30.4 | 32.1 |
| | Slight risk | 38.5 | 38.6 | 39.8 | 38.1 | 39.9 | 38.5 | 39.7 |
| | Moderate risk | 20.9 | 20.0 | 18.7 | 17.1 | 16.9 | 15.4 | 14.0 |
| | Great risk | 18.0 | 17.7 | 14.6 | 12.7 | 12.1 | 12.3 | 11.1 |
| Binge drink weekly | No Answer | 3.8 | 3.0 | 2.7 | 3.1 | 3.3 | 3.1 | 2.8 |
| | No risk | 7.0 | 5.6 | 5.7 | 5.5 | 4.5 | 4.7 | 4.3 |
| | Slight risk | 12.4 | 12.7 | 13.3 | 14.6 | 13.8 | 13.8 | 14.5 |
| | Moderate risk | 30.1 | 29.3 | 31.4 | 32.8 | 31.3 | 32.0 | 33.5 |
| | Great risk | 46.6 | 49.4 | 46.9 | 44.0 | 47.1 | 46.4 | 44.9 |

**Perceived Peer Approval of Using Selected Substances
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|---------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Smoke 1+ pack cigarettes per day | No Answer | 4.1 | 3.5 | 2.8 | 3.4 | 3.2 | 3.1 | 3.0 |
| | Strongly approve | 1.7 | 1.4 | 1.5 | 1.6 | 1.6 | 2.2 | 1.8 |
| | Approve | .9 | 1.7 | 3.1 | 4.2 | 4.7 | 5.7 | 6.0 |
| | Do not know | 8.6 | 9.8 | 12.7 | 13.8 | 14.4 | 13.7 | 15.2 |
| | Disapprove | 8.5 | 10.0 | 13.2 | 16.3 | 16.5 | 17.2 | 19.4 |
| | Strongly disapprove | 76.2 | 73.6 | 66.8 | 60.7 | 59.5 | 58.0 | 54.6 |
| Smoke marijuana occasionally | No Answer | 4.3 | 3.5 | 2.9 | 3.5 | 3.6 | 3.4 | 3.1 |
| | Strongly approve | 1.8 | 1.9 | 2.8 | 3.7 | 4.5 | 4.8 | 4.6 |
| | Approve | 1.1 | 3.5 | 7.5 | 11.3 | 15.0 | 16.6 | 17.8 |
| | Do not know | 7.2 | 8.6 | 12.0 | 14.3 | 15.5 | 15.6 | 16.6 |
| | Disapprove | 6.3 | 7.8 | 10.6 | 13.5 | 13.6 | 14.0 | 15.5 |
| | Strongly disapprove | 79.3 | 74.8 | 64.2 | 53.6 | 47.9 | 45.7 | 42.3 |
| Smoke marijuana regularly | No Answer | 4.6 | 3.7 | 3.2 | 4.0 | 3.8 | 3.6 | 3.5 |
| | Strongly approve | 1.9 | 1.8 | 2.9 | 3.3 | 4.0 | 4.4 | 3.9 |
| | Approve | .7 | 2.1 | 4.3 | 6.6 | 8.6 | 9.0 | 9.3 |
| | Do not know | 6.9 | 7.9 | 11.1 | 12.7 | 13.8 | 13.7 | 14.5 |
| | Disapprove | 4.5 | 5.7 | 8.2 | 11.3 | 12.7 | 13.3 | 14.9 |
| | Strongly disapprove | 81.4 | 78.7 | 70.3 | 62.1 | 57.0 | 56.0 | 53.8 |
| Occasionally consume 1-2 alcoholic drinks | No Answer | 4.5 | 3.7 | 3.3 | 3.7 | 3.7 | 3.5 | 3.3 |
| | Strongly approve | 2.0 | 2.4 | 3.5 | 4.5 | 5.2 | 6.3 | 7.4 |
| | Approve | 3.4 | 7.2 | 14.0 | 20.9 | 25.0 | 28.5 | 33.0 |
| | Do not know | 12.9 | 15.0 | 18.8 | 21.6 | 22.3 | 21.6 | 21.4 |
| | Disapprove | 12.2 | 12.8 | 14.2 | 14.8 | 14.5 | 13.3 | 12.2 |
| | Strongly disapprove | 65.0 | 58.8 | 46.2 | 34.5 | 29.1 | 26.8 | 22.6 |
| Binge drink weekly | No Answer | 4.4 | 3.7 | 3.1 | 3.6 | 3.6 | 3.5 | 3.2 |
| | Strongly approve | 1.9 | 1.8 | 2.4 | 2.7 | 3.0 | 3.3 | 3.2 |
| | Approve | .9 | 2.3 | 4.4 | 6.8 | 8.1 | 9.0 | 10.7 |
| | Do not know | 8.4 | 10.1 | 13.8 | 16.8 | 17.8 | 17.4 | 19.4 |
| | Disapprove | 6.9 | 8.3 | 12.1 | 15.8 | 16.8 | 18.4 | 19.8 |
| | Strongly disapprove | 77.5 | 73.7 | 64.2 | 54.3 | 50.8 | 48.4 | 43.6 |

**Parental Attitudes toward Drug Use and Antisocial Behavior
by Indiana Female 6th - 12th Graders, 2011**

(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Parents feel about alcohol | No Answer | 3.4 | 3.1 | 2.6 | 3.1 | 3.5 | 3.1 | 3.2 |
| | Very wrong | 86.9 | 83.8 | 77.7 | 70.9 | 67.0 | 63.3 | 55.3 |
| | Wrong | 5.7 | 7.4 | 10.7 | 13.6 | 15.0 | 15.8 | 17.5 |
| | A little bit wrong | 2.8 | 4.2 | 6.8 | 9.5 | 11.0 | 13.4 | 17.8 |
| | Not at all wrong | 1.2 | 1.5 | 2.2 | 2.9 | 3.6 | 4.4 | 6.2 |
| Parents feel about cigarettes | No Answer | 3.3 | 3.1 | 2.6 | 3.2 | 3.4 | 3.2 | 3.1 |
| | Very wrong | 90.1 | 87.9 | 85.0 | 80.7 | 78.6 | 74.9 | 66.6 |
| | Wrong | 4.8 | 6.1 | 8.1 | 10.0 | 10.7 | 12.3 | 16.2 |
| | A little bit wrong | 1.0 | 1.6 | 2.7 | 4.1 | 4.5 | 6.0 | 8.8 |
| | Not at all wrong | .9 | 1.2 | 1.7 | 1.9 | 2.8 | 3.6 | 5.3 |
| Parents feel about marijuana | No Answer | 3.5 | 3.1 | 2.7 | 3.3 | 3.5 | 3.3 | 3.2 |
| | Very wrong | 93.5 | 91.6 | 88.4 | 83.5 | 80.5 | 79.3 | 76.4 |
| | Wrong | 1.7 | 3.1 | 5.0 | 7.2 | 8.5 | 9.1 | 10.7 |
| | A little bit wrong | .5 | 1.1 | 2.1 | 3.6 | 4.7 | 5.3 | 6.1 |
| | Not at all wrong | .8 | 1.1 | 1.8 | 2.4 | 2.8 | 3.0 | 3.5 |
| Parents feel about stealing | No Answer | 3.6 | 3.2 | 2.8 | 3.4 | 3.6 | 3.4 | 3.2 |
| | Very wrong | 87.1 | 85.7 | 83.9 | 81.9 | 82.9 | 84.0 | 83.5 |
| | Wrong | 7.1 | 8.0 | 9.3 | 11.0 | 10.1 | 9.3 | 9.9 |
| | A little bit wrong | 1.4 | 2.0 | 2.5 | 2.5 | 2.2 | 1.9 | 2.0 |
| | Not at all wrong | .8 | 1.1 | 1.4 | 1.3 | 1.2 | 1.4 | 1.4 |
| Parents feel about graffiti | No Answer | 3.7 | 3.3 | 2.8 | 3.4 | 3.6 | 3.3 | 3.2 |
| | Very wrong | 86.9 | 85.1 | 82.4 | 79.6 | 80.4 | 81.1 | 79.9 |
| | Wrong | 6.5 | 7.8 | 9.8 | 11.6 | 11.0 | 10.7 | 11.9 |
| | A little bit wrong | 1.7 | 2.3 | 3.2 | 3.6 | 3.4 | 3.2 | 3.4 |
| | Not at all wrong | 1.2 | 1.5 | 1.7 | 1.8 | 1.6 | 1.6 | 1.7 |
| Parents feel about fighting | No Answer | 3.8 | 3.3 | 3.0 | 3.5 | 3.7 | 3.4 | 3.2 |
| | Very wrong | 71.2 | 66.0 | 60.1 | 59.3 | 59.6 | 62.5 | 61.6 |
| | Wrong | 16.4 | 18.4 | 20.8 | 21.5 | 21.8 | 21.8 | 23.2 |
| | A little bit wrong | 6.4 | 9.4 | 12.3 | 12.6 | 11.8 | 9.7 | 9.7 |
| | Not at all wrong | 2.3 | 2.9 | 3.8 | 3.2 | 3.1 | 2.6 | 2.4 |

**Family Management and Conflict
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Clear rules | No Answer | 2.3 | 2.8 | 2.3 | 3.7 | 3.7 | 3.6 | 3.6 |
| | YES! | 65.3 | 58.6 | 53.9 | 51.1 | 52.4 | 54.2 | 54.2 |
| | yes | 28.0 | 32.0 | 35.3 | 36.6 | 36.1 | 34.4 | 33.9 |
| | no | 3.1 | 5.0 | 6.5 | 6.6 | 6.1 | 5.7 | 6.2 |
| | NO! | 1.3 | 1.7 | 2.0 | 1.9 | 1.8 | 2.1 | 2.1 |
| Ask about homework | No Answer | 2.2 | 2.9 | 2.6 | 3.8 | 3.9 | 3.9 | 3.9 |
| | YES! | 70.5 | 60.3 | 50.6 | 44.4 | 38.9 | 35.0 | 29.0 |
| | yes | 21.3 | 25.4 | 30.7 | 32.0 | 33.8 | 33.1 | 33.2 |
| | no | 4.4 | 8.5 | 11.7 | 14.6 | 16.9 | 19.8 | 23.3 |
| | NO! | 1.6 | 2.9 | 4.4 | 5.1 | 6.5 | 8.2 | 10.6 |
| Parents know whereabouts | No Answer | 2.3 | 3.1 | 2.6 | 4.0 | 4.0 | 4.1 | 4.0 |
| | YES! | 81.7 | 74.5 | 67.7 | 60.7 | 58.6 | 55.9 | 50.8 |
| | yes | 11.6 | 16.0 | 21.4 | 26.3 | 28.6 | 29.8 | 31.9 |
| | no | 2.9 | 4.4 | 6.2 | 6.9 | 6.8 | 7.6 | 9.8 |
| | NO! | 1.5 | 2.0 | 2.1 | 2.1 | 2.1 | 2.7 | 3.5 |
| Clear rules about alcohol use | No Answer | 3.4 | 3.7 | 3.3 | 4.5 | 4.3 | 4.4 | 4.4 |
| | YES! | 82.1 | 76.2 | 67.6 | 60.7 | 58.0 | 56.4 | 52.1 |
| | yes | 9.4 | 12.6 | 18.1 | 22.5 | 24.2 | 25.3 | 27.1 |
| | no | 3.3 | 5.6 | 8.5 | 9.9 | 10.9 | 11.0 | 13.2 |
| | NO! | 1.8 | 1.9 | 2.5 | 2.5 | 2.6 | 2.8 | 3.2 |
| Parents ask before most family decisions made | No Answer | 5.5 | 5.0 | 4.2 | 5.2 | 5.1 | 5.1 | 5.0 |
| | YES! | 38.2 | 31.9 | 26.5 | 23.6 | 22.3 | 21.7 | 21.2 |
| | yes | 33.2 | 34.5 | 35.0 | 35.2 | 36.1 | 36.5 | 37.3 |
| | no | 13.4 | 17.0 | 20.9 | 22.7 | 23.3 | 23.6 | 24.1 |
| | NO! | 9.7 | 11.6 | 13.4 | 13.3 | 13.2 | 13.1 | 12.5 |
| Ask parents for help if problem | No Answer | 3.4 | 3.9 | 3.3 | 4.9 | 4.7 | 4.7 | 4.5 |
| | YES! | 65.3 | 54.2 | 45.6 | 41.0 | 40.2 | 41.3 | 43.3 |
| | yes | 19.7 | 23.5 | 27.2 | 28.8 | 31.0 | 31.2 | 32.6 |
| | no | 6.3 | 9.8 | 13.1 | 14.3 | 14.2 | 13.4 | 12.2 |
| | NO! | 5.4 | 8.6 | 10.8 | 10.9 | 9.9 | 9.5 | 7.4 |
| Parents give me chances for fun with them | No Answer | 3.8 | 4.3 | 3.7 | 5.2 | 5.1 | 4.9 | 4.8 |
| | YES! | 54.9 | 46.9 | 39.5 | 34.2 | 32.7 | 31.9 | 31.8 |
| | yes | 26.8 | 29.0 | 31.6 | 32.9 | 33.7 | 34.3 | 34.6 |
| | no | 10.1 | 13.5 | 17.4 | 19.5 | 20.4 | 20.0 | 20.5 |
| | NO! | 4.4 | 6.2 | 7.9 | 8.1 | 8.2 | 8.8 | 8.2 |

**Family Management and Conflict
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Parents know when home late | No Answer | 4.4 | 5.1 | 4.1 | 5.6 | 5.3 | 5.4 | 5.3 |
| | YES! | 72.5 | 65.3 | 61.0 | 55.6 | 53.7 | 51.2 | 46.6 |
| | yes | 16.7 | 20.3 | 23.7 | 27.0 | 28.6 | 28.7 | 29.7 |
| | no | 4.5 | 6.8 | 8.4 | 9.0 | 9.6 | 10.9 | 13.3 |
| | NO! | 1.9 | 2.6 | 2.8 | 2.8 | 2.8 | 3.8 | 5.2 |
| Parents catch you drinking | No Answer | 4.3 | 4.7 | 4.1 | 5.6 | 5.6 | 5.3 | 5.1 |
| | YES! | 75.1 | 61.7 | 47.2 | 34.4 | 29.4 | 26.7 | 22.7 |
| | yes | 11.1 | 15.6 | 19.5 | 21.4 | 21.5 | 20.2 | 19.3 |
| | no | 6.0 | 12.1 | 19.9 | 27.4 | 31.8 | 34.3 | 38.3 |
| | NO! | 3.6 | 6.0 | 9.3 | 11.1 | 11.8 | 13.5 | 14.7 |
| Parents catch you with gun | No Answer | 4.4 | 4.8 | 4.0 | 5.5 | 5.4 | 5.6 | 5.1 |
| | YES! | 84.0 | 77.5 | 71.3 | 64.0 | 60.0 | 58.5 | 52.4 |
| | yes | 7.4 | 11.1 | 14.9 | 18.6 | 20.1 | 20.6 | 23.3 |
| | no | 2.2 | 4.4 | 6.8 | 8.4 | 10.4 | 10.6 | 13.9 |
| | NO! | 2.0 | 2.2 | 2.9 | 3.5 | 4.1 | 4.7 | 5.3 |
| Parents catch you skip school | No Answer | 3.6 | 4.3 | 3.5 | 5.1 | 4.9 | 4.9 | 4.7 |
| | YES! | 82.3 | 75.9 | 69.5 | 62.5 | 58.1 | 55.9 | 49.4 |
| | yes | 9.6 | 12.9 | 16.8 | 21.0 | 23.4 | 24.1 | 26.2 |
| | no | 2.9 | 4.3 | 7.1 | 7.9 | 9.7 | 10.4 | 14.0 |
| | NO! | 1.7 | 2.5 | 3.0 | 3.5 | 3.9 | 4.6 | 5.8 |
| Family argue repetitively | No Answer | 4.6 | 4.3 | 3.5 | 4.6 | 4.6 | 4.6 | 4.4 |
| | YES! | 17.6 | 21.6 | 25.3 | 26.4 | 25.5 | 24.5 | 22.5 |
| | yes | 20.3 | 22.8 | 24.0 | 26.6 | 27.3 | 26.1 | 25.7 |
| | no | 31.4 | 31.9 | 32.5 | 30.9 | 31.1 | 31.9 | 33.7 |
| | NO! | 26.1 | 19.3 | 14.6 | 11.5 | 11.4 | 13.0 | 13.6 |
| Family have serious arguments | No Answer | 4.5 | 4.6 | 3.7 | 4.9 | 5.0 | 4.8 | 4.9 |
| | YES! | 13.2 | 15.8 | 18.2 | 19.7 | 18.8 | 18.1 | 16.3 |
| | yes | 14.5 | 15.3 | 18.2 | 18.7 | 19.5 | 18.4 | 18.0 |
| | no | 28.1 | 31.1 | 33.2 | 33.7 | 34.5 | 34.6 | 36.2 |
| | NO! | 39.7 | 33.3 | 26.7 | 23.0 | 22.3 | 24.2 | 24.7 |
| Family insult each other | No Answer | 5.7 | 5.1 | 4.6 | 5.6 | 6.0 | 5.7 | 5.7 |
| | YES! | 15.0 | 17.2 | 20.2 | 20.6 | 19.6 | 17.9 | 15.9 |
| | yes | 16.7 | 18.7 | 20.4 | 20.9 | 20.7 | 19.4 | 19.3 |
| | no | 25.0 | 27.5 | 28.0 | 29.2 | 29.8 | 30.4 | 30.8 |
| | NO! | 37.5 | 31.5 | 26.7 | 23.6 | 23.8 | 26.5 | 28.3 |

Interaction with Antisocial Peers
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Number of best friends suspended | No Answer | 3.4 | 4.1 | 3.6 | 5.4 | 5.2 | 4.9 | 4.7 |
| | None of my friends | 77.9 | 71.4 | 67.5 | 67.3 | 69.7 | 73.9 | 78.7 |
| | 1 of my friends | 11.5 | 14.1 | 16.0 | 15.5 | 14.4 | 13.0 | 10.2 |
| | 2 of my friends | 3.9 | 5.3 | 6.8 | 6.4 | 6.1 | 5.1 | 3.6 |
| | 3 of my friends | 1.2 | 1.9 | 2.5 | 2.2 | 2.0 | 1.4 | .9 |
| | 4 of my friends | 2.1 | 3.1 | 3.5 | 3.2 | 2.5 | 1.7 | 1.7 |
| Number of best friends carry guns | No Answer | 3.3 | 4.2 | 3.5 | 5.3 | 5.3 | 4.9 | 4.8 |
| | None of my friends | 94.4 | 92.2 | 92.1 | 90.1 | 89.2 | 89.6 | 89.6 |
| | 1 of my friends | 1.6 | 2.5 | 2.7 | 2.7 | 3.4 | 3.6 | 3.7 |
| | 2 of my friends | .3 | .6 | .8 | .9 | 1.0 | 1.0 | .9 |
| | 3 of my friends | .1 | .2 | .3 | .3 | .4 | .3 | .4 |
| | 4 of my friends | .2 | .4 | .4 | .7 | .7 | .6 | .6 |
| Number of best friends sold drugs | No Answer | 3.4 | 4.4 | 3.7 | 5.6 | 5.6 | 5.1 | 5.0 |
| | None of my friends | 93.7 | 88.7 | 83.7 | 76.9 | 75.2 | 74.3 | 75.6 |
| | 1 of my friends | 2.0 | 3.9 | 6.8 | 8.9 | 9.5 | 10.3 | 9.8 |
| | 2 of my friends | .5 | 1.4 | 2.9 | 3.9 | 4.7 | 5.1 | 5.1 |
| | 3 of my friends | .2 | .7 | 1.3 | 1.7 | 2.0 | 2.2 | 1.9 |
| | 4 of my friends | .3 | 1.0 | 1.6 | 3.0 | 3.0 | 3.0 | 2.5 |
| Number of best friends stolen vehicle | No Answer | 3.4 | 4.5 | 3.7 | 5.5 | 5.5 | 5.0 | 4.9 |
| | None of my friends | 94.7 | 92.3 | 90.9 | 87.6 | 88.3 | 90.3 | 91.4 |
| | 1 of my friends | 1.2 | 1.9 | 3.3 | 4.0 | 3.7 | 2.9 | 1.9 |
| | 2 of my friends | .3 | .5 | 1.1 | 1.4 | 1.3 | .8 | .8 |
| | 3 of my friends | .1 | .3 | .5 | .5 | .4 | .3 | .5 |
| | 4 of my friends | .2 | .4 | .6 | .9 | .8 | .6 | .6 |
| Number of best friends arrested | No Answer | 3.5 | 4.4 | 3.7 | 5.6 | 5.5 | 5.0 | 5.0 |
| | None of my friends | 89.5 | 84.6 | 80.4 | 77.3 | 76.4 | 77.5 | 78.4 |
| | 1 of my friends | 4.8 | 6.7 | 9.4 | 9.8 | 10.5 | 10.7 | 10.1 |
| | 2 of my friends | 1.1 | 2.2 | 3.1 | 3.6 | 4.0 | 3.6 | 3.4 |
| | 3 of my friends | .5 | .9 | 1.5 | 1.4 | 1.4 | 1.5 | 1.2 |
| | 4 of my friends | .5 | 1.2 | 1.9 | 2.2 | 2.1 | 1.8 | 1.8 |
| Number of best friends dropped out of school | No Answer | 3.6 | 4.5 | 3.8 | 5.6 | 5.5 | 5.0 | 5.0 |
| | None of my friends | 92.3 | 89.8 | 88.6 | 83.5 | 81.3 | 80.5 | 82.0 |
| | 1 of my friends | 3.1 | 3.7 | 4.9 | 7.1 | 8.3 | 9.3 | 8.1 |
| | 2 of my friends | .6 | 1.0 | 1.3 | 1.9 | 2.3 | 2.9 | 2.3 |
| | 3 of my friends | .2 | .4 | .6 | .8 | 1.0 | 1.1 | 1.2 |
| | 4 of my friends | .3 | .6 | .8 | 1.2 | 1.4 | 1.1 | 1.4 |
| Number of best friends been members of gangs | No Answer | 4.8 | 5.5 | 4.6 | 6.4 | 6.5 | 6.0 | 5.8 |
| | None of my friends | 88.1 | 84.7 | 82.9 | 82.8 | 83.7 | 85.9 | 87.6 |
| | 1 of my friends | 3.7 | 4.5 | 5.9 | 5.0 | 4.2 | 4.0 | 2.7 |
| | 2 of my friends | 1.2 | 1.9 | 2.5 | 2.0 | 2.2 | 1.7 | 1.6 |
| | 3 of my friends | .7 | 1.0 | 1.4 | 1.2 | 1.0 | .8 | .7 |
| | 4 of my friends | 1.5 | 2.5 | 2.7 | 2.6 | 2.5 | 1.7 | 1.6 |

Interaction with Prosocial Peers
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Number of best friends participated in school activities | No Answer | 5.9 | 6.1 | 5.5 | 6.7 | 6.8 | 6.0 | 6.2 |
| | None of my friends | 16.1 | 17.1 | 17.3 | 17.1 | 17.3 | 17.4 | 17.6 |
| | 1 of my friends | 9.8 | 8.8 | 9.1 | 8.7 | 8.7 | 8.9 | 8.7 |
| | 2 of my friends | 15.1 | 14.6 | 14.4 | 14.4 | 14.9 | 14.6 | 13.6 |
| | 3 of my friends | 13.1 | 13.6 | 14.6 | 13.9 | 13.7 | 12.3 | 12.1 |
| | 4 of my friends | 39.9 | 39.8 | 39.1 | 39.2 | 38.7 | 40.8 | 41.9 |
| Number of best friends commitment to stay drug-free | No Answer | 7.4 | 7.1 | 6.3 | 7.4 | 7.3 | 6.7 | 6.7 |
| | None of my friends | 15.3 | 18.6 | 23.3 | 26.2 | 27.4 | 30.4 | 32.0 |
| | 1 of my friends | 3.7 | 5.3 | 8.5 | 9.9 | 11.8 | 12.3 | 11.8 |
| | 2 of my friends | 4.0 | 5.5 | 8.0 | 10.2 | 10.6 | 10.6 | 10.8 |
| | 3 of my friends | 5.8 | 6.9 | 9.3 | 9.7 | 10.0 | 9.7 | 9.4 |
| | 4 of my friends | 63.6 | 56.6 | 44.5 | 36.4 | 32.8 | 30.4 | 29.4 |
| Number of best friends liked school | No Answer | 6.6 | 6.4 | 5.9 | 7.7 | 7.2 | 6.6 | 6.4 |
| | None of my friends | 19.1 | 23.2 | 27.1 | 29.4 | 31.1 | 31.9 | 31.0 |
| | 1 of my friends | 9.0 | 10.0 | 10.5 | 11.0 | 11.7 | 11.3 | 11.3 |
| | 2 of my friends | 13.8 | 14.0 | 14.8 | 14.8 | 15.8 | 15.4 | 15.8 |
| | 3 of my friends | 14.1 | 13.6 | 13.5 | 13.0 | 12.2 | 12.0 | 12.3 |
| | 4 of my friends | 37.5 | 32.8 | 28.2 | 24.1 | 22.2 | 22.9 | 23.3 |
| Number of best friends regularly attended religious activities | No Answer | 8.4 | 7.8 | 6.2 | 7.8 | 7.2 | 6.7 | 6.2 |
| | None of my friends | 19.1 | 19.6 | 22.6 | 23.2 | 24.7 | 25.9 | 27.4 |
| | 1 of my friends | 14.5 | 15.3 | 17.0 | 18.0 | 18.1 | 19.4 | 18.8 |
| | 2 of my friends | 18.2 | 18.9 | 19.5 | 18.8 | 19.1 | 18.6 | 19.5 |
| | 3 of my friends | 15.2 | 15.5 | 14.5 | 13.2 | 13.3 | 11.7 | 11.6 |
| | 4 of my friends | 24.6 | 23.0 | 20.2 | 19.0 | 17.7 | 17.7 | 16.4 |
| Number of best friends tried to do well in school | No Answer | 6.7 | 6.3 | 6.1 | 7.1 | 7.1 | 6.6 | 6.4 |
| | None of my friends | 5.6 | 6.8 | 7.2 | 7.4 | 7.7 | 7.7 | 7.4 |
| | 1 of my friends | 3.4 | 4.4 | 5.5 | 5.6 | 5.2 | 5.7 | 5.6 |
| | 2 of my friends | 5.0 | 6.4 | 8.1 | 8.9 | 10.0 | 9.3 | 9.3 |
| | 3 of my friends | 8.7 | 11.4 | 13.7 | 14.0 | 14.6 | 14.8 | 13.4 |
| | 4 of my friends | 70.6 | 64.8 | 59.4 | 57.0 | 55.3 | 55.9 | 57.8 |

Rewards for Antisocial Involvement
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|---------------------------------|--------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Cool if smoke cigarettes | No Answer | 4.9 | 5.7 | 4.4 | 6.9 | 6.5 | 6.1 | 5.8 |
| | No or very little chance | 79.1 | 71.7 | 66.6 | 61.5 | 62.2 | 64.0 | 64.9 |
| | Little chance | 6.9 | 9.7 | 12.4 | 13.8 | 14.3 | 13.9 | 14.4 |
| | Some chance | 4.5 | 7.2 | 9.8 | 11.3 | 11.1 | 10.2 | 10.2 |
| | Pretty good chance | 2.1 | 3.0 | 4.1 | 4.2 | 3.7 | 3.6 | 2.8 |
| | Very good chance | 2.6 | 2.7 | 2.7 | 2.3 | 2.3 | 2.2 | 1.9 |
| Cool if drink alcohol regularly | No Answer | 5.0 | 5.9 | 4.6 | 7.0 | 6.6 | 6.2 | 6.0 |
| | No or very little chance | 78.9 | 70.0 | 61.5 | 52.0 | 48.9 | 50.3 | 48.6 |
| | Little chance | 7.1 | 9.7 | 12.6 | 13.7 | 14.2 | 13.9 | 14.1 |
| | Some chance | 4.0 | 7.5 | 11.1 | 13.6 | 14.6 | 14.3 | 15.1 |
| | Pretty good chance | 2.1 | 3.8 | 6.2 | 9.1 | 10.5 | 10.2 | 11.1 |
| | Very good chance | 2.8 | 3.1 | 4.0 | 4.5 | 5.2 | 5.1 | 5.1 |
| Cool if smoke marijuana | No Answer | 5.0 | 5.9 | 4.6 | 6.9 | 6.6 | 6.3 | 6.0 |
| | No or very little chance | 83.1 | 74.9 | 66.4 | 57.3 | 54.7 | 56.6 | 56.4 |
| | Little chance | 4.0 | 6.5 | 9.0 | 10.6 | 11.8 | 11.3 | 12.8 |
| | Some chance | 2.5 | 4.8 | 8.2 | 11.6 | 12.0 | 12.1 | 12.9 |
| | Pretty good chance | 1.8 | 3.5 | 5.8 | 8.0 | 8.5 | 8.2 | 7.4 |
| | Very good chance | 3.5 | 4.3 | 6.0 | 5.7 | 6.5 | 5.5 | 4.5 |
| Cool if carry a gun | No Answer | 5.1 | 6.0 | 4.6 | 7.0 | 6.7 | 6.2 | 5.9 |
| | No or very little chance | 82.4 | 79.7 | 78.7 | 76.3 | 78.1 | 80.2 | 80.4 |
| | Little chance | 4.7 | 6.0 | 7.4 | 7.9 | 7.2 | 6.8 | 7.2 |
| | Some chance | 2.8 | 3.4 | 4.3 | 4.6 | 4.0 | 3.9 | 3.5 |
| | Pretty good chance | 1.7 | 1.9 | 2.0 | 1.8 | 1.8 | 1.1 | 1.3 |
| | Very good chance | 3.2 | 2.9 | 2.9 | 2.4 | 2.3 | 1.9 | 1.7 |

School Commitment
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|--|----------------------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Days of school skipped during last month | No Answer | 3.8 | 4.6 | 4.0 | 5.7 | 5.4 | 4.8 | 4.8 |
| | None | 89.2 | 85.5 | 82.6 | 80.1 | 78.7 | 77.9 | 71.8 |
| | 1 | 3.3 | 4.3 | 5.8 | 6.3 | 7.3 | 7.6 | 9.8 |
| | 2 | 1.6 | 2.4 | 3.0 | 3.1 | 3.4 | 4.0 | 5.8 |
| | 3 | .9 | 1.3 | 1.8 | 2.1 | 2.0 | 2.4 | 3.3 |
| | 4-5 | .8 | 1.0 | 1.6 | 1.6 | 1.9 | 1.9 | 2.6 |
| | 6-10 | .2 | .4 | .5 | .5 | .5 | .7 | 1.1 |
| | 11 or more | .3 | .5 | .6 | .7 | .7 | .5 | .8 |
| Feel schoolwork is meaningful | No Answer | 4.1 | 5.1 | 4.0 | 6.0 | 5.6 | 5.1 | 4.9 |
| | Almost always | 46.4 | 34.5 | 28.6 | 23.3 | 21.0 | 20.5 | 17.3 |
| | Often | 20.4 | 22.5 | 22.6 | 21.8 | 21.2 | 21.0 | 20.6 |
| | Sometimes | 19.2 | 22.5 | 26.8 | 29.1 | 31.2 | 31.3 | 33.0 |
| | Seldom | 4.7 | 9.1 | 11.7 | 13.1 | 14.6 | 15.9 | 17.6 |
| | Never | 5.1 | 6.2 | 6.3 | 6.6 | 6.5 | 6.3 | 6.6 |
| How interesting are courses | No Answer | 8.0 | 6.7 | 4.7 | 6.5 | 5.9 | 5.4 | 5.1 |
| | Very interesting and stimulating | 19.5 | 12.7 | 9.3 | 7.8 | 7.2 | 8.0 | 7.6 |
| | Quite interesting | 27.7 | 24.2 | 21.7 | 21.1 | 20.4 | 20.4 | 20.3 |
| | Fairly interesting | 27.5 | 32.2 | 36.6 | 37.6 | 37.8 | 38.0 | 37.9 |
| | Slightly dull | 10.5 | 13.6 | 16.1 | 16.1 | 17.4 | 16.9 | 16.7 |
| | Very dull | 6.7 | 10.5 | 11.6 | 10.9 | 11.3 | 11.2 | 12.5 |
| Importance of school to later life | No Answer | 5.2 | 6.0 | 4.6 | 6.7 | 6.0 | 5.6 | 5.3 |
| | Very important | 64.9 | 51.3 | 43.8 | 33.6 | 28.9 | 26.6 | 22.9 |
| | Quite important | 15.2 | 19.9 | 20.8 | 22.0 | 22.0 | 21.5 | 21.7 |
| | Fairly important | 8.0 | 12.4 | 16.5 | 20.2 | 21.9 | 23.4 | 24.3 |
| | Slightly important | 5.2 | 7.8 | 11.1 | 13.5 | 16.7 | 18.2 | 20.2 |
| | Not at all important | 1.6 | 2.6 | 3.1 | 4.0 | 4.5 | 4.8 | 5.6 |
| Grades better than other students | No Answer | 5.4 | 6.3 | 4.6 | 7.1 | 6.4 | 6.0 | 5.4 |
| | YES! | 30.1 | 28.2 | 27.6 | 25.9 | 25.3 | 26.5 | 29.7 |
| | yes | 45.2 | 44.1 | 44.5 | 43.8 | 46.5 | 47.1 | 46.2 |
| | no | 15.1 | 16.1 | 18.4 | 18.3 | 17.6 | 16.9 | 16.1 |
| | NO! | 4.1 | 5.2 | 4.9 | 5.0 | 4.2 | 3.5 | 2.6 |
| How often did you enjoy school | No Answer | 4.8 | 5.7 | 4.4 | 6.7 | 6.3 | 5.7 | 5.5 |
| | Never | 5.1 | 6.3 | 7.2 | 7.7 | 8.4 | 8.7 | 8.3 |
| | Seldom | 7.0 | 9.6 | 11.3 | 12.7 | 14.8 | 16.5 | 16.0 |
| | Sometimes | 21.7 | 24.1 | 27.6 | 29.5 | 31.7 | 32.8 | 33.1 |
| | Often | 27.3 | 29.2 | 28.7 | 28.3 | 26.7 | 25.7 | 26.7 |
| | A lot | 34.1 | 25.0 | 20.9 | 15.0 | 12.1 | 10.6 | 10.3 |

School Commitment
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|-------------------------------------|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| How often did you hate school | No Answer | 5.0 | 5.9 | 4.5 | 6.8 | 6.3 | 5.7 | 5.4 |
| | Never | 25.5 | 16.3 | 12.0 | 8.2 | 6.7 | 5.9 | 5.5 |
| | Seldom | 27.8 | 29.3 | 28.4 | 25.3 | 23.8 | 22.1 | 22.6 |
| | Sometimes | 23.7 | 25.5 | 28.7 | 31.0 | 31.6 | 33.1 | 34.0 |
| | Often | 9.0 | 11.5 | 13.8 | 15.7 | 17.9 | 18.7 | 18.7 |
| | A lot | 9.0 | 11.5 | 12.7 | 13.0 | 13.8 | 14.4 | 13.8 |
| How often did you do your best work | No Answer | 5.0 | 5.9 | 4.5 | 7.0 | 6.5 | 5.7 | 5.4 |
| | Never | 1.0 | 1.6 | 1.5 | 1.5 | 1.6 | 1.5 | 1.2 |
| | Seldom | 1.8 | 3.4 | 4.1 | 4.1 | 3.7 | 3.7 | 3.1 |
| | Sometimes | 6.0 | 9.2 | 12.7 | 14.1 | 15.5 | 15.2 | 15.8 |
| | Often | 15.6 | 20.3 | 25.1 | 26.7 | 28.4 | 29.5 | 32.6 |
| | A lot | 70.5 | 59.6 | 52.0 | 46.5 | 44.4 | 44.3 | 42.0 |

**Opportunities for Prosocial Involvement
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Students have chances to help decide activities and rules | No Answer | 7.6 | 7.8 | 6.1 | 9.2 | 8.5 | 7.8 | 7.2 |
| | YES! | 28.7 | 23.0 | 19.8 | 21.1 | 18.6 | 20.0 | 17.1 |
| | yes | 35.4 | 33.7 | 34.1 | 34.1 | 32.6 | 31.7 | 33.0 |
| | no | 19.7 | 24.5 | 28.3 | 26.3 | 28.8 | 28.5 | 29.8 |
| | NO! | 8.6 | 11.0 | 11.7 | 9.3 | 11.5 | 11.9 | 12.9 |
| Teachers ask to work on special class projects | No Answer | 7.8 | 8.0 | 6.4 | 9.6 | 8.9 | 8.1 | 7.3 |
| | YES! | 24.3 | 18.2 | 15.7 | 11.4 | 10.7 | 10.9 | 11.4 |
| | yes | 32.0 | 31.4 | 31.5 | 28.4 | 29.6 | 29.2 | 32.1 |
| | no | 25.2 | 30.6 | 34.5 | 38.9 | 38.3 | 39.3 | 37.8 |
| | NO! | 10.6 | 11.6 | 12.0 | 11.7 | 12.5 | 12.5 | 11.4 |
| Chances for kids to get involved in activities outside of class | No Answer | 7.8 | 8.4 | 6.5 | 9.9 | 9.1 | 8.0 | 7.7 |
| | YES! | 70.0 | 67.6 | 63.4 | 59.0 | 55.3 | 55.1 | 52.8 |
| | yes | 17.4 | 18.9 | 24.1 | 24.9 | 28.3 | 29.7 | 31.5 |
| | no | 3.3 | 3.6 | 4.1 | 4.4 | 5.3 | 5.0 | 5.6 |
| | NO! | 1.6 | 1.6 | 1.8 | 1.8 | 1.9 | 2.1 | 2.4 |
| Chances for kids to talk with teacher one-on-one | No Answer | 8.5 | 8.8 | 7.0 | 10.2 | 9.4 | 8.5 | 8.0 |
| | YES! | 51.7 | 43.8 | 40.4 | 33.9 | 30.9 | 32.1 | 32.4 |
| | yes | 27.7 | 30.7 | 33.5 | 36.7 | 38.7 | 38.9 | 40.9 |
| | no | 8.8 | 12.3 | 14.5 | 14.8 | 16.5 | 15.8 | 14.3 |
| | NO! | 3.3 | 4.4 | 4.6 | 4.3 | 4.4 | 4.7 | 4.3 |
| Chances to be a part of class discussion and activities | No Answer | 9.7 | 9.6 | 8.0 | 11.0 | 10.1 | 8.9 | 8.4 |
| | YES! | 48.7 | 42.5 | 36.9 | 29.9 | 26.3 | 27.4 | 28.2 |
| | yes | 29.8 | 34.2 | 39.8 | 43.9 | 47.1 | 48.2 | 49.2 |
| | no | 8.6 | 10.0 | 11.5 | 11.8 | 12.7 | 12.0 | 10.8 |
| | NO! | 3.2 | 3.8 | 3.8 | 3.3 | 3.8 | 3.5 | 3.4 |

Rewards for Prosocial Involvement
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Teachers notice when doing a good job and let me know | No Answer | 8.1 | 8.5 | 6.8 | 10.0 | 9.3 | 8.3 | 7.8 |
| | YES! | 43.9 | 33.9 | 29.0 | 21.5 | 18.4 | 18.7 | 18.7 |
| | yes | 31.3 | 34.4 | 37.5 | 38.3 | 39.7 | 40.1 | 41.7 |
| | no | 11.5 | 16.2 | 19.4 | 22.3 | 23.9 | 24.7 | 23.7 |
| | NO! | 5.1 | 7.0 | 7.3 | 7.9 | 8.7 | 8.3 | 8.2 |
| Feel safe at school | No Answer | 8.0 | 8.6 | 6.8 | 9.9 | 9.1 | 8.3 | 7.7 |
| | YES! | 59.3 | 47.4 | 40.4 | 31.5 | 30.0 | 32.9 | 34.2 |
| | yes | 24.0 | 31.5 | 37.6 | 42.5 | 44.1 | 43.8 | 45.0 |
| | no | 5.6 | 8.2 | 10.4 | 11.2 | 11.6 | 10.7 | 9.3 |
| | NO! | 3.2 | 4.3 | 4.9 | 4.9 | 5.2 | 4.2 | 3.8 |
| School lets parents know when do good | No Answer | 8.8 | 9.1 | 7.2 | 10.3 | 9.5 | 8.6 | 8.0 |
| | YES! | 27.5 | 22.3 | 18.2 | 13.1 | 11.7 | 12.0 | 11.7 |
| | yes | 27.5 | 24.9 | 24.0 | 20.8 | 21.0 | 21.3 | 21.5 |
| | no | 24.2 | 28.6 | 32.3 | 36.5 | 37.6 | 37.2 | 37.0 |
| | NO! | 12.0 | 15.1 | 18.2 | 19.2 | 20.3 | 21.0 | 21.7 |
| Teachers praise when do well in school | No Answer | 10.4 | 10.0 | 8.1 | 11.2 | 10.1 | 9.2 | 8.5 |
| | YES! | 30.4 | 25.0 | 21.1 | 15.7 | 13.5 | 14.5 | 13.8 |
| | yes | 31.2 | 30.4 | 31.3 | 31.0 | 31.2 | 31.4 | 34.8 |
| | no | 19.0 | 24.0 | 27.6 | 29.8 | 32.4 | 32.6 | 30.8 |
| | NO! | 9.0 | 10.7 | 11.9 | 12.3 | 12.8 | 12.3 | 12.0 |

Participation in Afterschool Activities
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|---|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Participation in ROCK, LEAD, SADD, etc. | No Answer | 12.5 | 12.3 | 9.6 | 12.6 | 11.6 | 10.6 | 10.2 |
| | Yes | 20.1 | 14.1 | 10.6 | 9.4 | 9.2 | 10.4 | 10.4 |
| | No | 67.4 | 73.6 | 79.9 | 78.0 | 79.2 | 79.0 | 79.4 |

**Favorable Attitudes Toward Drug Use
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Wrong to take a gun to school | No Answer | 9.6 | 9.9 | 7.1 | 10.5 | 9.5 | 8.3 | 8.1 |
| | Very wrong | 85.6 | 82.9 | 84.0 | 80.6 | 82.6 | 85.0 | 85.7 |
| | Wrong | 3.5 | 5.3 | 6.4 | 6.5 | 5.7 | 4.9 | 4.4 |
| | A little bit wrong | .9 | 1.3 | 1.7 | 1.7 | 1.5 | 1.1 | 1.1 |
| | Not at all wrong | .5 | .6 | .8 | .7 | .8 | .7 | .8 |
| | | | | | | | | |
| Wrong to steal something more than \$5 | No Answer | 9.6 | 10.2 | 7.3 | 10.7 | 9.7 | 8.6 | 8.2 |
| | Very wrong | 70.5 | 64.1 | 60.0 | 54.2 | 57.2 | 61.1 | 63.2 |
| | Wrong | 15.2 | 17.9 | 22.3 | 24.1 | 23.1 | 22.0 | 21.7 |
| | A little bit wrong | 3.7 | 6.2 | 8.4 | 9.3 | 8.3 | 6.9 | 5.6 |
| | Not at all wrong | 1.0 | 1.6 | 1.9 | 1.7 | 1.7 | 1.4 | 1.3 |
| | | | | | | | | |
| Wrong to attack someone | No Answer | 9.8 | 10.2 | 7.5 | 11.0 | 9.8 | 8.7 | 8.3 |
| | Very wrong | 80.3 | 74.4 | 72.1 | 67.3 | 70.2 | 74.0 | 76.4 |
| | Wrong | 6.4 | 9.7 | 13.5 | 14.4 | 13.8 | 12.5 | 11.3 |
| | A little bit wrong | 2.3 | 4.0 | 5.2 | 5.4 | 4.6 | 3.8 | 2.9 |
| | Not at all wrong | 1.2 | 1.6 | 1.7 | 1.9 | 1.6 | 1.1 | 1.1 |
| | | | | | | | | |
| Wrong to pick a fight | No Answer | 9.9 | 10.5 | 7.6 | 11.0 | 9.9 | 8.9 | 8.4 |
| | Very wrong | 63.2 | 55.3 | 48.8 | 44.4 | 46.2 | 51.1 | 52.5 |
| | Wrong | 18.0 | 20.4 | 25.1 | 26.8 | 28.3 | 26.9 | 28.4 |
| | A little bit wrong | 6.8 | 10.3 | 14.5 | 14.1 | 12.9 | 11.0 | 9.0 |
| | Not at all wrong | 2.1 | 3.6 | 4.0 | 3.7 | 2.8 | 2.2 | 1.7 |
| | | | | | | | | |
| Wrong to skip school | No Answer | 10.1 | 10.5 | 7.7 | 11.2 | 10.0 | 9.1 | 8.4 |
| | Very wrong | 77.6 | 68.5 | 61.5 | 51.1 | 50.3 | 50.1 | 47.4 |
| | Wrong | 8.6 | 13.7 | 19.9 | 23.1 | 24.9 | 25.2 | 26.5 |
| | A little bit wrong | 2.6 | 5.1 | 8.5 | 11.5 | 12.0 | 12.8 | 14.7 |
| | Not at all wrong | 1.0 | 2.1 | 2.5 | 3.1 | 2.9 | 2.8 | 3.1 |
| | | | | | | | | |
| Wrong to drink alcohol regularly | No Answer | 10.1 | 10.5 | 7.7 | 11.2 | 10.1 | 9.1 | 8.5 |
| | Very wrong | 80.7 | 71.9 | 62.5 | 49.9 | 47.0 | 45.7 | 41.4 |
| | Wrong | 5.4 | 9.1 | 14.4 | 17.3 | 19.1 | 20.2 | 21.3 |
| | A little bit wrong | 2.7 | 6.0 | 11.1 | 15.5 | 17.6 | 18.7 | 21.4 |
| | Not at all wrong | 1.1 | 2.6 | 4.2 | 6.1 | 6.3 | 6.3 | 7.4 |
| | | | | | | | | |
| Wrong to smoke cigarettes | No Answer | 10.1 | 10.6 | 7.8 | 11.2 | 10.2 | 9.2 | 8.5 |
| | Very wrong | 81.6 | 74.7 | 67.6 | 57.1 | 54.2 | 51.4 | 43.6 |
| | Wrong | 5.2 | 7.9 | 12.5 | 15.1 | 17.2 | 17.1 | 18.7 |
| | A little bit wrong | 2.0 | 4.3 | 8.1 | 10.8 | 12.0 | 14.1 | 16.9 |
| | Not at all wrong | 1.1 | 2.6 | 4.1 | 5.7 | 6.4 | 8.2 | 12.3 |
| | | | | | | | | |

Favorable Attitudes Toward Drug Use
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|----------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Wrong to smoke marijuana | No Answer | 10.2 | 10.7 | 7.8 | 11.3 | 10.2 | 9.2 | 8.6 |
| | Very wrong | 85.3 | 78.6 | 71.1 | 59.5 | 55.3 | 53.3 | 50.3 |
| | Wrong | 2.4 | 5.0 | 9.3 | 12.3 | 14.0 | 14.9 | 16.4 |
| | A little bit wrong | 1.2 | 3.4 | 7.1 | 10.0 | 12.3 | 13.6 | 15.5 |
| | Not at all wrong | 1.0 | 2.3 | 4.6 | 6.9 | 8.2 | 8.9 | 9.3 |
| Wrong to use illegal drugs | No Answer | 10.4 | 10.7 | 7.9 | 11.4 | 10.2 | 9.2 | 8.5 |
| | Very wrong | 86.7 | 82.9 | 81.7 | 74.9 | 75.6 | 76.2 | 76.9 |
| | Wrong | 1.9 | 3.9 | 6.2 | 8.3 | 9.0 | 8.8 | 9.6 |
| | A little bit wrong | .6 | 1.5 | 2.6 | 3.4 | 3.2 | 3.8 | 3.0 |
| | Not at all wrong | .5 | 1.0 | 1.6 | 2.0 | 2.0 | 2.0 | 2.0 |

Rebelliousness
by Indiana Female 6th - 12th Graders, 2011
 (Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|----------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Like to see what can get away with | No Answer | 11.6 | 11.6 | 9.1 | 12.4 | 11.1 | 9.9 | 9.3 |
| | Very false | 63.1 | 53.0 | 44.9 | 39.4 | 38.7 | 40.4 | 41.8 |
| | Somewhat false | 13.8 | 17.7 | 20.9 | 21.1 | 22.4 | 21.8 | 22.9 |
| | Somewhat true | 8.6 | 13.4 | 18.8 | 20.4 | 21.4 | 21.7 | 20.8 |
| | Very true | 2.9 | 4.3 | 6.2 | 6.7 | 6.5 | 6.2 | 5.2 |
| Ignore rules | No Answer | 11.7 | 11.8 | 9.2 | 12.6 | 11.2 | 9.9 | 9.5 |
| | Very false | 66.2 | 58.1 | 51.6 | 45.6 | 46.7 | 48.2 | 48.2 |
| | Somewhat false | 13.2 | 16.6 | 19.9 | 21.2 | 21.9 | 22.6 | 23.7 |
| | Somewhat true | 6.4 | 9.6 | 14.1 | 15.5 | 15.3 | 15.1 | 15.0 |
| | Very true | 2.5 | 4.0 | 5.3 | 5.1 | 4.9 | 4.2 | 3.6 |
| Do opposite of what I am told | No Answer | 11.8 | 11.8 | 9.3 | 12.7 | 11.2 | 9.9 | 9.4 |
| | Very false | 65.9 | 60.0 | 55.8 | 52.4 | 55.2 | 58.7 | 61.2 |
| | Somewhat false | 12.8 | 15.5 | 18.6 | 18.8 | 19.2 | 18.4 | 17.8 |
| | Somewhat true | 7.0 | 9.2 | 11.9 | 11.9 | 10.7 | 9.7 | 8.9 |
| | Very true | 2.5 | 3.6 | 4.5 | 4.3 | 3.7 | 3.3 | 2.6 |

**Laws and Norms Favorable to Drug Use
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|------------------------------------|--------------------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Caught by police alcohol | No Answer | 13.2 | 12.7 | 9.7 | 13.6 | 12.0 | 11.0 | 10.4 |
| | YES! | 28.0 | 21.6 | 14.8 | 10.1 | 8.2 | 7.7 | 7.7 |
| | yes | 28.6 | 25.9 | 22.0 | 18.2 | 15.9 | 15.0 | 14.6 |
| | no | 20.8 | 28.6 | 38.0 | 41.3 | 45.6 | 46.7 | 47.9 |
| | NO! | 9.4 | 11.1 | 15.5 | 16.8 | 18.3 | 19.6 | 19.3 |
| Caught by police marijuana | No Answer | 13.2 | 12.8 | 9.9 | 13.9 | 12.2 | 11.1 | 10.4 |
| | YES! | 38.5 | 29.7 | 20.2 | 12.7 | 9.7 | 8.7 | 8.4 |
| | yes | 26.5 | 26.8 | 26.0 | 21.5 | 18.9 | 17.7 | 17.1 |
| | no | 14.1 | 21.2 | 30.3 | 35.9 | 41.5 | 43.4 | 45.2 |
| | NO! | 7.7 | 9.6 | 13.6 | 15.9 | 17.7 | 19.1 | 18.8 |
| Caught by police gun | No Answer | 13.2 | 12.9 | 9.9 | 13.8 | 12.2 | 11.1 | 10.5 |
| | YES! | 42.9 | 35.1 | 28.2 | 20.8 | 18.1 | 16.7 | 15.8 |
| | yes | 24.1 | 26.8 | 28.8 | 29.0 | 29.0 | 29.0 | 29.1 |
| | no | 12.9 | 17.5 | 23.4 | 26.1 | 29.7 | 31.7 | 32.8 |
| | NO! | 6.9 | 7.7 | 9.7 | 10.2 | 11.1 | 11.5 | 11.8 |
| Caught by police smoke cigarettes | No Answer | 13.6 | 13.1 | 10.4 | 14.2 | 12.4 | 11.1 | 10.9 |
| | YES! | 24.7 | 19.0 | 13.2 | 8.9 | 7.2 | 6.8 | 6.1 |
| | yes | 23.0 | 20.7 | 17.6 | 14.5 | 12.3 | 10.9 | 9.8 |
| | no | 24.6 | 30.5 | 36.9 | 38.1 | 42.2 | 42.2 | 43.1 |
| | NO! | 14.2 | 16.6 | 21.9 | 24.3 | 25.9 | 29.0 | 30.1 |
| Neighborhood feel about marijuana | No Answer | 12.4 | 12.2 | 9.4 | 13.6 | 11.8 | 10.7 | 10.1 |
| | Very wrong | 77.7 | 71.6 | 64.6 | 53.9 | 50.7 | 50.0 | 47.9 |
| | Wrong | 7.0 | 11.0 | 16.3 | 20.0 | 22.5 | 23.4 | 24.9 |
| | A little bit wrong | 2.1 | 3.9 | 7.3 | 9.5 | 11.0 | 12.2 | 13.0 |
| | Not at all wrong | .8 | 1.3 | 2.4 | 3.1 | 4.0 | 3.6 | 4.1 |
| Neighborhood feel about alcohol | No Answer | 12.4 | 12.2 | 9.6 | 13.7 | 12.0 | 10.8 | 10.4 |
| | Very wrong | 69.5 | 62.0 | 53.7 | 42.3 | 39.0 | 37.2 | 32.9 |
| | Wrong | 11.6 | 15.3 | 20.8 | 23.5 | 25.6 | 26.3 | 26.5 |
| | A little bit wrong | 4.8 | 7.6 | 11.8 | 15.0 | 17.4 | 19.1 | 22.4 |
| | Not at all wrong | 1.6 | 2.8 | 4.1 | 5.4 | 6.1 | 6.6 | 7.7 |
| Neighborhood feel about cigarettes | No Answer | 12.6 | 12.2 | 9.6 | 13.8 | 12.0 | 10.7 | 10.5 |
| | Very wrong | 69.7 | 63.1 | 55.9 | 45.6 | 42.1 | 39.3 | 31.8 |
| | Wrong | 11.3 | 14.8 | 19.9 | 22.9 | 24.8 | 25.1 | 25.5 |
| | A little bit wrong | 4.7 | 7.0 | 10.3 | 12.5 | 14.9 | 17.2 | 20.7 |
| | Not at all wrong | 1.8 | 2.9 | 4.3 | 5.2 | 6.2 | 7.7 | 11.4 |

**Community Rewards for Prosocial Involvement
by Indiana Female 6th - 12th Graders, 2011**
(Values expressed as percentages)

| | | Grade | | | | | | |
|--|-----------|-------|------|------|------|------|------|------|
| | | 6th | 7th | 8th | 9th | 10th | 11th | 12th |
| Neighbors notice good job and let me know | No Answer | 13.6 | 12.8 | 10.4 | 14.2 | 12.5 | 11.4 | 10.9 |
| | YES! | 21.8 | 16.4 | 12.5 | 9.6 | 8.0 | 8.0 | 7.5 |
| | yes | 21.3 | 19.9 | 17.0 | 15.0 | 14.4 | 14.2 | 14.0 |
| | no | 23.5 | 27.3 | 30.6 | 30.9 | 32.2 | 31.3 | 32.3 |
| | NO! | 19.9 | 23.6 | 29.6 | 30.3 | 32.9 | 35.1 | 35.3 |
| | | | | | | | | |
| People in neighborhood who are proud of me | No Answer | 13.6 | 12.9 | 10.5 | 14.3 | 12.6 | 11.5 | 11.0 |
| | YES! | 27.6 | 21.3 | 16.4 | 12.6 | 11.2 | 10.7 | 10.9 |
| | yes | 24.3 | 25.4 | 23.9 | 22.1 | 22.1 | 23.5 | 24.3 |
| | no | 18.3 | 20.2 | 23.9 | 24.5 | 25.4 | 24.0 | 24.1 |
| | NO! | 16.1 | 20.2 | 25.3 | 26.5 | 28.6 | 30.4 | 29.7 |
| | | | | | | | | |
| People in neighborhood who encourage to do my best | No Answer | 13.6 | 13.0 | 10.7 | 14.5 | 12.8 | 11.4 | 11.2 |
| | YES! | 33.5 | 26.4 | 19.7 | 14.9 | 12.9 | 12.3 | 11.8 |
| | yes | 23.2 | 24.1 | 23.9 | 22.9 | 22.7 | 23.4 | 23.7 |
| | no | 14.5 | 17.3 | 21.6 | 22.3 | 23.9 | 23.3 | 24.1 |
| | NO! | 15.2 | 19.1 | 24.1 | 25.5 | 27.7 | 29.6 | 29.2 |
| | | | | | | | | |



APPENDIX: INSTRUMENT

ALCOHOL, TOBACCO, AND OTHER DRUG USE SURVEY

© COPYRIGHT 2011, 2010, 2009, 2008, 2007, 2005, 2002, 2001, 1995, 1993, 1991 The Trustees of Indiana University. All rights reserved.
Questions #18 - 19 used with permission from the Minnesota Department of Education.

This survey is funded, in part, by a contract with the Indiana Family and Social Services Administration, Division of Mental Health and Addiction, HHS/SAMHSA, CSAP, SAPT Block Grant, and Indiana Gambler's Assistance Fund.

PRIVACY STATEMENT

This survey is completely confidential. Complete the survey in private and deposit your completed survey in the privacy envelope, as instructed by your teacher or supervisor. Do not place your name or any identifying marks on the survey form or envelope. There is no way for anyone to identify your individual responses. The survey envelopes will not be opened until they are received at the statewide processing center.

YOUR PARTICIPATION IS VOLUNTARY

Your participation in this survey is voluntary. If you do not wish to answer a question, just leave it blank. If you do not wish to participate in this survey, you may:

- Return the entire survey form blank.
- Answer questions with random answers and then answer "not truthfully at all" to the last question. Your form will be discarded and no one will know that you did not participate.
- Inform your teacher or supervisor that you choose not to participate in the survey.

You will not be penalized in any way for deciding not to participate in this survey. We do value your participation; however, accurate results are dependent upon getting as many students as possible to volunteer to participate in this confidential survey. The results are used by many state and federal agencies to evaluate the effectiveness of drug programs in your community. Your help is important to this effort.

Thank you for your assistance.

[illegible]

Please enter the unique ten-digit school code provided by your teacher or supervisor.


PLEASE DO NOT WRITE IN THIS AREA

serial #

MARKING INSTRUCTIONS

2

- Use a No. 2 pencil only.
- Do not use ink, ballpoint, or felt tip pens.
- Make solid marks that fill the response completely.
- Erase cleanly any marks you wish to change.
- Make no stray marks on this form.



CORRECT:

INCORRECT:



PERSONAL AND FAMILY INFORMATION

1. SEX:

- ☐ Male
☐ Female

2. ETHNICITY:

- ☐ Hispanic
☐ Non-Hispanic

3. RACE:

- ☐ White
☐ Black or African-American
☐ Asian
☐ Race not known or other
- ☐ Native Hawaiian/Pacific Islander
☐ American Indian/Alaskan Native
☐ More than one race

4. GRADE:

- ☐ 6th ☐ 7th ☐ 8th ☐ 9th
☐ 10th ☐ 11th ☐ 12th

5. AGE:

- ☐ 10 years old or less ☐ 11 years old ☐ 12 years old
☐ 13 years old ☐ 14 years old ☐ 15 years old
☐ 16 years old ☐ 17 years old ☐ 18 years or older

6. What grades do you generally get in school?

- ☐ Mostly A's
☐ Mostly B's
☐ Mostly C's
☐ Mostly D's
☐ Mostly F's

7. Do you live with...

- ☐ Both parents
☐ Mother only
☐ Father only
☐ Mother & stepfather
☐ Father & stepmother
☐ Other

8. Do you or your parents have a job?

| | YES, FULL TIME | YES, PART TIME | NO |
|--------|-----------------------|-----------------------|-----------------------|
| You | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Mother | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Father | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

LIFETIME USE:

9. Have you ever used.....?

| | NEVER | 1-5 TIMES | 6-19 TIMES | 20-39 TIMES | 40 TIMES OR MORE |
|---|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Cigarettes | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smokeless tobacco (chew, snuff, etc.) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Cigars | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Pipe (tobacco used in pipe, water-pipe, hookah) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Alcohol (beer, wine, liquor, wine coolers) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Marijuana (pot, hash, weed) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Cocaine | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Crack | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Inhalants | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Amphetamines (uppers) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Methamphetamines (meth, crystal, ice, crank) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Heroin | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Steroids (non-prescribed) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Ecstasy (MDMA, X, XTC) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Tranquilizers (downers) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Hallucinogens (LSD, PCP) | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Prescription painkillers (Vicodin, OxyContin, Percocet) to get high | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Prescription drugs (such as Ritalin, Adderall, Xanax) to get high | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Over-the-counter drugs to get high | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

10. How many times in the last month (30 days) have you used.....

AGE AT FIRST USE:

NEVER 10 OR 11 12 13 14 15 16 17 OR
USED YOUNGER OLDER

12. How easy would it be for you to get...

PLEASE DO NOT WRITE IN THIS AREA

serial #

ALCOHOL USE

4

13. How many times in the last two weeks have you had 5 or more for males / 4 or more for females alcoholic drinks (beer, wine, liquor) at a sitting?

- | | | |
|------------------------------------|------------------------------------|--|
| <input type="radio"/> None | <input type="radio"/> Once | <input type="radio"/> Twice |
| <input type="radio"/> 3 to 5 Times | <input type="radio"/> 6 to 9 Times | <input type="radio"/> 10 or more Times |

14. During the past month (30 days), how did you USUALLY get your alcohol?

- | | |
|---|---|
| <input type="radio"/> I did not drink alcohol during the past month (30 days). | <input type="radio"/> A person 21 years old or older gave it to me. |
| <input type="radio"/> I bought it at a restaurant, bar, or club. | <input type="radio"/> I took it from a store. |
| <input type="radio"/> I bought it at a public event (concert, sporting event). | <input type="radio"/> I got it from a family member. |
| <input type="radio"/> I gave someone else money to buy it for me. | <input type="radio"/> I got it some other way. |
| <input type="radio"/> I bought it at a store (liquor store, convenience store, supermarket, gas station). | |

15. What have been the most important reasons for your drinking alcoholic beverages? (Mark all that apply.)

- | | |
|---|--|
| <input type="radio"/> I did not drink alcoholic beverages | <input type="radio"/> Because of boredom, nothing else to do |
| <input type="radio"/> To experiment (to see what it's like) | <input type="radio"/> Because of anger or frustration |
| <input type="radio"/> To relax or relieve tension | <input type="radio"/> To get through the day |
| <input type="radio"/> To feel good or get high | <input type="radio"/> To increase the effects of some other drug(s) |
| <input type="radio"/> To seek deeper insights and understanding | <input type="radio"/> To decrease (offset) the effects of some other drug(s) |
| <input type="radio"/> To have a good time with my friends | <input type="radio"/> To get to sleep |
| <input type="radio"/> To fit in with a group I like | <input type="radio"/> Because it tastes good |
| <input type="radio"/> To get away from my problems or troubles | <input type="radio"/> Because I am "hooked" (I feel I have to drink) |

CONSEQUENCES OF USE

16. How many times have you experienced the following due to your drinking or drug use during the past year?

| | NEVER | ONCE | TWICE | 3-5 TIMES | 6-10 TIMES | 11 TIMES OR MORE |
|--|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Had a hangover | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Got nauseated or vomited | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Had memory loss | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Performed poorly on a test or project | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Missed school | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Got into a fight or argument | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Damaged property, pulled fire alarms, etc. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

17. Please answer the following questions:

| | YES | NO |
|---|-----------------------|-----------------------|
| Have you ever ridden in a car driven by someone (including yourself) who was high or had been using alcohol or drugs? | <input type="radio"/> | <input type="radio"/> |
| Do you use alcohol or drugs to relax, feel better about yourself, or fit in? | <input type="radio"/> | <input type="radio"/> |
| Do you ever use alcohol or drugs while you are by yourself alone? | <input type="radio"/> | <input type="radio"/> |
| Do you ever forget things you did while using alcohol or drugs? | <input type="radio"/> | <input type="radio"/> |
| Do your family or friends ever tell you that you should cut down on your drinking or drug use? | <input type="radio"/> | <input type="radio"/> |
| Have you ever gotten into trouble while you were using alcohol or drugs? | <input type="radio"/> | <input type="radio"/> |

GAMBLING

5

18. During the last 12 months, how often have you done these activities for money?

| | NOT AT ALL | LESS THAN ONCE A MONTH | ABOUT ONCE A MONTH | ABOUT ONCE A WEEK | DAILY |
|--|-----------------------|------------------------|-----------------------|-----------------------|-----------------------|
| Played cards | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Played games of personal skill like pool, golf, or bowling | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Bet on sports teams | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Played lottery tickets or scratch offs | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Played at a casino | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Played online | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

19. During the last 12 months, have you ever felt....

| | YES | NO | I DON'T BET FOR MONEY |
|--|-----------------------|-----------------------|-----------------------|
| Bad about the amount of money you bet, or about what happens when you bet money? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| That you would like to stop betting money but didn't think you could? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

BELIEFS AND PERCEPTIONS

20. How much do you think people risk harming themselves (physically or in other ways) if they...

| | NO RISK | SLIGHT RISK | MODERATE RISK | GREAT RISK |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| Smoke one or more packs of cigarettes per day | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke marijuana (pot) occasionally | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke marijuana regularly | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Take one or two drinks of alcohol (beer, wine, liquor) occasionally | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Have five or more drinks once or twice a week | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

21. How do you think your close friends feel (or would feel) about you doing each of the following things?

| | STRONGLY APPROVE | APPROVE | DON'T KNOW | DISAPPROVE | STRONGLY DISAPPROVE |
|---|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Smoke one or more packs of cigarettes per day | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke marijuana (pot) occasionally | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke marijuana regularly | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Take one or two drinks of alcohol (beer, wine, liquor) occasionally | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Have five or more drinks once or twice a week | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

22. How wrong do your parents feel it would be for you to...

| | VERY WRONG | WRONG | A LITTLE BIT WRONG | NOT AT ALL WRONG |
|--|-----------------------|-----------------------|-----------------------|-----------------------|
| Drink beer, wine or hard liquor (vodka, whiskey, or gin) regularly (at least once or twice a month)? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke cigarettes? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke marijuana? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Steal something worth more than \$5? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Draw graffiti, or write things or draw pictures on buildings or other property (without the owner's permission)? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Pick a fight with someone? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

23. The next few questions ask about your family and friends.

Mark the Big "YES!" if you think the statement is definitely true for you.
 Mark the little "yes" if you think the statement is mostly true for you.
 Mark the little "no" if you think the statement is mostly not true for you.
 Mark the Big "NO!" if you think the statement is definitely not true for you.

| | YES! | yes | no | NO! |
|--|-----------------------|-----------------------|-----------------------|-----------------------|
| The rules in my family are clear. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| My parents ask if I've gotten my homework done. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| When I am not at home, one of my parents knows where I am and who I am with. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| My family has clear rules about alcohol and drug use. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| We argue about the same things in my family over and over. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| People in my family have serious arguments. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| People in my family often insult or yell at each other. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| My parents ask me what I think before most family decisions affecting me are made. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If I had a personal problem, I could ask my mom or dad for help. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| My parents give me lots of chances to do fun things with them. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Would your parents know if you did not come home on time? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If you drank some beer or wine or hard liquor (for example, vodka, whiskey, or gin) without your parents' permission, would you be caught by your parents? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If you carried a handgun without your parents' permission, would you be caught by your parents? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If you skipped school would you be caught by your parents? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

24. Think of your four best friends (the friends you feel closest to).

In the past year (12 months), how many of your best friends have:

| | NONE OF MY FRIENDS | 1 OF MY FRIENDS | 2 OF MY FRIENDS | 3 OF MY FRIENDS | 4 OF MY FRIENDS |
|---|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Been suspended from school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Carried a handgun? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Sold illegal drugs? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Stolen or tried to steal a motor vehicle such as a car or motorcycle? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Been arrested? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Dropped out of school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Been members of a gang? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Participated in clubs, organizations, or activities at school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Made a commitment to stay drug-free? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Liked school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Regularly attended religious services? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Tried to do well in school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

25. What are the chances you would be seen as cool if you....

| | NO OR VERY LITTLE CHANCE | LITTLE CHANCE | SOME CHANCE | PRETTY GOOD CHANCE | VERY GOOD CHANCE |
|--|--------------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Smoked cigarettes? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Began drinking alcoholic beverages regularly, that is, at least once or twice a month? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoked marijuana? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Carried a handgun? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

PLEASE DO NOT WRITE IN THIS AREA



serial #

26. During the LAST FOUR WEEKS how many whole days of school have you missed because you skipped or "cut" school?

- ☐ None ☐ 1 ☐ 2 ☐ 3 ☐ 4-5 ☐ 6-10 ☐ 11 or more

27. How often do you feel that the schoolwork you are assigned is meaningful and important?

- ☐ Almost always ☐ Often ☐ Sometimes ☐ Seldom ☐ Never

28. How interesting are most of your courses to you?

- ☐ Very interesting and stimulating ☐ Quite interesting ☐ Fairly interesting
☐ Slightly dull ☐ Very dull

29. How important do you think the things you are learning in school are going to be for you later in life?

- ☐ Very important ☐ Quite important ☐ Fairly important
☐ Slightly important ☐ Not at all important

30. Are your school grades better than the grades of most students in your class?

- ☐ YES! ☐ yes ☐ no ☐ NO!

31. Now thinking back over the past year in school, how often did you....

| | NEVER | SELDOM | SOMETIMES | OFTEN | A LOT |
|-------------------------------------|-----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| Enjoy being in school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Hate being in school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Try to do your best work in school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

32. Please answer the following questions:

| | YES! | yes | no | NO! |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| In my school, students have lots of chances to help decide things like class activities and rules. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Teachers ask me to work on special classroom projects. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| My teacher(s) notices when I am doing a good job and lets me know about it. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| There are lots of chances for students in my school to get involved in sports, clubs, and other school activities outside of class. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| There are lots of chances for students in my school to talk with a teacher one-on-one. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| I feel safe at my school. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| The school lets my parents know when I have done something well. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| My teachers praise me when I work hard in school. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| I have lots of chances to be part of class discussions or activities. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Are your school grades better than the grades of most students in your class? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

33. During the past year, did you attend or participate in an "Afternoons R.O.C.K. in Indiana" after-school prevention program, L.E.A.D. Initiative, SADD or other prevention group at your school?

- ☐ Yes ☐ No

INDIVIDUAL PERCEPTIONS

8

34. How wrong do you think it is for SOMEONE YOUR AGE to.....?

| | VERY WRONG | WRONG | A LITTLE BIT WRONG | NOT WRONG AT ALL |
|--|-----------------------|-----------------------|-----------------------|-----------------------|
| Take a handgun to school | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Steal something worth more than \$5 | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Attack someone with the idea of seriously hurting them | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Pick a fight with someone | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Stay away from school all day when their parents think they are at school? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Drink beer, wine or hard liquor (for example vodka, whiskey, or gin) regularly | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke cigarettes | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Smoke marijuana | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| Use LSD, cocaine, amphetamines or another illegal drug | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

35. Please answer the following questions.

| | VERY FALSE | SOMEWHAT FALSE | SOMEWHAT TRUE | VERY TRUE |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| I like to see how much I can get away with. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| I ignore rules that get in my way. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| I do the opposite of what people tell me, just to get them mad. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

COMMUNITY

36. These questions ask about the neighborhood and community where you live.

| | YES! | yes | no | NO! |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| If a kid drank some beer, wine, or hard liquor (for example, vodka, whiskey, or gin) in your neighborhood, would he or she be caught by the police? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If a kid smoked marijuana in your neighborhood, would he or she be caught by the police? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If a kid carried a handgun in your neighborhood, would he or she be caught by the police? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| If a kid smoked a cigarette in your neighborhood, would he or she be caught by the police? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

37. How wrong would most adults (over 21) in your neighborhood think it is for kids your age.....?

| | VERY WRONG | WRONG | A LITTLE BIT WRONG | NOT AT ALL WRONG |
|----------------------|-----------------------|-----------------------|-----------------------|-----------------------|
| To use marijuana? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| To drink alcohol? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| To smoke cigarettes? | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

38. Please answer the following questions.

| | YES! | yes | no | NO! |
|---|-----------------------|-----------------------|-----------------------|-----------------------|
| My neighbors notice when I am doing a good job and let me know. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| There are people in my neighborhood who are proud of me when I do something well. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |
| There are people in my neighborhood who encourage me to do my best. | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> | <input type="radio"/> |

HONESTY

39. How truthfully have you answered these questions?

☐ Not truthfully at all
 ☐ Somewhat truthfully
 ☐ Completely truthfully

PLEASE DO NOT WRITE IN THIS AREA



serial #

THE INDIANA PREVENTION RESOURCE CENTER
2011 PREVALENCE STATISTICS
MAIN FINDINGS

ALCOHOL, TOBACCO AND OTHER DRUG USE BY INDIANA CHILDREN AND ADOLESCENTS

Survey Conducted February through April 2011
Report Dated August 2011

Conducted by:

Indiana Prevention Resource Center

Published by:

Institute for Drug Abuse Prevention

Department of Applied Health Science

School of Health, Physical Education, and Recreation

Indiana University

Funded, in part by a contract with the

Indiana Family and Social Services Administration

Division of Mental Health and Addiction, and the

School of Health, Physical Education, and Recreation

Contact:

Indiana Prevention Resource Center

501 North Morton Street Suite 110

Bloomington, IN 47404

<http://www.drugs.indiana.edu>

drugprc@indiana.edu

Toll Free: 1-800-346-3077

Fax: 812-855-4940



INDIANA UNIVERSITY
SCHOOL OF HEALTH, PHYSICAL EDUCATION,
AND RECREATION
Bloomington